

The source of copper, gold and molybdenum in giant porphyry-type ore deposits from western North America

Other Conference Item**Author(s):**

Pettke, Thomas; Oberli, Felix; [Heinrich, Christoph A.](#) 

Publication date:

2009-06

Permanent link:

<https://doi.org/10.3929/ethz-b-000162197>

Rights / license:

[Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International](#)

Originally published in:

Geochimica et Cosmochimica Acta 73(13 / Supplement), <https://doi.org/10.1016/j.gca.2009.05.012>

The source of copper, gold and molybdenum in giant porphyry-type ore deposits from western North America

THOMAS PETTKE^{1,2*}, FELIX OBERLI²
AND CHRISTOPH A. HEINRICH²

¹University of Bern, Institute of Geological Sciences,
Baltzerstrasse 1+3, CH-3012 Bern, Switzerland
(*correspondence, pettke@geo.unibe.ch)

²ETH Zürich, Department of Earth Sciences, Clausiusstrasse
25, CH-8092 Zürich, Switzerland

Calcalkaline magmas producing porphyry-type ore deposits are thought to be generated by contemporaneous subduction zone processes. We show here that this is not required.

Lead isotope ratios measured in individual fluid inclusions from 37 Ma old high-temperature hydrothermal ore veins at Bingham Canyon, USA, are uniform and conspicuously non-radiogenic, for both the Cu-Au and the distinctly later Mo stage. These data lie at the non-radiogenic end of a linear array described by *in situ* Pb isotope data from feldspars of shallow intrusions bracketing the ore-forming event. It indicates mixing between components from depleted mantle and non-radiogenic, ancient Pb.

Lead isotope forward modeling of the non-radiogenic Pb component reveals a history dating back to the Archean. Three events characterize this evolution. (1) Archean extraction of crustal components, that then evolved with elevated μ - and κ -values in isolation for >1 Ga. (2) During the assembly of Laurentia at ca. 1.8 Ga, this material was subducted and released a Pb-rich but U- and Th-poor fluid component that metasomatized the overlying depleted mantle. This metasomatized mantle then evolved in isolation until incipient extensional tectonics triggered its partial melting in the late Eocene to form the Bingham magmas and associated giant Cu-Mo-Au deposit.

Such distinctly non-radiogenic, lead-rich, ancient source domains are extensive beneath western North America and have been identified at Butte, Henderson, Questa, and at ore deposits in SE Arizona. We propose that Cenozoic melting of lithospheric mantle metasomatized by subduction fluids during the early Proterozoic has provided the metal endowment and subduction flavor for the formation of giant magmatic-hydrothermal ore deposits in western North America, which together constitute the world's major molybdenum province.