SnAP reagents for the synthesis of piperazines and morpholines

Journal Article

Author(s):

Luescher, Michael U.; Vo, Cam-Van T.; Bode, Jeffrey W.

Publication date:

2014-02-21

Permanent link:

<https://doi.org/10.3929/ethz-b-000260433>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Originally published in:

Organic Letters 16(4), <https://doi.org/10.1021/ol500210z>

SnAP Reagents for the Synthesis of Piperazines and Morpholines

Michael U. Luescher, Cam-Van T. Vo, and Jeffrey W. Bode*

Laboratorium für Organische Chemie, ETH Zürich, CH-8093 Zürich, Switzerland

Supporting Information Placeholder

ABSTRACT: Substituted piperazines and morpholines are valuable structural motifs in biologically active compounds, but are not easily prepared by contemporary cross-coupling approaches. In this report, we introduce SnAP reagents for the transformation of aldehydes into *N*-unprotected piperazines and morpholines. This approach offers simple, mild conditions compatible with aromatic, heteroaromatic, aliphatic, and glyoxylic aldehydes and provides mono- and disubstituted *N*-heterocycles in a single step.

We have recently introduced SnAP¹ reagents for the simple synthesis of saturated *N*-heterocycles including thiomorpholines² and medium-sized rings.³ This process allows widely available aliphatic, aromatic, and heteroaromatic aldehydes to be converted to various *N*-heterocycles by a simple, general reaction protocol. It affords directly *N*-unprotected products, has outstanding substrate scope and functional group tolerance, and offers an easily recognized retrosynthetic disconnection. Mechanistically, we current favor a Cu^{II}-promoted generation of a stabilized radical that undergoes endo cyclization with the intermediate imine. We envision that readily prepared SnAP reagents and their equivalents will become commercially available building blocks for direct incorporation of saturated *N*-heterocycles via aldehyde synthetic handles (Figure 1).

In this report, we document SnAP reagents **1–6** for the synthesis of substituted, *N*-unprotected piperazines and morpholines (Figure 2). We also introduce Me-substituted SnAP reagents for the diastereoselective synthesis of disubstituted morpholines and piperazines.

SnAP reagents **1–6** were prepared on a multigram scale by straightforward and efficient routes (see Supporting Information for full synthetic details). The SnAP reagents are easily handled, air- and moisture-stable liquids that can be stored for several weeks without decomposition.⁴ For the purposes of evaluation, a single reaction protocol was used in all of the cyclization reactions.

Piperazines prepared with SnAP Pip **1** were obtained in good yield using electron-poor and electron-rich aromatic, heteroaromatic and aliphatic aldehydes (Table 1). Sterically demanding 2-chloro-4-fluorobenzaldehyde (**7b**) and pivalaldehyde (**7f**) afforded the 6-*endo* products in good yield.

Figure 1. SnAP reagents for the synthesis of *N*-heterocycles from aldehydes.

Products containing groups suitable for further elaboration, including esters, organohalides and a variety of heterocycles, were easily prepared. Additionally, we were pleased to discover that unprotected indole (**7j**) was incorporated in good yield. Piperazines from unbranched aliphatic aldehydes were observed in lower yields, presumably due to the facile enamine formation. The major side products observed in the cyclization reactions were the protodestannylated imines.

Figure 2. SnAP reagents. ^aOverall yield from commercially available starting materials.

Table 1. Piperazine synthesis with SnAP Pip 1^a

^aConditions: SnAP Pip **1** (1.0 equiv, 0.5 mmol), aldehyde (1.0 equiv, 0.5 mmol), MS 4A, CH₂Cl₂ (0.2 M), 2 h, rt; Cu(OTf)₂ (1.0 equiv, 0.5 mmol), 2,6-lutidine (1.0 equiv, 0.5 mmol), 4:1 CH₂Cl₂:HFIP (0.05 M), 12 h, rt; isolated yield.

The synthesis of morpholines using SnAP M **4** showed again a broad substrate scope, functional group tolerance, and afforded the desired *N*-unprotected heterocycles in good to excellent yields from electron-poor and electron-rich aromatic, heteroaromatic and aliphatic aldehydes (Table 2). The reaction with bulky 2,4,6-mesitylaldehyde (**8d**) was slow, promoting the formation of destannylated side product. Elevated temperature did not improve the overall yield; only traces of the desired product were observed. As reported for the synthesis of the thiomorpholines,² 2-pyridinaldehyde (**8k**), which can chelate the Cu^{II}, also showed poor reactivity and the starting material was recovered.

Table 2. Morpholine synthesis with SnAP M 4^a

^aConditions: SnAP M **4** (1.0 equiv, 0.5 mmol), aldehyde (1.0 equiv, 0.5 mmol), MS 4A, CH₂Cl₂ (0.2 M), 2 h, rt; Cu(OTf)₂ (1.0 equiv, 0.5 mmol), 2,6-lutidine (1.0 equiv, 0.5 mmol), 4:1 CH₂Cl₂:HFIP (0.05 M), 12 h, rt; isolated yield. ^b4:1 DCE:HFIP, 12 h, 60 °C; yield determined by ¹H NMR with 1,3,5-trimethoxybenzene as internal standard. ^cAs a mixture of diastereomers.

In general, the formation of the piperazines and morpholines proved more efficient for a broad range of aldehydes and occurred in overall higher yields compared to the thiomorpholine² synthesis. CaSO₄, which we employed in the thiomorpholine synthesis as drying agent, can be left out without deterioration of product formation. We anticipated that substrate specific optimization will be possible if higher yields or faster reaction times are necessary.⁵ An advantage of this protocol for the preparation of diverse *N*-heterocycles is the simple reaction setup: the SnAP reagent is combined with the aldehyde in the presence of molecular sieves to afford the corresponding imine, which is cyclized with stoichiometric Cu(OTf)₂ and 2,6-lutidine in 4:1 CH₂Cl₂:HFIP at rt for 12 h. The imines were processed by filtration over a glass sintered funnel and evaporated to ensure clean and full conversion before subsection to the cyclization. Alternatively, the imine formation can be diluted with additional CH₂Cl₂ and transferred to the heterogeneous copper/ligand suspension by a syringe equipped with an HPLC filter.

The introduction of additional substituents into the targeted piperazine or morpholine can dramatically complicate their synthesis by traditional methods. To examine the feasibility of our approach for the facile diastereoselective synthesis of disubstituted morpholines and piperazines, we prepared methyl substituted SnAP reagents **2–3** and **5–6** via short reaction sequences. Cyclization under standard conditions afforded the representative disubstituted *N*-unprotected saturated heterocycles in moderate to good yields with the usual broad substrate scope (Table 3).

Table 3. Synthesis of disubstituted piperazines and morpholines with SnAP reagents^{a,b,c}

^aConditions: SnAP reagent 2–3, 5–6 (1.0 equiv, 0.5 mmol), aldehyde (1.0 equiv, 0.5 mmol), MS 4A, CH₂Cl₂ (0.2 M), 2 h, rt; Cu(OTf)₂ (1.0 equiv, 0.5 mmol), 2,6-lutidine (1.0 equiv, 0.5 mmol), 4:1 CH₂Cl₂:HFIP (0.05 M), 12 h, rt; isolated yield. ^bDiastereomeric ratio was determined by ¹H NMR spectroscopy of the unpurified reaction mixtures. ^cRelative stereochemistry was confirmed by X-ray analysis of (±)-**9b** and (±)-**11b**; others assigned by analogy and NOESY spectroscopy. ^dYield of major and minor diastereomers combined. ^eYield of major diastereomer. ^fEnantiomer-enriched (*S*)-SnAP 3Me-M **6** (ee > 98%) was used.

The diastereomeric ratio varied depending on the SnAP reagents used. 3,5-disubstituted piperazines (**10a–c**) and morpholines (**12a–b**) bearing substituents in 1,3-relationship generally showed a high diastereomeric ratio towards the thermodynamically favored isomer. SnAP 2Me-M **5** gave predominantly *trans*-configured 3,6-disubstituted morpholines (**11a–b**) (Figure 3a) and SnAP 2Me-Pip **2** afforded the 1,4-diazacyclohexanes (**9a–9c**) as a mixture of separable diastereomers, favoring the *cis*-isomers (Figure 3b). We attribute the reduced diastereoselectivity with SnAP 2Me-Pip **2** to the presence of the *N*-*tert*-butoxycarbonyl group next to the methyl, which is forced into the axial position to reduce steric interactions. No racemization during the synthesis of **12b** was detected using enantiomerically pure SnAP 3Me-M **6**,⁶ permitting fast access to enantiomerically pure building blocks.⁷

Figure 3. X-ray structure of (±)-**11b** and (±)-**9b** and stereochemical assignment by NOESY spectroscopy.

Piperazines and morpholines are important elements in natural products and small bioactive molecules.⁸ Despite the importance of these building blocks, their applications in medicinal chemistry are challenging due to poor commercial availability and the often-laborious synthetic routes that rely on ring closure⁹ or α -functionalization.¹⁰ Successful approaches using vinyl sulfonium salts,¹¹ vinyl selenones,¹² epoxides,¹³ aziridines,¹⁴ or glyoxal derivatives¹⁵ require rather advanced precursors, making the preparation of diverse substitution patterns or heterocycle types laborious. These new SnAP reagents should therefore fulfill a need for the rapid preparation of morpholines and piperazines not currently met by existing methods. We also anticipate that the reagent design and reactions can easily be adapted to accommodate SnAP reagents with additional substituents and functional groups.

In summary, we have developed new SnAP reagents for the synthesis mono- and disubstituted piperazines and morpholines. The cyclization takes place under mild conditions at room temperature and affords the *N*-unprotected, saturated *N*-heterocycles in good to excellent yields. This methodology accepts a wide range of electronically and sterically diverse aromatic, heteroaromatic, glyoxylic, and aliphatic aldehydes bearing variety of functional groups including esters, protected amines, organohalides, ethers, and various heterocycles were tolerated.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures, characterizations, spectral data for all new compounds, and the data for single-crystal X-ray diffraction of compounds (±)-**9b** and (±)-**11b** (CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*Email: bode@org.chem.ethz.ch

Author Contributions

M.U.L. and C. V. V. performed the experiments, compound characterization and data analysis. All authors contributed in experiment design and discussions.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This work was supported by an ETH Research Grant (ETH-12 11-1) and the European Research Council (ERC Starting Grant No. 306793 – CASAA). The authors acknowledge the LOC Mass Spectrometry Service, the LOC NMR Service, and the Small Molecule Crystallography Center of ETH Zurich for the X-ray measurement.

REFERENCES

- (1) SnAP = tin(Sn) Amine Protocol
- (2) Vo, C.-V. T.; Mikutis, G.; Bode, J. W. *Angew. Chem. Int. Ed.* **2013**, *52*, 1705–1708.
- (3) Vo, C.-V. T.; Luescher, M. U.; Bode, J. W. *Nat. Chem. in press*.
- (4) SnAP reagents are usually stored neat at $-10\text{ }^{\circ}\text{C}$ for months without detectable decompositions.
- (5) The formation of **8c**, for example, is completed after 2 h at rt; all reactions are completed after 12 h at rt.
- (6) Enantiomerically pure SnAP 3Me-M **6** was synthesized in one-step from enantiomerically pure (*S*)-(+)-2-amino-1-propanol without detectable racemization.
- (7) See Supporting Information.
- (8) (a) Bruncko, M.; Oost, T. K.; Belli, B. A.; Ding, H.; Joseph, M. K.; Kunzer, A.; Martineau, D.; McClellan, W. J.; Mitten, M.; Ng, S.-C.; Nimmer, P. M.; Oltersdorf, R.; Park, C.-M.; Petros, A. M.; Shoemaker, A. R.; Song, X.; Wang, X.; Wendt, M. D.; Zhang, H.; Fesik, S. W.; Rosenberg, S. H.; Elmore, S. W. *J. Med. Chem.* **2007**, *50*, 641–662. (b) Ward, S. E.; Harrington, F. P.; Gordon, L. J.; Hopley, S. C.; Scott, C. M.; Watson, J. M. *J. Med. Chem.* **2005**, *48*, 3478–3480. (c) Leopoldo, M.; Lacivita, E.; Colabufo, N. A.; Contino, M.; Berardi, F.; Perrone, R. *J. Med. Chem.* **2005**, *48*, 7919–7922. (d) Wijtmans, R.; Vink, M. K. S.; Schoemaker, H. E.; van Delft, F. L.; Blaauw, R. H.; Rutjes, F. P. J. T. *Synthesis* **2004**, 641–662. For natural products, see: (e) Lin, C.; Huang, P.; Lu, C.; Wu, R.; Hu, W.; Wang, J. *Tetrahedron* **1997**, *53*, 2025–2028. (f) Fujita, T.; Hayashi, H. *Biosci. Biotechnol. Biochem.* **2004**, *68*, 820–826. (g) Kohmoto, S.; Kashman, Y.; McConnell, O. J., Jr.; Rinehart, K. L., Jr.; Wright, A.; Koehn, F. J. *J. Org. Chem.* **1988**, *53*, 3116–3118.
- (9) For some recent examples, see: (a) Ruider, S. A.; Müller, S.; Carreira, E. M. *Angew. Chem. Int. Ed.* **2013**, *52*, 11908–11911. (b) Zhai, H.; Borzenko, A.; Lau, Y. Y.; Ahn, S. H.; Schafer, L. L. *Angew. Chem. Int. Ed.* **2012**, *51*, 12219–12223. (c) Lu, Z.; Stahl, S. S. *Org. Lett.* **2012**, *14*, 1234–1237. (d) Leathen, M. L.; Rosen, B. R.; Wolfe, J. P. *J. Org. Chem.* **2009**, *74*, 5107–5110. (e) Cochran, B. M.; Michael, F. E. *Org. Lett.* **2008**, *10*, 329–332. (f) Nakhla, J. S.; Wolfe, J. P. *Org. Lett.* **2007**, *9*, 3279–3282. (g) Lanman, B. A.; Myers, A. G. *Org. Lett.* **2004**, *6*, 1045–1047.
- (10) For reviews, see: (a) Mitchell, E. A.; Peschiulli, A.; Lefevre, N.; Meerpoel, L.; Maes, B. U. W. *Chem. Eur. J.* **2012**, *18*, 10092–10142. (b) Campos, K. R. *Chem. Soc. Rev.* **2007**, *36*, 1069–1084.
- (11) (a) Yar, M.; McGarrigle, E. M.; Aggarwal, V. K. *Angew. Chem. Int. Ed.* **2008**, *47*, 3784–3786. (b) Yar, M.; McGarrigle, E. M.; Aggarwal, V. K. *Org. Lett.* **2009**, *11*, 257–260. (c) Yar, M.; Fritz, S. P.; Gates, P. J.; McGarrigle, E. M.; Aggarwal, V. K. *Eur. J. Org. Chem.* **2012**, 160–166.
- (12) Bagnoli, L.; Scarponi, C.; Rossi, M. G.; Testaferri, L.; Tiecco, M. *Chem. Eur. J.* **2011**, *17*, 993–999.
- (13) Breuning, M.; Winnacker, M.; Steiner, M. *Eur. J. Org. Chem.* **2007**, 2100–2106.
- (14) Wang, L.; Liu, Q.-B.; Wang, D.-S.; Li, X.; Han, X.-W.; Xiao, W.-J.; Zhou, Y.-G. *Org. Lett.* **2009**, *11*, 1119–1122.
- (15) Berrée, F.; Debache, A.; Marsac, Y.; Carboni, B. *Tetrahedron Lett.* **2001**, *42*, 3591–3594.

-
- (1) SnAP = tin(Sn) Amine Protocol
- (2) Vo, C.-V. T.; Mikutis, G.; Bode, J. W. *Angew. Chem. Int. Ed.* **2013**, *52*, 1705–1708.
- (3) Vo, C.-V. T.; Luescher, M. U.; Bode, J. W. *Nat. Chem.* *in press*.
- (4) SnAP reagents are usually stored neat at –10 °C for months without detectable decompositions.
- (5) The formation of **8c**, for example, is completed after 2 h at rt; all reactions are completed after 12 h at rt.
- (6) Enantiomerically pure SnAP 3Me-M **6** was synthesized in one-step from enantiomerically pure (S)-(+)-2-amino-1-propanol without detectable racemization.
- (7) See Supporting Information.
- (8) (a) Bruncko, M.; Oost, T. K.; Belli, B. A.; Ding, H.; Joseph, M. K.; Kunzer, A.; Martineau, D.; McClellan, W. J.; Mitten, M.; Ng, S.-C.; Nimmer, P. M.; Oltersdorf, R.; Park, C.-M.; Petros, A. M.; Shoemaker, A. R.; Song, X.; Wang, X.; Wendt, M. D.; Zhang, H.; Fesik, S. W.; Rosenberg, S. H.; Elmore, S. W. *J. Med. Chem.* **2007**, *50*, 641–662. (b) Ward, S. E.; Harrington, F. P.; Gordon, L. J.; Hopley, S. C.; Scott, C. M.; Watson, J. M. *J. Med. Chem.* **2005**, *48*, 3478–3480. (c) Leopoldo, M.; Lacivita, E.; Colabufo, N. A.; Contino, M.; Berardi, F.; Perrone, R. *J. Med. Chem.* **2005**, *48*, 7919–7922. (d) Wijtmans, R.; Vink, M. K. S.; Schoemaker, H. E.; van Delft, F. L.; Blaauw, R. H.; Rutjes, F. P. J. T. *Synthesis* **2004**, 641–662. For natural products see: (e) Lin, C.; Huang, P.; Lu, C.; Wu, R.; Hu, W.; Wang, J. *Tetrahedron* **1997**, *53*, 2025–2028. (f) Fujita, T.; Hayashi, H. *Biosci. Biotechnol. Biochem.* **2004**, *68*, 820–826. (g) Kohmoto, S.; Kashman, Y.; McConnell, O. J., Jr.; Rinehart, K. L., Jr.; Wright, A.; Koehn, F. J. *Org. Chem.* **1988**, *53*, 3116–3118.
- (9) For some recent examples see: (a) Ruider, S. A.; Müller, S.; Carreira, E. M. *Angew. Chem. Int. Ed.* **2013**, *52*, 11908–11911. (b) Zhai, H.; Borzenko, A.; Lau, Y. Y.; Ahn, S. H.; Schafer, L. L. *Angew. Chem. Int. Ed.* **2012**, *51*, 12219–12223. (c) Lu, Z.; Stahl, S. S. *Org. Lett.* **2012**, *14*, 1234–1237. (d) Leathen, M. L.; Rosen, B. R.; Wolfe, J. P. *J. Org. Chem.* **2009**, *74*, 5107–5110. (e) Cochran, B. M.; Michael, F. E. *Org. Lett.* **2008**, *10*, 329–332. (f) Nakhla, J. S.; Wolfe, J. P. *Org. Lett.* **2007**, *9*, 3279–3282. (g) Lanman, B. A.; Myers, A. G. *Org. Lett.* **2004**, *6*, 1045–1047.
- (10) For reviews see: (a) Mitchell, E. A.; Peschiulli, A.; Lefevre, N.; Meerpoel, L.; Maes, B. U. W. *Chem. Eur. J.* **2012**, *18*, 10092–10142. (b) Campos, K. R. *Chem. Soc. Rev.* **2007**, *36*, 1069–1084.
- (11) (a) Yar, M.; McGarrigle, E. M.; Aggarwal, V. K. *Angew. Chem. Int. Ed.* **2008**, *47*, 3784–3786. (b) Yar, M.; McGarrigle, E. M.; Aggarwal, V. K. *Org. Lett.* **2009**, *11*, 257–260. (c) Yar, M.; Fritz, S. P.; Gates, P. J.; McGarrigle, E. M.; Aggarwal, V. K. *Eur. J. Org. Chem.* **2012**, 160–166.
- (12) Bagnoli, L.; Scarponi, C.; Rossi, M. G.; Testaferri, L.; Tiecco, M. *Chem. Eur. J.* **2011**, *17*, 993–999.
- (13) Breuning, M.; Winnacker, M.; Steiner, M. *Eur. J. Org. Chem.* **2007**, 2100–2106.
- (14) Wang, L.; Liu, Q.-B.; Wang, D.-S.; Li, X.; Han, X.-W.; Xiao, W.-J.; Zhou, Y.-G. *Org. Lett.* **2009**, *11*, 1119–1122.
- (15) Berrée, F.; Debache, A.; Marsac, Y.; Carboni, B. *Tetrahedron Lett.* **2001**, *42*, 3591–3594.