The receptor locus for Escherichia coli F4ab/F4ac in the pig maps distal to the MUC4-LMLN region

Journal Article**Author(s):**

Rampoldi, Antonio; Jacobsen, Mette J.; Bertschinger, Hans U.; Joller, David; Bürgi, Ester; Vögeli, Peter; Andersson, Leif; Archibald, Alan L.; Fredholm, Merete; Jørgensen, Claus B.; Neuenschwander, Stefan

Publication date:

2011-02

Permanent link:

<https://doi.org/10.3929/ethz-b-000026882>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Originally published in:

Mammalian Genome 22(1-2), <https://doi.org/10.1007/s00335-010-9305-3>

The receptor locus for *Escherichia coli* F4ab/F4ac in the pig maps distal to the *MUC4*–*LMLN* region

Antonio Rampoldi · Mette J. Jacobsen · Hans U. Bertschinger · David Joller · Esther Bürgi · Peter Vögeli · Leif Andersson · Alan L. Archibald · Merete Fredholm · Claus B. Jørgensen · Stefan Neuenschwander

Received: 2 July 2010 / Accepted: 3 November 2010 / Published online: 7 December 2010
© Springer Science+Business Media, LLC 2010

Abstract Enterotoxigenic *Escherichia coli* (ETEC) with fimbriae of the F4 family are one of the major causes of diarrhea and death among neonatal and young piglets. Bacteria use the F4 fimbriae to adhere to specific receptors expressed on the surface of the enterocytes. F4 fimbriae exist in three different antigenic variants, F4ab, F4ac, and F4ad, of which F4ac is the most common. Resistance to ETEC F4ab/F4ac adhesion in pigs has been shown to be inherited as an autosomal recessive trait. In previous studies the ETEC F4ab/F4ac receptor locus (*F4bcR*) was mapped to the q41 region on pig chromosome 13. A polymorphism within an intron of the mucin 4 (*MUC4*) gene, which is one of the possible candidate genes located in this region, was shown earlier to cosegregate with the *F4bcR* alleles. Recently, we

discovered a Large White boar from a Swiss experimental herd with a recombination between *F4bcR* and *MUC4*. A three-generation pedigree including 45 offspring was generated with the aim to use this recombination event to refine the localization of the *F4bcR* locus. All pigs were phenotyped using the microscopic adhesion test and genotyped for a total of 59 markers. The recombination event was mapped to a 220-kb region between a newly detected SNP in the leishmanolysin-like gene (*LMLN* g.15920) and SNP *ALGA0072075*. In this study the six SNPs *ALGA0072075*, *ALGA0106330*, *MUC13-226*, *MUC13-813*, *DIA0000584*, and *MARC0006918* were in complete linkage disequilibrium with *F4bcR*. Based on this finding and earlier investigations, we suggest that the locus for *F4bcR* is located between the *LMLN* locus and microsatellite S0283.

A. Rampoldi and M. J. Jacobsen contributed equally to this work.

A. Rampoldi · H. U. Bertschinger · D. Joller · P. Vögeli (✉) · S. Neuenschwander
Institute of Plant, Animal and Agroecosystem Sciences,
ETH Zurich, 8092 Zurich, Switzerland
e-mail: peter.voegeli@inw.agr.ethz.ch

M. J. Jacobsen · M. Fredholm · C. B. Jørgensen
Basic Animal and Veterinary Sciences, University
of Copenhagen, 1870 Frederiksberg, Copenhagen, Denmark

E. Bürgi
Department of Farm Animals, University of Zurich,
8057 Zurich, Switzerland

L. Andersson
Department of Medical Biochemistry and Microbiology,
Uppsala University, 751 23 Uppsala, Sweden

A. L. Archibald
Division of Genetics and Genomics, The Roslin Institute
and R(D)SVS, University of Edinburgh, Roslin,
Midlothian EH25 9PS, UK

Introduction

Close association to host surface is an essential requirement for successful induction of disease by a broad variety of pathogens. This association is often mediated by lectin recognition of host cell glycoconjugates (Karlsson 1998). Breeding for genetic resistance is an established routine in plant sciences and has become an attractive goal in animal sciences. Thus, selection for absence of receptor molecules or for genetic variants of receptor molecules to which the pathogen cannot bind would represent an attractive route to genetic resistance.

The two most economically dominant porcine enteric *E. coli* infections comprise diarrhea induced by diverse enterotoxins and enterotoxemia mediated by a shigatoxin. Both diseases have in common that the bacteria have to colonize the small intestine in high numbers where they produce the deleterious toxins. To build up huge

populations, the bacteria are forced by the short transit time of ingesta to adhere to the small intestine's inner surface. This adhesion is the result of a highly specific interaction between fimbriae produced by the bacteria and receptors present exclusively on the brush border of small-intestine enterocytes (Bertschinger and Fairbrother 1999). Of the five pig-specific types of fimbriae, designated F4, F5, F6, F18, and F41 (Osek 1999), so far it is known that only two, F18 and F4, are highly prevalent. Antigen variants have been detected for both types of fimbriae. Whereas the two variants F18ab and F18ac recognize the same receptor (Rippinger et al. 1995), the three variants of F4 vary in their receptor specificities (Bijlsma et al. 1982). However, F4ac is by far the most prevalent variant all over the world with the exception of central China, where F4ad is reported to be predominant (Wang et al. 2006). Fortunately, the causative mutation for *E. coli* F18 susceptibility is known (Meijerink et al. 1997, 2000) and elimination of the *E. coli* F18 susceptibility allele from the porcine population is currently done in Switzerland and other countries to increase the number of resistant piglets in commercial populations (Luther et al. 2009).

In pigs the resistant and susceptible phenotypes for adherence or infection by *E. coli* with F4ac fimbriae are inherited as a monogenetic trait, the susceptibility allele being dominant over the resistance one (Gibbons et al. 1977). The receptor for F4ac or a closely linked locus binds F4ab as well (Jørgensen et al. 2003; Python et al. 2002). As yet, attempts to develop practical methods for identification of breeding stock that inherit resistance have not been completely successful. Several investigators have mapped the ETEC F4ab/F4ac receptor gene (*F4bcR*) to SSC13 (Edfors-Lilja et al. 1995; Python et al. 2002). An overview of the marker positions and mapped genes encompassing the *F4bcR* is given in Fig. 1. Subsequent studies mapped the ETEC F4ab/F4ac susceptibility locus to the interval *SW207-S0075* refined in a 5.7-cM region around the *F4bcR* locus (Joller et al. 2009). Jørgensen et al. (2004) reported an association between resistance and a mutation in intron 7 of the mucin 4 (*MUC4* g.8227 G > C) gene. However, data presented later by Rasschaert et al. (2007) as well as observations in Switzerland (Joller 2009) raised doubt as to whether this mutation in the *MUC4* gene is in some rare cases not in complete linkage disequilibrium (LD) with the F4ac receptor locus.

Fig. 1 Location of 59 markers in the *F4bcR* candidate region at SSC13. The gene order and the scale are deduced from the Sscrofa9 assembly. The positions of the microsatellites are shown to the right of the chromosome 13 idiogram. The approximate positions of candidate genes are indicated at the far right side. The number of markers in each gene is given in parenthesis. The position of the six SNPs *ALGA0072075*, *ALGA0106330*, *MUC13-226*, *MUC13-813*, *DIA0000584*, and *MARC000691*, which were in complete LD, are shown in bold

breeding. Samples of blood and intestine were taken at that time. The interval between killing and sampling of intestine was between 10 and 30 min to avoid degeneration of the epithelial intestinal cells.

All animals were phenotyped with the antigenic variant F4ac fimbriae by the microscopic adhesion test according to the method of Vögeli et al. (1996). A more detailed description of the preparation of bacterial strains, sampling of intestinal tissues, purification of enterocytes, and the procedure of the microscopic adhesion test can be found in Joller (2009). In brief, starting at the upper mesentery root, the intestine was separated from the mesenterium and an empty segment in the middle part was taken anywhere between 3.5 and 7.5 m in young pigs and between 5 and 10 m in older pigs. Twenty intact enterocytes were scored under an optical microscope by the same person. A pig was classified as adhesive to F4ac if more than 15% of the enterocytes bound more than five bacteria on the brush borders.

Screening for genetic polymorphisms

Genomic DNA was extracted from EDTA-anticoagulated whole blood as described by Vögeli et al. (1994). A total of 59 markers on chromosome 13q41 were used to characterize the three-generation pedigree (53 SNPs listed in Tables 1 and 2) along with 6 microsatellite markers: *SW207*, *MUC4GT*, *S0283*, *S0075*, *SW1876*, and *SW698*. Twenty-nine markers from Table 1 were selected from Jacobsen et al. (2010), nine SNPs in *MUC13* were taken from Zhang et al. (2008), and *LMLN* g.15920 was detected in this study by BLASTing the human sequence (NM_001136049) to the pig genome and by sequencing the genomic porcine homolog. The 14 SNPs listed in Table 2 were analyzed using the Illumina PorcineSNP60 BeadChip (Illumina, Inc., San Diego, CA, USA) (Ramos et al. 2009). The microsatellite markers were selected from Python et al. (2002).

PCR was carried out in a reaction volume of 25 µl containing 100 ng of genomic DNA, 10 mM Tris-HCl (pH 9), 50 mM KCl, 1.5 mM MgCl₂, 200 µM of each deoxynucleotide, 0.4 µM of forward and reverse primers, and 1.25 U of *Taq* DNA polymerase. The annealing temperature was chosen according to the melting temperature of the primers. For microsatellites, the PCR consisted of 25 cycles and for the SNPs 35 cycles followed by a final extension at 72°C for 2 min. Thermal cycling conditions were incubation at 94°C for 2 min followed by incubation at the specific annealing temperature for 30 sec and extension at 72°C for 30 sec. The subsequent denaturation was at 94°C for 30 sec. DNA samples were then purified with Montage PCR Centrifugal Filter Devices (p36461) (Millipore, Billerica, MA, USA) and sequenced. Alleles of the microsatellites were analyzed on an ABI377 DNA

analyzer (Applied Biosystems, Foster City, CA, USA). Haplotypes were determined using MERLIN (Abecasis et al. 2002) and HaploPainter software (Thiele and Nürnberg 2005).

Results

The distribution of the 59 markers in the candidate region of the *F4bcR* locus is shown in Fig. 1, and the approximate positions of genes and markers are according to the Sscrofa9 assembly and the physical map (http://pre.ensembl.org/Sus_scrofa_map/). These markers cover the 35-Mb interval from microsatellite SW207 to SW698. Fifty-five markers map to the core interval 100–104 Mb on chromosome 13.

Boar B698 was mated with three sows resistant to ETEC F4ab/F4ac and hence putative homozygotes for the absence of an adherent ETEC F4ab/F4ac receptor, generating a total of 45 offspring. All of the offspring were adhesive to ETEC F4ab/F4ac according to the microscopic adhesion test. Thus, all the offspring are presumed to have received (from boar B698) a susceptibility allele encoding the presence of a receptor to which *E. coli* F4ac can adhere. Therefore, it can be deduced that boar B698 is homozygous for the susceptibility allele. However, the genotyping data showed that 23 piglets (51.1%) possessed the *MUC4* g.8227 C/C genotype, which was previously reported to be associated with the absence of F4ab/F4ac adhesion (Jacobsen et al. 2010; Jørgensen et al. 2004), and 22 piglets were C/G (48.9%). Haplotype analysis of the genotype data reveals a putative recombination event between *LMLM* g.15920 and *ALGA0072075* SNP marker loci (Fig. 2). The recombinant chromosome was transmitted to 23 of B698's offspring. The recombinant chromosome was identical to all other susceptible haplotypes for the interval *ALGA0072075*–*SW698*. The *LMLN*–*S0075* region was also covered with 11 SNPs of the PorcineSNP60 BeadChip (Table 2), 2 microsatellites, and 14 SNPs in *MUC13*. In contrast to the *MUC4* markers, *ALGA0072075*, *ALGA0106330*, *MUC13*-226, *MUC13*-813, *DIA0000584*, and *MARC0006918* were all in complete LD with *F4bcR* in our family (Fig. 2).

Discussion

Previous studies identified *MUC4* as the most probable candidate gene for *F4bcR* based on its chromosomal location and evidence that the molecule to which the *E. coli* F4ac adheres is a mucin-like sialoglycoprotein (Erickson et al. 1994; Jacobsen et al. 2010; Joller et al. 2009; Jørgensen et al. 2004). As reported earlier, the resistance

Table 1 Description of the SNPs

BAC Clone	GenBank reference	SNP name	Alleles ^a <i>R</i> > <i>S</i>	Position on SSC13 (Sscrofa9)	Forward primer	Reverse primer	Amplicon length
CH242 182C12	NM_001039617	<i>ZDHH19</i> g.4043 ^b	C>T	100681071	TCCTACGAGGGCAAGGTATGTG	GAACTGAATCTGGTCTGGAATAGC	171
CH242 221N4	NM_001010983	<i>TNK2</i> g.7075 ^b	C>A	100914782	GGAAGCCACTTTGATTGTTCTC	GACAGGGACACCAACAGCTAA	625
		<i>TNK2</i> g.7717 ^b	C>T	100915124	GAAGAGCTGGTGTCTCTGCTT	AGCGGACGTGCATACCTTGA	551
		<i>TNK2</i> g.11142 ^b	G>A	100924258	CCTGTGAGTGAAGACCAAGACC	CCCCATCTCCTCCTCACTTGA	751
CH242 240L11	NM_004532	<i>MUC4</i> g.6242 ^b	G>A	101034210	GTTACTGGCCTCGACTCTCC	AGGTTGTACCCCTTGGCAATTC	749
		<i>MUC4</i> g.6308 ^b	G>T	101034276			
		<i>MUC4</i> g.6317 ^b	G>A	101034285			
		<i>MUC4</i> g.6321 ^b	G>C	101034289			
		<i>MUC4</i> g.6609 ^b	T>A	101034577			
		<i>MUC4</i> g.6616 ^b	G>T	101034584			
		<i>MUC4</i> g.6634 ^b	A>C	101034602			
		<i>MUC4</i> g.6690 ^b	A>T	101034658			
		<i>MUC4</i> g.6745 ^b	T>C	101034713			
		<i>MUC4</i> g.6770 ^b	G>T	101034738			
		<i>MUC4</i> g.7947 ^b	A>G	101035915	GGTCCTACGCCCTTGTCTC	CCTTCATGGGGTGTGTGAATA	462
		<i>MUC4</i> g.8227 ^b	C>G	101036195	GTGCCCTTGGGTGAGAGGTGA	CACTCTGCCGTTCTCTTCC	367
CH242 123C16	NM_001145642	<i>KIAA0226</i> g.62250 ^b	G>A	101112249	GCGGGAGCTGCTAGCCAA	GGGCTTCGGTGGGTTCCCT	70
	NM_032773	<i>LRCH3-212</i> ^b	G>C	101264808	TGCCTGACATTTTGCTAACG	CTGCACCTTGTGGTGGAGAAC	643
		<i>LRCH3-213</i> ^b	G>C	101264855			
		<i>LRCH3-214</i> ^b	T>C	101264894			
		<i>LRCH3-215</i> ^b	T>A	101265038			
		<i>LRCH3-217</i> ^b	G>A	101270027	TTGAGGAGAGTTGCATGTGT	TCCTGCTCAGTGGATTAAAGG	405
		<i>LRCH3-218</i> ^b	A>G	101270038			
		<i>LRCH3-219</i> ^b	G>T	101270040			
CH242 101B19	NM_001136049	<i>MLN</i> g.15920 ^d	C>G	101361940	GGCACTATCTTACTTAGCAG	TGGTTTGTGACACATTGT	528
	NM_033049.2	<i>MUC13-227</i> ^b	C>T	101846954	TGAGCAAGATGAGTGCCCCAGT	TAGCCAGGCAGGCACAAGCA	536
		<i>MUC13-226</i> ^b	G>A	101847134			
		<i>MUC13-225</i> ^b	G>C	101847197			
		<i>MUC13-224</i> ^b	A>G	101847222			
		<i>MUC13-223</i> ^b	T>C	101847300			

Table 1 continued

BAC Clone	GenBank reference	SNP name	Alleles ^a <i>R</i> > <i>S</i>	Position on SSC13 (Sscrofa9)	Forward primer	Reverse primer	Amplicon length
		<i>MUC13</i> -813 ^c	C>T	101853611	ATGTGGAAGAACAGAACTTGATTGAG	ATAGTCAGGGCGGGGTATACTACC	176
		<i>MUC13</i> -814 ^c	T>G	101853612			
		<i>MUC13</i> -829 ^c	C>A	101853627			
		<i>MUC13</i> -895 ^c	C>A	101853693			
		<i>MUC13</i> -905 ^c	A>G	101853703			
		<i>MUC13</i> -908 ^c	A>G	101853706			
		<i>MUC13</i> -920 ^c	G>A	101853718			
		<i>MUC13</i> -933 ^c	T>C	101853731			
		<i>MUC13</i> -935 ^c	C>A	101853733			

The positions of the SNPs are given according to their position in the reference sequence (second column)

^a *R* alleles are associated with the resistant and *S* alleles with the susceptible haplotypes, respectively

^b SNPs from Jacobsen et al. (2010); ^c SNPs from Zhang et al. (2008); ^d SNP from this study

Table 2 SNP list from the PorcineSNP60 BeadChip

SNP name	Position on SSC13 (Sscrofa9)	Sequence and allelic variants ^a
<i>MAR0067282</i>	101488856	AGAAAAAGAACTCTATTGAATGACGGT [C/T] TCAGAGTATACCTGTTCCTGTTCACCTA
<i>ALGA0072075</i>	101582070	CAGCTTGGAAAGCCATGCGTTGGAGAAATGG [C/T] AGAGCCACTCAGAGTATGGAGACCTGAGTG
<i>ASGA0058925</i>	101659492	TGTGTCTGTCTCTCCCGATGGGGCTCTTC [C/T] TCTACAGGAGAGCTGACTGCAGCATGTTGG
<i>ALGA0122555</i>	101783014	CTTGAAGAAAGCTAATTGGGCAACACTGCT [C/T] GAAGCCTTAATGTCTTTACTGTGTAATTAG
<i>ASGA0091537</i>	101818006	GATAGGAAAATAAAGTATGATCAACTTTATTTA [C/T] TATCCAAAATAAGATATTGATAATTAGCTTT
<i>ALGA0106330</i>	101846502	TAAACACAGACTTTAAGTACAAATGGGAATTC [A/G] GGGGCTCAGCTACCTAAAGGTAACATTGTG
<i>ALGA0106230</i>	101860064	CATGTCACATATATCTCAATAAAGCTGTG [G/T] TTTTITTTAAGAGGAAGGTGGCTGCTCATG
<i>H3GA0037348</i>	101925778	GGAGGCCACCCAGGTATGGAGAGCATTTTCG [C/A] TGCAAGCCTTTCTAAGTGGCTCTGGCTCA
<i>H3GA0037371</i>	102371992	CAAAAGCAGGCTGCTTTTCAAGTTAAAGTCTG [G/A] GACCAATACATGGTAAATAAAAAGTAAGAA
<i>DIAS0000584</i>	102430792	CCCAGAAACCCACCCAGCAGATGCTCTTCT [G/C] CAGTGTCTTAGATCACACCCCGCTGGAC
<i>MARC0006918</i>	102797817	CATCCAGAAATTGCTGACCCAGCATGCTGA [G/A] GTGATTTGTGCAGCCTCTTCTCCCTGTTTT
<i>MARC0045417</i>	103289127	CCTTACCTCCTTCAAGCTTATGTTGGAATC [A/G] AGAATAATCTGCCCTGAGAAAGCAATAAAA
<i>H3GA0037402</i>	103303135	GGCCTTGTTTCACTACCTTTGAAAATAATTC [C/T] ATCTTCCTTTCAATACGTAAGAGCAGTAAA
<i>ASGA0059048</i>	107026948	TCTAGGCTCATGTTGTTTGAAGGGAAA [G/A] CAGAGAGCTGGCTCCTTTGTGGAAGCTTT

^a The SNP is given in bold and in square brackets. The first nucleotide of the SNP indicates the resistant and the second the susceptible allele

allele is presumed to represent the ancestral allele and is associated with greater haplotype heterogeneity (see 798B and 739B, Fig. 2) (Jacobsen et al. 2010). The large shared haplotype block associated with the susceptibility allele handicaps efforts to locate and identify the causative mutation. In addition, determination of the *F4bcR* genotypes is time-consuming as it has to be done by progeny phenotyping which leads to the loss of potential breeding animals. Thus, when we detected a boar with a recombination between the *MUC4* markers and *F4bcR*, it represented an opportunity to refine the search for the causative mutation. Therefore, we mated this boar to homozygous resistant sows and analyzed more markers distal to *MUC4*.

Boar 698B inherited a recombinant haplotype from its father, boar 381B. The recombinant allele was transmitted from 698B to half of its progeny, as seen in pigs 1977 and 2734 (Fig. 2), and was not found in other pigs from the same herd. The recombination in 698B occurred in an interval of about 220 kb, between SNPs *LMLN* g.15920 and *ALGA0072075*. Boars 381B and 698B were found to be homozygous for the SNP *MARC0067282*, making it impossible to determine the position of the crossing over in 698B more precisely.

These results support the hypothesis that the locus for *F4bcR* is located distal to *MUC4*. Besides the experimental pigs, we phenotyped 78 unrelated pigs of 38 litters of the Swiss performing station and determined their *MUC4* g.8227 genotype (Joller 2009). Five pigs were phenotyped as resistant, whereas the genotype was *MUC4* g.8227 C/G and one susceptible pig was *MUC4* g.8227 C/C. These pigs were typed with the markers used in this study and the resistant pigs were homozygous A/A in *ALGA0106330*, G/G in *MUC13-226*, and C/C in *MUC13-813*, consistent with a resistant genotype. The susceptible pig was heterozygous A/G, A/G, and C/T, respectively, for the same markers. These results suggest stronger LD between these markers and *F4bcR* than between *MUC4* g.8227 and *F4bcR* and give further evidence that *F4bcR* is located distal to *MUC4*.

BLAST analyses of the markers that were in complete LD with *F4bcR* showed that only *MUC13-813* was located in an exon. However, this C > T transition is silent. *ALGA0072075* was mapped between *SLC12A8* and *HEG1*. *ALGA0106330* was located in an intron of *MUC13*. *DIAS0000584* was mapped to an intron of gene *KALRN* and *MARC0006918* to an intron of *MYLK*. These findings indicate that these markers may not be causative for ETEC F4ab/F4ac adhesion.

Joller et al. (2006) had initially mapped the most probable position of the *F4bcR* locus in the *SW207–S0283* interval. Unfortunately, the family in this study did not allow the identification of a distal border closer to the unknown *F4bcR* than microsatellite *S0283*. Therefore, we

propose the interval between *LMLN* g.15920 and *S0283*, consisting of around 620 kb, to be the most probable region for *F4bcR*.

Mucin-type proteins and lipids are present on almost every epithelial tissue and are important for defense against pathogens. However, mucins and other glycosylated structures are also targets for microbial attachment as they contain a variety of oligosaccharide structures providing binding sites for bacteria. In the case of bacteria expressing the F18-type fimbriae, the adhesion is dependent on the activity of fucosyltransferase 1 and the presence of a yet unknown fucosylated structure (Meijerink et al. 2000). Although the causative mutation in F4ac is still unknown, the genetic analyses performed in 698B's family have determined that the causal mutation for ETEC F4ab/F4ac susceptibility is downstream of the gene *LMLN* and probably may be located around the region containing the *MUC13* gene.

Acknowledgments We thank Prof. Martien Groenen and his group (Animal Breeding and Genetics, Wageningen University, Wageningen, The Netherlands) for providing the SNP sequences and their positions in the Sscrofa9 assembly. We thank Michael Goe for editing the manuscript and the Genomic Diversity Center (ETH Zurich) for allowing us to use their infrastructure. The study was supported by the Swiss National Science Foundation (grant No. 3100A0-120255/1), by the ETH Zurich, and the SUISAG, Sempach (Dr. Andreas Hofer).

References

- Abecasis GR, Cherny SS, Cookson WO, Cardon LR (2002) Merlin—rapid analysis of dense genetic maps using sparse gene flow trees. *Nat Genet* 30:97–101
- Bertschinger HU, Fairbrother JM (1999) *Escherichia coli* infections. In: Straw BE, D'Allaire S, Mengeling WL, Taylor DJ (eds) *Diseases of Swine*, 8th edn. Iowa State University Press, Ames, IA, pp 431–468
- Bijlsma IG, de Nijs A, van der Meer C, Frik JF (1982) Different pig phenotypes affect adherence of *Escherichia coli* to jejunal brush borders by K88ab, K88ac, or K88ad antigen. *Infect Immun* 37:891–894
- Edfors-Lilja I, Gustafsson U, Duval-Iflah Y, Ellergren H, Johansson M et al (1995) The porcine intestinal receptor for *Escherichia coli* K88ab, K88ac: regional localization on chromosome 13 and influence of IgG response to the K88 antigen. *Anim Genet* 26:237–242
- Erickson AK, Baker DR, Bosworth BT, Casey TA, Benfield DA et al (1994) Characterization of porcine intestinal receptors for the K88ac fimbrial adhesion of *Escherichia coli* as mucin-type sialoglycoproteins. *Infect Immun* 62:5404–5410
- Gibbons RA, Sellwood R, Burrows M, Hunter PA (1977) Inheritance of resistance to neonatal *E. coli* diarrhoea in the pig: Examination of the genetic system. *Theor Appl Genet* 51:65–70
- Jacobsen M, Kracht SS, Estes G, Cirera S, Edfors I et al (2010) Refined candidate region specified by haplotype sharing for *Escherichia coli* F4ab/F4ac susceptibility alleles in pigs. *Anim Genet* 41:21–25
- Joller D (2009) Comparative molecular approaches to identify host determinants mediating adhesion of *E. coli* F4 strains in pigs.

- Dissertation ETH Zürich number 18518. Available at <http://e-collection.ethbib.ethz.ch/eserv/eth:445/eth-445-02.pdf#search=%22Joller%20David%22>
- Joller D, Jørgensen CB, Bertschinger HU, Bürgi E, Stannarius C et al (2006) Refined linkage mapping of the *Escherichia coli* F4ac receptor gene on pig chromosome 13. Proceedings of the 30th International Conference for Animal Genetics, Porto Seguro, Brazil, August 20–25, 2006
- Joller D, Jørgensen CB, Bertschinger HU, Python P, Edfors I et al (2009) Refined localization of the *Escherichia coli* F4ab/F4ac receptor locus on pig chromosome 13. *Anim Genet* 40:749–752
- Jørgensen CB, Cirera S, Archibald AL, Anderson L, Fredholm M et al (2004) Porcine polymorphisms and methods for detecting them. International application published under the patent cooperation treaty (PCT), WO2004/048606A2
- Jørgensen CB, Cirera S, Anderson SI, Archibald AL, Raudsepp T et al (2003) Linkage and comparative mapping of the locus controlling susceptibility towards *E. coli* F4ab/ac diarrhoea in pigs. *Cytogenet Genome Res* 102:157–162
- Karlsson KA (1998) Meaning and therapeutic potential of microbial recognition of host glycoconjugates. *Mol Microbiol* 29:1–11
- Luther H, Vögeli P, Hofer A (2009) Increasing genetic *E. coli* F18 resistance in Swiss pigs. Available at http://www.eaap.org/Barcelona/Book_Abstracts.pdf
- Meijerink E, Fries R, Vögeli P, Masabanda J, Wigger G et al (1997) Two alpha(1, 2) fucosyltransferase genes on porcine chromosome 6q11 are closely linked to the blood group inhibitor (S) and *Escherichia coli* F18 receptor (ECF18R) loci. *Mamm Genome* 8:736–741
- Meijerink E, Neuenschwander S, Fries R, Dinter A, Bertschinger HU et al (2000) A DNA polymorphism influencing alpha(1, 2)fucosyltransferase activity of the pig FUT1 enzyme determines susceptibility of small intestinal epithelium to *Escherichia coli* F18 adhesion. *Immunogenetics* 52:129–136
- Osek J (1999) Prevalence of virulence factors of *Escherichia coli* strains isolated from diarrheic and healthy piglets after weaning. *Vet Microbiol* 68:209–217
- Python P, Jorg H, Neuenschwander S, Hagger C, Stricker C et al (2002) Fine-mapping of the intestinal receptor locus for enterotoxigenic *Escherichia coli* F4ac on porcine chromosome 13. *Anim Genet* 33:441–447
- Ramos AM, Crooijmans RP, Affara NA, Amaral AJ, Archibald AL et al (2009) Design of a high density SNP genotyping assay in the pig using SNPs identified and characterized by next generation sequencing technology. *PLoS One* 4:e6524
- Rasschaert K, Verdonck F, Goddeeris BM, Duchateau L, Cox E (2007) Screening of pigs resistant to F4 enterotoxigenic *Escherichia coli* (ETEC) infection. *Vet Microbiol* 123:249–253
- Rippinger P, Bertschinger HU, Imberechts H, Nagy B, Sorg I et al (1995) Designations F18ab and F18ac for the related fimbrial types F107, 2134P and 8813 of *Escherichia coli* isolated from porcine postweaning diarrhoea and from oedema disease. *Vet Microbiol* 45:281–295
- Thiele H, Nürnberg P (2005) HaploPainter: a tool for drawing pedigrees with complex haplotypes. *Bioinformatics* 21:1730–1732
- Vögeli P, Bolt R, Fries R, Stranzinger G (1994) Co-segregation of the malignant hyperthermia and the Arg615-Cys615 mutation in the skeletal muscle calcium release channel protein in five European Landrace and Pietrain pig breeds. *Anim Genet* 25 Suppl 1:59–66
- Vögeli P, Bertschinger HU, Stamm M, Stricker C, Hagger C et al (1996) Genes specifying receptors for F18 fimbriated *Escherichia coli*, causing oedema disease and postweaning diarrhoea in pigs, map to chromosome 6. *Anim Genet* 27:321–328
- Wang J, Jiang SW, Chen XH, Liu ZL, Peng J (2006) Prevalence of fimbrial antigen (K88 variants, K99 and 987P) of enterotoxigenic *Escherichia coli* from neonatal and post-weaning piglets with diarrhea in central China. *Asian Austral J Anim Sci* 19(9):1342–1346
- Zhang B, Ren J, Yan X, Huang X, Ji H et al (2008) Investigation of the porcine *MUC13* gene: isolation, expression, polymorphisms and strong association with susceptibility to enterotoxigenic *Escherichia coli* F4ab/ac. *Anim Genet* 39:258–266