Countries influence the trade-off between crop yields and nitrogen pollution

Journal Article

Author(s):

Wüpper, David Johannes ; Le Clech, Solen; Zilberman, David; Mueller, Nathaniel; Finger, Robert

Publication date:

2020-11-11

Permanent link:

<https://doi.org/10.3929/ethz-b-000452561>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Originally published in:

Nature Food 1, <https://doi.org/10.1038/s43016-020-00185-6>

Countries Influence the Trade-Off between Crop Yields and Nitrogen Pollution

David Wuepper^{1*}, Solen Le Clech², David Zilberman³, Nathaniel Mueller⁴, Robert Finger¹

¹ ETH Zürich, Switzerland, ² Wageningen University, Netherlands,

³ UC Berkeley, USA, ⁴ Colorado State University, USA

*corresponding author: dwuepper@ethz.ch

POSTPRINT to

Wuepper, D., Le Clech, S., Zilberman, D. et al. Countries influence the trade-off between crop yields and nitrogen pollution. Nat Food 1, 713–719 (2020).

<https://doi.org/10.1038/s43016-020-00185-6>

Abstract: National institutions and policies could provide powerful levers to steer the global food system towards higher agricultural production and lower environmental impact. However, causal evidence of countries' influence is scarce. Using global geospatial datasets and a regression discontinuity design, we provide causal quantifications how much crop yield gaps, nitrogen pollution, and nitrogen pollution per crop yield, are influenced by country-level factors, such as institutions and policies. We find that countries influence nitrogen pollution much more than crop yields and there is only a small trade-off between reducing nitrogen pollution and increasing yields. Overall, countries that cause 35% less nitrogen pollution than their neighbors only cause a 1 percent larger yield gap (the difference between attainable and attained yield). Explanations which countries cause the most pollution relative to their crop yields include economic development, population size, institutional quality, foreign financial flows to land resources, as well as countries' overall agricultural intensity and its share in the economy. Our findings suggest that many national governments have an impressive capacity to reduce global nitrogen pollution without having to sacrifice much agricultural production.

Main

The global food system is at the epicenter of many of this century's greatest challenges¹⁻⁴. To match growing demand, crop production will need to increase 25–70% from 2015 to 2050⁴⁻⁷. Because natural

ecosystems must be simultaneously protected, increased production cannot mainly come from agricultural area expansion, but per-area production must increase^{3,8,9}. Yet, increasing input intensity can negatively affect water and air quality, climate, biodiversity, and human health¹⁰⁻¹⁵.

The main solution to this food-water-environment nexus is to improve agricultural input use efficiency alongside the necessary increase in inputs^{2,6,13}. However, the opportunities to increase yields while keeping environmental impacts low are context and country specific. Socio-economic circumstances, policies, institutions, and regulations are a few examples of the country-level variables that affect crop mixes, input use, technologies and thus the resulting yields and environmental effects of crop production^{13,16,17}. For example, for many years, nitrogen fertilizer was heavily subsidized in China^{13,18}. More recently, China phased out these subsidies and started to fund improvements in nitrogen and manure management¹⁹⁻²². However, there are still policies in place that negatively affect nitrogen use efficiency^{21,22}. Overall, China uses >30% of all global fertilizer on only 9% of global cropland while achieving intermediate yields²².

Because we now live on a “cultivated planet”², we can now often already see the impact of country-level factors on satellite images^{17,23} (**Figure 1**). The fields in China are visibly greener than the fields in Kazakhstan and the fields in Turkey are visibly greener than the fields in Syria, and importantly, the changes pop up right at the border. In both examples, greener fields generally indicate higher agricultural production intensity. The only reason why we see these border discontinuities is that the neighboring countries- as political entities - influence farmer decisions where and how to grow what – and, as we establish below, there is no natural discontinuity of environmental conditions at these borders.

FIGURE 1 HERE

Fig.1. Revealing borders. Right at the political borders between (a) Kazakhstan and China and between (b) Turkey and Syria, the color of the agricultural fields changes discontinuously, revealing the impact of countries on agricultural production decisions. Greener colors indicate higher production intensity. Credit: (a) NASA Earth Observatory image by Robert Simmon, using Landsat data from the U.S. Geological Survey, (b) Copernicus Sentinel image, retrieved from Google Earth Engine

Here, we propose an approach to estimate countries’ causal effect on their crop yields, expressed as yield gaps that account for differences in local attainable yields (for wheat, maize, rice, potato, soy, sorghum, and cassava), their nitrogen balances on croplands and their nitrogen pollution in freshwater,

as well the relationship between countries' effect on their yields and their pollution. Our approach is a formal econometric framework that is based on the logic of the examples shown above, applied at the global scale (examining 289 land-borders around the world). Our analysis also allows us to investigate the driving forces behind the empirical patterns.

Results

In general, most countries are not comparable to each other and they also do not have 100% influence on all the agricultural and environmental outcomes on their territories. For example, agricultural and environmental outcomes are strongly influenced by a range of natural factors that are mostly outside the influence of the countries. Moreover, in most countries around the world, there is at least some degree of cultural and institutional variation that also affects agricultural and environmental outcomes that pre-dates the current countries²⁴⁻²⁶. Considering the examples from above (**Figure 1**), Kazakhstan and China, and Turkey and Syria are very different in terms of e.g. their weather, soils, and other natural characteristics. It is only close to their borders where they become more and more comparable.

Our approach is to analyze only observations within a narrow band around political borders, where natural conditions tend to be more comparable, control for the general spatial distribution of yields and pollution, and estimate whether there are statistically significant discontinuities right at the countries' borders that can only be explained by country-level characteristics and actions, and not e.g. local or regional confounders^{17,27,28}. This is a regression discontinuity design^{29,30} which we estimate as a system of simultaneous regressions^{31,32}. We describe and discuss this framework and all its assumptions in the **Materials and Methods**. We can visually illustrate the main mechanics, showing the spatial distribution of four outcomes, locally averaged in 200 bins of 300 meters length on each border side, as a function of border distance, for approximately all land-borders and countries around the world (**Figure 2**). The outcomes are the nitrogen balance on croplands (**a**)³, nitrogen pollution in freshwater (**b**)¹⁴, average yield gaps (the difference between a location's attainable yield versus what is actually attained)(**c**)⁸, and the natural vegetation mixture that we would observe without human impact, which is a valuable summary indicator for overall environmental differences, expressed as percentage of naturally occurring tree cover from 0 to 100% (**d**)³³. The observations are sorted the same way in all four plots (**a – d**), namely

by each country's comparative water pollution (see **Materials and Methods**). In each plot, the bins shown on the left of the border (vertical, dashed line) are from countries that cause more nitrogen water pollution than their neighboring countries. Then, the fitted linear trends (solid, black line on each side of the border) indicates the general spatial pattern (e.g. the yield gaps around a particular border might continuously change from west to east and north to south, because of continuously changing rainfall and soil fertility). A discontinuity right at the border suggests an effect of the individual countries, whereas continuity right at the border would suggest no effect. It can be seen that globally, we see a sharp border discontinuity in cropland nitrogen balances (**a**), nitrogen water pollution (**b**), and yield gaps (**c**) - but not in the natural vegetation potential (**d**). The last finding is important for the interpretation of our other results. If the border areas of countries with more nitrogen pollution and smaller yield gaps were to be naturally different from the border areas of countries with less nitrogen water pollution and larger yield gaps, this would compromise our identification strategy, because then border discontinuities could either be explained by the effect of the countries *or* the effect of their natural environment. Our data suggests that, in general, the border sides are naturally comparable, and countries' characteristics and actions (which we explore below) cause significant differences in nitrogen pollution and crop yields. However, we do find natural discontinuities at some borders and we carefully test whether these borders affect our results and control for these natural discontinuities when analyzing individual borders further below.

FIGURE 2 HERE

Fig 2. Spatial distributions of nitrogen balances, water pollution, yield gaps and the natural vegetation potential around international borders. Border discontinuities were examined in (**a**) cropland nitrogen balances (n=151,232), (**b**) nitrogen in water pollution (n=200,367), (**c**) yield gaps (n= 115,901), and (**d**) to test our main identifying assumption, natural vegetation potential (expressed as percentage of naturally occurring tree cover from 0 to 100%)(n= 200,367). Shown are all global data points that fall within a bandwidth of 60km to at least one of 289 land-borders all around the world, here averaged in 200 bins of 300m width. All outcomes are shown as standard deviations from their own mean. Border distances are shown in kilometers.

Global Estimates of Countries Influence on their Yield gaps and Nitrogen Pollution

We obtain three main results (**Figure 3**). First, we estimate that on global average, countries cause a much larger discontinuity in the spatial distribution of nitrogen water pollution than in the spatial distribution of yield gaps. The discontinuity is around 35% for nitrogen pollution, but only between 1

and 1.5% for yield gaps. Second, the two discontinuities are inversely related, i.e. countries that achieve lower yield gaps (higher yields) tend to cause more nitrogen pollution. Third, the estimates are robust, only varying slightly in magnitude between different specifications and sub-samples.

FIGURE 3 HERE

Fig 3. Estimated effect of countries on their yield gaps and nitrogen pollution.

Circles show point estimates and bars show the 95% confidence interval. In general, the countries that achieve smaller yield gaps cause disproportionately more nitrogen pollution than their neighboring countries, and vice versa. In our baseline specification (**1a**, $n=91,472$), we control for border distance, separately on each side of each border, fixed border effects, and a linear polynomial of longitude and latitude. Standard errors are clustered by border. As one robustness check, we then add covariates regarding environmental characteristics (altitude, depth of bedrock, precipitation during wet and dry season as well as overall precipitation, and soil organic carbon) in the yield gap equation and human and animal densities (pigs, cattle, and chicken) in the nitrogen pollution equation (**1b**, $n=91,472$). With this, we test whether it matters that sometimes the natural environment is different on one side of the border than it is on the other, even if this is not so on average, and whether potential discontinuities in human and animal population densities might confound our nitrogen pollution estimates. Next, we exclude borders at which we find any discontinuity in the natural environment, first without (**2a**, $n=73,754$) and then with covariates (**2b**, $n=73,754$). This is an alternative test for the influence of “natural” borders versus “purely political” borders, and - as we explain below - also for the influence of data density around each border. Finally, we exclude borders with low-resolution input data (for some countries, important input data is only available at high aggregation), again without (**3a**, $n=88,692$) and then with covariates (**3b**, $n=88,692$). With this, we probe how measurement errors (both random and systematic) influence our estimates. We find only little variation in our results across specifications.

The Spatial Distribution of Countries’ Effect on their Yield Gaps and Nitrogen Pollution

Moving beyond global averages, **Figure 4** shows a global map of each country’s estimated effect on its crop yields compared to its effect on nitrogen pollution (an aggregation of each country’s estimated effect on nitrogen pollution minus its estimated effect on crop yields).

FIGURE 4 HERE

Fig.4. Countries’ estimated effect on their yield gaps versus their nitrogen pollution.

Quantification of how much nitrogen pollution countries are causing compared to how much they reduce their yield gaps, relative to directly neighboring countries. Darker colors reflect larger increases in nitrogen pollution compared to the closing of yield gaps, lighter colors reflect larger decreases in nitrogen pollution compared to widening yield gaps.

By far the highest value is estimated for China (170%, which can be interpreted as China is causing 170% more nitrogen pollution than it is reducing its crop yield gaps, both compared to all its neighboring countries). Other countries with less but still particularly high nitrogen pollution estimates compared to

their neighboring countries include Brazil, Mexico, Colombia, Israel, Thailand, and Georgia, whereas countries such as the United States, Germany, France, South Korea, and Austria achieve relatively high yields with comparably less nitrogen pollution. Important caveats are that all estimates are based completely on comparisons between neighboring countries, so they are strictly relative. This means e.g. that the positive, comparative yield effect of South Korea can be as much attributed to its own high yields as it can be attributed to the particularly low yields in North Korea, and the comparative nitrogen pollution effects of Kazakhstan and Mongolia, e.g., are mostly attributable to the fact that they are being compared to China. It should also be noted that these relative values are an unweighted average of all discontinuities, i.e. a country might have a positive effect compared to one neighbor and a negative effect of a similar magnitude compared to another, and its final value then is close to zero. Finally, we have no results for the few countries worldwide that have no direct neighbors (e.g. Australia).

Explanations for Countries' Effect on their Yield Gaps and Nitrogen Pollution

Countries that produce a disproportionate amount of nitrogen pollution relative to their yield performance likely have potential to reduce pollution without large sacrifices in terms of yield. To learn what distinguishes the countries that cause more pollution per yield from those that cause less, we regress countries' estimated effect from above on regional fixed effects (e.g. Sub-Saharan Africa, North America, etc.) and individual explanatory variables, which are shown together with their estimated coefficients below (**Figure 5**).

FIGURE 5 HERE

Fig.5. Explaining countries' estimated pollution versus yield gaps effect. This figure shows the results from linear regressions of countries' estimated effect on their nitrogen pollution versus their effect on their crop yields (**Fig.4**) on regional fixed effects and a broad range of potential explanatory variables (n=143). To test a potential non-linear relationship with gdp and/or gdp growth, these variables were included linearly as well as squared.

Previous studies have found an asymmetric para-curve relationship between nitrogen pollution and economic development^{13,34} (an Environmental Kuznets curve). Consistent with this, we find that middle income countries cause the most nitrogen pollution compared to the yields they achieve (e.g. China or Brazil) whereas several richer countries cause less nitrogen pollution compared to the yields they achieve (e.g. Germany or the US). Overall, however, we find an approximately linearly increasing

relationship between countries' gdp and their pollution per yield (specifications **1a** and **1b**). Simply put, richer countries cause considerably more nitrogen pollution than poorer countries and this is not matched by commensurate yield advantages. In contrast, there is no associations with gdp *growth* (**2a** and **2b**). However, countries with larger populations cause more pollution compared to their yield effect than countries with smaller populations (**3**). There is no association with population *growth* (**4**).

We also find a positive association with the quality of countries' economic institutions³⁵ (**5**). This suggests that globally, better institutions are more tightly associated with heightened environmental impact than better environmental regulation.

A small, but statistically significant, positive association is found with foreign investments into countries' land resources (**6**). There is no association with countries' agricultural gdp share (**7**) but we find that more intensive farming systems are associated with more pollution per yield (**8**). At the same time, more pollution per yield is also associated with a significantly higher availability and affordability of food (**9** and **10**), which cautions to consider food security issues in this context.

Future research might moreover consider the influence of specific policies^{19,22}, national legislations³⁶, behavioral factors such as culture²⁶, and different farm sizes^{21,37}.

Discussion

Our global food system is more productive than ever before in human history. However, productivity growth may still be slightly below what we need to match projected future demand⁴⁻⁶. At the same time, environmental impacts, foremost nitrogen pollution, are far beyond any safe level^{10,38}. Both are in large parts because nitrogen use is currently spatially inefficiently distributed^{11,39,40}. We here have exploited a natural experiment created by spatial discontinuities at international borders, to identify the role of countries. We find that they have a much larger effect on nitrogen pollution than they have on yield gaps and many countries cause very high pollution for the yield that they actually achieve. Importantly, closing yield gaps and mitigating nitrogen pollution is not a technically necessary trade-off. In particular, our empirical results suggest that nitrogen surpluses can be reduced by ~35% if the more polluting countries around the world only achieve the pollution levels of their less polluting neighboring countries and even without any adjustments, this would only increase yield gaps by ~1%. Important to note, this

1 1% increase in yield gaps is not necessary, because countries can commonly adjust and even synergies
2 exists. As we find, this is largely under the control of national governments.

3 An important lever for national governments is the ratio of fertilizer price to agricultural output price.
4 Countries with nitrogen surpluses might re-allocate financial resources from agricultural subsidies that
5 increase environmental impacts to those that incentivize a more environmentally friendly production.
6 Another option is the introduction of taxation schemes that raise the relative price of nitrogen compared
7 to its production value^{41,42}. There are also other possibilities for countries to support the adoption of new,
8 more sustainable technologies and farming practices, e.g. via improving extension systems, and
9 changing environmental and tenure regulations^{13,16,19-22}. The opposite applies for countries with large
10 yield gaps, where it is important to support (sustainable) intensification, e.g. via input subsidies^{43,44}. An
11 increasingly important role might be played by precision farming and new plant breeding in the future,
12 which require national government support and could vastly increase nitrogen use efficiency^{45,46}.

13 **Materials and Methods**

14 To understand countries' effect on yield gaps and nitrogen pollution, as well as their relationship, we
15 analyze large, global datasets with a combination of a regression discontinuity design^{17,30,47} and a
16 seemingly unrelated regression framework^{31,32}.

17 We begin with a discussion of our analytical framework (I), its assumptions, and how we test them (II).
18 Then we describe our four main datasets (III), and the data we use to explain our findings (IV).

19 **Analytical Framework**

20 Our analytical framework is a combination of a regression discontinuity design^{17,30,47} and a seemingly
21 unrelated regression framework^{31,32}. The basic idea is that under a set of falsifiable assumptions, spatial
22 discontinuities right at political borders reveal the influence of the countries that are separated there.
23 Often, political borders can be easily recognized in the landscape, even though natural environmental
24 conditions are otherwise homogenous (see e.g. **Figure 1**). In a sense, the division of the world into
25 different countries functions like a “natural experiment” in many places around the globe.
26 Technically, we simultaneously estimate whether there is a statistically significant discontinuity at
27 political borders in yield gaps and nitrogen pollution, while controlling for all continuously distributed

confounders via linear polynomials of border distance and longitude and latitude, as well as covariates.

For this, we first estimate at each border which country potentially causes more nitrogen pollution:

$$\log N\text{ Pollution}_i = \beta_1 D_i^a + \beta_2 \text{dist}_i^A + \beta_3 \text{dist}_i^B + \beta_4 \text{Vegetation}_i + \epsilon_i \text{ if } \text{dist}_i \leq \varphi^* \quad (1)$$

where $\log N\text{ Pollution}_i$ is the natural logarithm of nitrogen pollution in water¹⁴ of pixel i on a gridded global map at 5 arc-minute resolution, D_i^a indicates which of the two countries A or B at every border might pollute its waters more with nitrogen¹⁴, dist_i^A and dist_i^B control for border-distance, separately in country A and B, *Vegetation* controls for natural vegetation potential, expressed as naturally occurring tree cover in percentage from 0 to 100%, which summarizes a large number of environmental factors³³, ϵ_i is an error-term, and φ^* is the estimated optimal bandwidth (the optimal maximum distance to each border, defining our sample), balancing bias and precision⁴⁸. In our case, this is 20km on each side. We then estimate a global model, in which we simultaneously estimate whether there is a significant border discontinuity in the average yield gap between the countries with more nitrogen pollution and those with less and whether there is a significant border discontinuity in nitrogen pollution between the same countries:

$$\text{Yield Gap}_i = \beta_1 D_i^a + \beta_2 \text{dist}_i^H + \beta_3 \text{dist}_i^L + \beta_4 \text{location}_i + \beta_5 \theta_i^a + \beta_6 \vartheta_j + \epsilon_i^a \text{ if } \text{dist}_i \leq \varphi^* \quad (2a)$$

$$\log N\text{ Pollution}_i = \beta_7 D_i^b + \beta_8 \text{dist}_i^H + \beta_9 \text{dist}_i^L + \beta_{10} \text{location}_i + \beta_{11} \theta_i^b + \beta_{12} \vartheta_j + \epsilon_i^b \text{ if } \text{dist}_i \leq \varphi^* \quad (2b)$$

where Yield Gap_i is the average yield gap⁸ in percentage (the difference between a place's achievable yield as a function of environmental constraints and the actually achieved yield). ϑ_j indicates to which pairwise border the observations belongs, θ_i^a is a vector of six environmental covariates³³ summarizing the influence 58 individual environmental characteristics, such as topography and bio-climate, location_i is described by longitude, latitude, and their interaction, dist_i^H and dist_i^L are linear polynomials of border distance, fit separately on both side of each border, and D_i^a quantifies the border discontinuity. Again, φ^* is the estimated optimal maximum border distance (the “bandwidth”) that minimizes omitted variable bias with the largest sample possible. The error term of this first equation is assumed to be correlated with the error term of the second equation, as the observations are from the same place. For the second equation, which is simultaneously estimated with the first, the left side variable is $\log N\text{ Pollution}_i$ which is alternatively the average nitrogen footprint¹⁴ in the freshwater or

the nitrogen balance on agricultural land³ of pixel i on a gridded global map at 5 arc-minute resolution. Then, θ_i^b is a vector of four population densities (chicken, cattle, pigs, and people)^{49,50}, D_i^b quantifies again the border discontinuity, and all other variables are defined as above. Throughout, standard errors are clustered at the border, accounting for common unobservables and spatial auto-correlation. We also always transform the estimated effects so they are expressed as percentage changes (using the inverse of the logarithmic function). Finally, we estimate individually at each border the effect of the countries on their yield gaps and their nitrogen pollution, to understand the trade-off between mitigating nitrogen pollution and closing yield gaps. Here, we use cropland N-balances as our measure for nitrogen pollution³, to avoid confounding by non-agricultural sources, which is more likely when analyzing individual borders. We then aggregate all estimated effects by country (ignoring discontinuities from borders with natural discontinuities) and regress each country's pollution versus yield effect on regional fixed effects and hypothesized explanations:

$$\text{pollution versus yield effect} = \beta_1 X_i + \beta_2 \text{Region}_i + \epsilon_i \quad (3)$$

Where X_i are possible explanations, such as countries' gross domestic product, institutional quality, and several others, and Region_i are fixed effects for Sub-Saharan Africa, the Middle East and North-Africa, Europe and Central Asia, South Asia, East Asia and the Pacific, and North and Latin America.

Assumptions and Tests

Our main identifying assumptions are that we can distinguish between exogenous and endogenous borders (**a**), that we can learn something about countries from focusing on their border areas (**b**), that we have sufficient data near borders (**c**), and that systematic and random measurement error in our input data is not a first-order problem (**d**). We discuss each assumption and how we probe it below.

a. Exogenous Borders

Our central assumption is that border discontinuities in nitrogen pollution and yield gaps reveal the causal influence of countries because there are no “compound treatments”. The “treatment” we are interested in is that one side of each border belongs to one country, and the other side belongs to another country. If, however, also one side has one type of soil and the other side another, then we cannot

interpret a border discontinuity as country-effect but it is possibly in part or in full the result of the difference in soil type. Thus, we must establish that international borders mostly divide naturally homogenous areas, i.e. that they are exogenous to differences in yields and nitrogen pollution.

In **Figure 2d** it is already shown that the countries that cause more nitrogen pollution than their neighboring countries do not have systematically different environmental and geographical conditions close to their borders (summarized by their hypothetical natural vegetation). To test this statistically, we focus on three main determinants of the natural vegetation, which are altitude, temperature, and precipitation and estimate – similar to how we quantify the border discontinuities in yield gaps and nitrogen pollution – whether there is a border discontinuity in any of these indicators:

$$Indicator_i = \beta_1 D_i + \beta_2 dist_i^H + \beta_3 dist_i^L + \beta_4 \theta_i + \epsilon_i \text{ if } dist_i \leq \varphi^+ \quad (4)$$

Where the $Indicator_i$ is alternatively altitude, temperature, and rainfall, D_i reveals whether there is a “jump” in the relevant indicator right at the border, $dist_i^H$ and $dist_i^L$ are linear polynomials of border distance, separately fitted on the two sides of each border, and θ_i are fixed effects for each border. As before, standard errors are clustered at the border. Neither altitude, temperature, nor precipitation exhibit any discontinuity at the average border that we use for our analysis. Moreover, moving from a maximum border distance of 60km to one of 30km, the point estimates move closer to zero, consistent with the idea that we increase the environmental comparability of observations by excluding observations further away from the border. This is shown in **Supplementary Figure 1** in the Supplementary Materials.

At individual borders, we sometimes do find natural discontinuities, so these borders are less reliable for our analysis and we investigate this issue further below. First, we examine our second assumption, which is that we can estimate the causal effect of countries in border areas.

b. Representative Borders

We achieve high internal validity - among others reasons - by only analyzing already quite comparable observations close to borders. This, however begs the question how representative international border areas are for countries’ interiors⁵¹. We examine this with a simple correlational analysis of the nitrogen pollution found in border areas and the nitrogen pollution of the entire countries. **Supplementary Figure 2** shows that there is generally quite a high correlation, even though there are outliers in this pattern, and, as we discuss below, there is an in-built bias in the data towards this pattern. Overall, this

is suggestive that we can learn something about the countries by only estimating what happens in their border areas.

c. Sufficient Data Density

Our third assumption is that we have sufficiently high densities of croplands left and right of the border. This is important because we use observations just left and right of the border as counterfactuals, and if there are not many observations, our sample is small and our estimator imprecise, and if we compare observations far away from the border, we risk increasing omitted variable bias. We test this assumption together with our first assumption that natural discontinuities at international borders are small and seldom. For this, we individually estimate at each border, whether we find a discontinuity in the hypothetical natural vegetation, as predicted by Bastin, et al. ³³. Finding a border discontinuity in the hypothetical natural vegetation comes either from an actual discontinuity in environmental characteristics right at the border, or alternatively, it comes from the fact that our observations at this particular border are actually not right at the border, but further apart. It should be noted that the natural vegetation depends on many environmental and geographic characteristics that also affect yield and nitrogen pollution potentials, so it is quite a general indicator. We estimate, similar to before:

$$Vegetation_i = \beta_1 D_i + \beta_2 dist_i^H + \beta_3 dist_i^L + \beta_4 \theta_i + \epsilon_i \text{ if } dist_i \leq \varphi^* \quad (5)$$

Where $Vegetation_i$ is the vegetation we would see all around the world if there was no human impact. All other variables are defined as above.

We mark all borders at which we find a statistically significant border discontinuity as “less reliable” (19% of all borders) and all others as “more reliable” (81% of all borders).

d. Measurement Error

All else equal, modelled data is often less precise than remote sensing data because it involves at least one more processing step. Thus, there is at least one more source of measurement error. Often, our data is based on multiple processing steps and it is unlikely that any one is error free. Second, due to data availability constraints, the global distribution of nitrogen pollution has a rather low resolution. It is not possible currently to model this at the same resolution like e.g. deforestation ⁵² or soil erosion ⁵³. For issues such as nutrient pollution ¹⁴ or greenhouse gas emissions ⁵⁴, a 5-arc-minute resolution is currently

1 the highest available resolution. This, however, means that a certain degree of measurement error it
2 unavoidable.

3 It is also noteworthy that both yields and water pollution by nitrogen are related to each other but also
4 caused by third variables. For example, yields are strongly affected by water availability^{3,5,8} (e.g. via
5 irrigation) and nitrogen in water also comes from non-cropland and non-agricultural sources^{14,34,55,56}.

6 Whereas random measurement error might lead to a bias towards statistically insignificant and/or
7 smaller border discontinuities, our data could also contain systematic measurement error with the
8 opposite effect. At first, one would not expect any systematic measurement error in our data because all
9 our main variables are from datasets that have been created globally homogenously. However, there is
10 a hidden source of systematic measurement error and that is low-resolution, statistical data that has been
11 used as input data. For example, in the N-pollution data of Mekonnen and Hoekstra¹⁴, the distribution
12 of cropland and production systems, manure input and N-output are all sub-national, but mineral N-
13 fertilizer applications are only available at the country level⁵⁷⁻⁵⁹ and for some countries also the yield
14 data is only available at this level⁶⁰. The cropland nitrogen balance data³ incorporate some subnational
15 fertilizer application rate data from Mueller, et al.⁸, but only for a subset of countries. The resolution of
16 our source datasets could bias estimated border discontinuities. Similar to the rather low resolution
17 increasing random measurement error, this issue cannot be solved because for many countries in the
18 world there exists no reliable data sub-national fertilizer application rates. However, even if the ratio of
19 random to systematic measurement error is such that we over or underestimate the average border
20 discontinuity in yield gaps and/or especially nitrogen pollution, the estimated association between
21 countries effect on pollution and yields can still be correctly estimated.

22 To take into account that yield gaps and nitrogen water pollution are caused by other sources than each
23 other, we estimate all our specifications once without covariates and then including potential
24 confounding factors, such as the population densities of humans and several animal species. This allows
25 us to test how sensitive our estimates are e.g. to the influence of nitrogen pollution from domestic and
26 industrial sources or yield differences caused by rainfall patterns. Specifically for nitrogen pollution in
27 water¹⁴, we also examine cropland nitrogen balances³, which are the intermediate channel (see e.g.

Figure 2 above and **Supplementary Figure 3**). Moreover, Monfreda, et al. ⁶⁰ provide data on the resolution of their utilized agricultural information, which is both used for the modelling of yield gaps and nitrogen pollution. Thus, we are able to test the robustness of our estimates by excluding observations at the bottom end of the quality spectrum (see **Figure 3** above and **Supplementary Figure 4** in the **Supplementary Materials**).

Main Data

Our main data sources are Mueller, et al. ⁸ for the global distribution of yield gaps, Mekonnen and Hoekstra ¹⁴ for the global distribution of nitrogen pollution, West, et al. ³ for the global distribution of nitrogen balances on croplands (the direct connection between closing yield gaps and increasing nitrogen-pollution in freshwater), and Bastin, et al. ³³ for global data on the distribution of the natural environment without human impact (to evaluate our “border exogeneity” assumption). Moreover, we use environmental characteristics also provided by Bastin, et al. ³³, and data on the human population density from SEDAC ⁴⁹ and animal population densities from Gilbert, et al. ⁵⁰. We re-sampled all datasets at a 5-arc-minute resolution which is the resolution of the dataset of Mekonnen and Hoekstra ¹⁴ and dropped all observations further than 100 km away from any border.

To compute yield gaps, first attainable yields are needed. For this, Mueller, et al. ⁸ created 100 zones of similar annual precipitation ⁶¹ and growing degree-day characteristics ⁶². Then, they defined attainable yield as the area-weighted 95th percentile observed yield within each bin. The yield data is from Monfreda, et al. ⁶⁰. The yield gap of each place is then defined as the difference between attainable and actually attained yield. An important “proximate” (in contrast to “fundamental”) explanation for the existence of yield gaps are differences in nitrogen fertilizer applications⁸. The main advantage of working with yield gaps in this study - and not simply with yields - is that the yield gaps already incorporate an estimate of the yield *potential* of each observation; working with yields directly would be subject to the confounding influence of changes in yield potential due to agro-ecological differences. For this study, we computed the average yield gap, by aggregating the individual yield gaps of the six major crops (maize, wheat, potato, cassava, sorghum, soy).

1 The nitrogen pollution is defined by Mekonnen and Hoekstra ¹⁴ as greywater “footprint”, which are all
2 anthropogenic N emissions divided by the difference between the ambient water quality standard for N
3 and the natural concentration of N in the receiving water body. This approximates how much pristine
4 water is necessary to assimilate the entire nitrogen pollution. To compute N-inputs, they combined data
5 on the global distribution of croplands from Monfreda, et al. ⁶⁰, which also provides information on N-
6 fixation by legumes, then fertilizer and manure applications ^{57-59,63}, atmospheric N deposition ⁶⁴, the N
7 content of irrigation water ⁶⁵, as well as a large number of point emission sources. To compute N-
8 removal, harvests ⁶⁰, soil erosion ⁶⁶, ammonia ⁶⁷, N₂O and NO emissions ⁶⁸ are considered. Data on soil
9 parameters comes from Batjes ⁶⁹, the rooting depths of individual crops is from Allen, et al. ⁷⁰, and
10 precipitation data is from Mitchell and Jones ⁷¹.

11 To model nitrogen balances at the landscape level, West, et al. ³ first modelled nitrogen input by adding
12 crop-specific nitrogen fertilizer applications⁸ and their own estimate of manure applications, based on
13 the distribution livestock density, and crop- and pasture land, similar to the approach of Foley, et al. ²
14 and Potter, et al. ⁷². Then, nitrogen fixation by legumes was added, using the data of Smil ⁷³, and
15 atmospheric nitrogen deposition, using the data of Dentener, et al. ⁷⁴. To then model nutrient removal,
16 the nutrient density data of the USDA ⁷⁵ was combined with the harvest data from Monfreda, et al. ⁶⁰.
17 The difference between input and removal is then the estimated nitrogen balance.

18 Finally, to evaluate which border around the world are endogenously drawn, i.e. along environmental
19 discontinuities, we use the data of Bastin, et al. ³³. The most sophisticated indicator for environmental
20 border discontinuities is their globally mapped *natural* vegetation distribution. For this, they let a
21 random forest algorithm ⁷⁶ learn how differences in natural environmental characteristics predict
22 differences in natural vegetation. For this, they trained the algorithm with photo-interpretations from
23 protected areas all around the world, under the assumption that protected areas are the best available
24 demonstration for the natural vegetation in each region. Then, they used global data on summary
25 measures of 58 environmental characteristics to predict the natural vegetation all around the world. For
26 our analysis, we both use their natural vegetation map and their predictor variables, all available via
27 Google Earth Engine ²³, see also Gorelick, et al. ⁷⁷. We show the entire global distributions of all four
28 datasets discussed above in **Supplementary Figure 5**. For a visual illustration how our initial sampling

1 along international borders looks like, please see **Supplementary Figure 6** for the example of landscape
2 nitrogen balances in Asia and Latin America.

3 For a description of our data to explain the estimated global patterns, see **Supplementary Materials**
4 **Section S2**.

6 **Data availability**

7 Data can be retrieved from Wuepper, et al. ⁷⁸ and from the corresponding author upon reasonable request

9 **Code availability**

10 Code can be retrieved from Wuepper, et al. ⁷⁸ and from the corresponding author upon reasonable request

Supplementary Materials for

Countries Influence the Trade-Off between Crop Yields and Nitrogen Pollution

David Wuepper^{1*}, Solen Le Clech², David Zilberman³, Nathaniel Mueller⁴, Robert Finger¹

¹ ETH Zürich, Switzerland, ² Wageningen University, Netherlands,

³ UC Berkeley, USA, ⁴ Colorado State University, USA

*corresponding author: dwuepper@ethz.ch

This PDF file includes:

Fig. S1 to S6 and a discussion of our main explanatory variables and their data sources

Section S1. Supplementary Figures (Page S2)

Supplementary Figure 1. Testing for Environmental Border Discontinuities. (Page S2)

Supplementary Figure 2. Comparing Pollution in Border Areas and Country Averages. (Page S3)

Supplementary Figure 3. Comparing Nitrogen Pollution in Water and on Land. (Page S4)

Supplementary Figure 4. Maps of Exemplary Yield Data Resolutions. (Page S5)

Supplementary Figure 5. N Pollution, Crop Yields, and Hypothetical Natural Vegetation. (Page S6)

Supplementary Figure 6. By Analyzing Data Close to International Borders Only (Page S7)

Section S2. Data for the Exploration of Explanations (Page S8)

Supplementary Materials for Wuepper, et al. ⁷⁹. All Data and Code can be retrieved from

Wuepper, et al. ⁷⁸ and from the corresponding author upon reasonable request

Section S1. Supplementary Figures

Supplementary Figure 1. Testing for Environmental Border Discontinuities. There are no statistically significant environmental border discontinuities, on average, between the countries. Diamonds, squares, and circles indicate the point estimates, the bars in blue, green, and brown show the 95 % confidence intervals, all of which range from positive upper bounds to negative lower bounds.

1

2

3

4

5

6

7

8

9

10

Supplementary Figure 2. Comparing Water Pollution in Border Areas and Country Averages. For our analysis, we focus exclusively on international border areas, because the areas just left and right of a border are more likely to be comparable environmentally and geographically than the areas further inland. Whether we can interpret our results as extrapolatable to the rest of the countries depends on how closely related are nitrogen water pollution in border areas and further inland. The graph above correlates country averages (x-axis) with border averages (y-axis). We see a strong positive relationship, suggesting that our findings in border areas are probably relevant beyond our sample.

1

2

3 **Supplementary Figure 3. Comparing Nitrogen Pollution in Water and Nitrogen Balances on**
4 **Cropland.** We are most concerned about nitrogen pollution in water, but not all nitrogen pollution in
5 water comes from croplands. Here, however, we show that nitrogen pollution in freshwater (y-axis) and
6 nitrogen balances on croplands (x-axis) are closely related. All observations are averaged in small bins
7 (green) and their relationship is approximated with a linear regression line (blue).

The maps in **Supplementary Figure 4** below show two-resolutions of the yield data from Monfreda, et al.⁶⁰, once for maize (a), and once for potatoes (b), as examples. The yield data is used both for the modelling yield gaps and nitrogen pollution and, as can be seen below, the resolution of the data is distributed heterogeneously around the world. Above, we tested whether this influences our estimates and found that it does not. For these tests, we used the data shown in the maps below and that for all other crops that we used to compute local average yield gaps and once included this as a control variable in the baseline specification, and once excluded the countries with the lowest resolution (bottom 25%).

Supplementary Figure 4. Maps of Exemplary Yield Data Resolutions. Maize and Potatoes

The maps in **Supplementary Figure 5** show the global distributions of nitrogen balances on croplands³ (a), water pollution¹⁴ (b), attained yield as percentage of attainable⁸ (c), and the predicted natural vegetation we would see without human impact³³ (d).

Supplementary Figure 5. Nitrogen Balances and Pollution, Crop Yields, and Hypothetical Natural Vegetation. The maps a to d show the global distributions of our main input data.

1

2

3 **Supplementary Figure 6. By Analyzing Data Close to International Borders Only, We can**

4 **Reduce the Impact of Environmental Influences.** Shown here is the nitrogen balance on

5 agricultural lands close to international borders in Asia and Latin America. Darker reds reflect

6 larger nitrogen surpluses while darker greens reflect larger nitrogen deficits. The closer together

7 the observations, the more similar are topography, weather, soils, and other confounders that are

8 not caused by the countries. Still, comparing e.g. the land just in China with that just in Mongolia,

9 or land just in Brazil with that just in Bolivia, there are apparent differences that reveal the

10 influence of the countries.

11

Section S2. Data for the Exploration of Explanations

For our examination of what explains the global variation in countries' nitrogen pollution per yield, we require data on various country characteristics that relate directly or indirectly to farmers' incentives and constraints, what to grow where and how. First of all, we use data on countries gross domestic product, their value added in agriculture as percentage of gross domestic product, and their populations, all from the World Bank ⁸⁰. Previous studies have found that yield gaps close with increasing development^{3,8} whereas nitrogen pollution first increases up to a point and then decreases^{13,34}. However, most countries in the world are still on the increasing part of the curve and overall, richer countries cause more nitrogen pollution than poorer countries^{13,34}. For population size and growth, the literature suggests lower yield gaps and more nitrogen pollution. From a policy point of view highly relevant, we also examine the role of institutional quality. The data comes from Kunčič ³⁵. The influence of institutional quality is ambiguous ex ante. Agricultural productivity is clearly positively associated with better institutions ^{81,82}. On the other hand, the effect on nitrogen pollution could be positive, if they mostly increase fertilizer use, or negative, if they also improve regulatory frameworks and environmental policies ¹⁶. Then, we use data from the United Nations Food and Agriculture Organization - FAO ⁸³ – on global development flows towards land resources. Previous studies have shown that financial flows can have large effects on agricultural and environmental outcomes⁸⁴. However, again the sign is not clear again ex-ante, because less financial constraints are empirically associated both with more fertilizer input and higher yields⁸⁵. We also use data on countries' agricultural intensity, as measured by fertilizer use, with data from FAO ⁸³. Higher agricultural intensity is both associated with higher yields and more pollution and overall, this should be strongly associated with more pollution per yield, given the prior literature and our own empirical evidence in this study. Finally, two especially interesting and policy-relevant variables are countries' availability and affordability of food, provided by Chaudhary, et al. ⁸⁶. Thinking about the trade-off between yield gaps and nitrogen pollution as quantifying how much countries "buy" higher yields with environmental damage, a natural question is whether perhaps more available and affordable food for all is a benefit of lowering the costs of agricultural production by externalizing costs to the environment.

References

- 1 Rockström, J., Edenhofer, O., Gaertner, J. & DeClerck, F. Planet-proofing the global food system. *Nature Food* **1**, 3-5, doi:10.1038/s43016-019-0010-4 (2020).
- 2 Foley, J. A. *et al.* Solutions for a cultivated planet. *Nature* **478**, 337 (2011).
- 3 West, P. C. *et al.* Leverage points for improving global food security and the environment. *Science* **345**, 325-328 (2014).
- 4 Hunter, M. C., Smith, R. G., Schipanski, M. E., Atwood, L. W. & Mortensen, D. A. Agriculture in 2050: recalibrating targets for sustainable intensification. *Bioscience* **67**, 386-391 (2017).
- 5 Ray, D. K., Mueller, N. D., West, P. C. & Foley, J. A. Yield trends are insufficient to double global crop production by 2050. *PloS one* **8**, e66428 (2013).
- 6 Tilman, D., Balzer, C., Hill, J. & Befort, B. Global food demand and the sustainable intensification of agriculture. *Proceedings of the National Academy of Sciences* **108**, 20260-20264 (2011).
- 7 Godfray, H. C. J. *et al.* Food security: the challenge of feeding 9 billion people. *Science* **327**, 812-818 (2010).
- 8 Mueller, N. D. *et al.* Closing yield gaps through nutrient and water management. *Nature* **490**, 254 (2012).
- 9 Folberth, C. *et al.* The global cropland-sparing potential of high-yield farming. *Nature Sustainability* **3**, 281-289, doi:10.1038/s41893-020-0505-x (2020).
- 10 Steffen, W. *et al.* Planetary boundaries: Guiding human development on a changing planet. *Science* **347**, 1259855 (2015).
- 11 Stevens, C. J. Nitrogen in the environment. *Science* **363**, 578-580 (2019).
- 12 Seitzinger, S. P. & Phillips, L. Nitrogen stewardship in the Anthropocene. *Science* **357**, 350-351, doi:10.1126/science.aao0812 (2017).
- 13 Zhang, X. *et al.* Managing nitrogen for sustainable development. *Nature* **528**, 51, doi:10.1038/nature15743 (2015).
- 14 Mekonnen, M. M. & Hoekstra, A. Y. Global gray water footprint and water pollution levels related to anthropogenic nitrogen loads to fresh water. *Environmental science & technology* **49**, 12860-12868 (2015).
- 15 Townsend, A. R. *et al.* Human health effects of a changing global nitrogen cycle. *Frontiers in Ecology and the Environment* **1**, 240-246 (2003).
- 16 Kanter, D. R. *et al.* Nitrogen pollution policy beyond the farm. *Nature Food*, doi:10.1038/s43016-019-0001-5 (2019).
- 17 Wuepper, D., Borrelli, P. & Finger, R. Countries and the global rate of soil erosion. *Nature Sustainability* **3**, 51-55, doi:10.1038/s41893-019-0438-4 (2020).
- 18 Yu, C. *et al.* Managing nitrogen to restore water quality in China. *Nature* **567**, 516-520, doi:10.1038/s41586-019-1001-1 (2019).
- 19 Searchinger, T. D. *et al.* Revising Public Agricultural Support to Mitigate Climate Change. (World Bank, Washington D.C., 2020).
- 20 Cui, Z. *et al.* Pursuing sustainable productivity with millions of smallholder farmers. *Nature* **555**, 363-366 (2018).
- 21 Ju, X., Gu, B., Wu, Y. & Galloway, J. N. Reducing China's fertilizer use by increasing farm size. *Global environmental change* **41**, 26-32 (2016).
- 22 Wu, Y. *et al.* Policy distortions, farm size, and the overuse of agricultural chemicals in China. *Proceedings of the National Academy of Sciences* **115**, 7010-7015 (2018).
- 23 Google Earth Engine. <<https://earthengine.google.com/>> (2020).
- 24 Alesina, A., Tabellini, G. & Trebbi, F. Is Europe an optimal political area? , (National Bureau of Economic Research, 2017).
- 25 Alesina, A. & Giuliano, P. Culture and institutions. *Journal of Economic Literature* **53**, 898-944 (2015).

- 26 Wuepper, D. Does culture affect soil erosion? Empirical evidence from Europe. *European Review of Agricultural Economics* **47**, 619-653 (2020).
- 27 Pinkovskiy, M. Growth discontinuities at borders. *Journal of Economic Growth* **22**, 145-192 (2017).
- 28 Keele, L. & Titiunik, R. Natural experiments based on geography. *Political Science Research and Methods* **4**, 65-95 (2016).
- 29 Cattaneo, M. & Escanciano, J. *Regression Discontinuity Designs: Theory and Applications* (Emerald Group Publishing, 2017).
- 30 Lee, D. S. & Lemieux, T. Regression discontinuity designs in economics. *Journal of Economic Literature* **48**, 281-355 (2010).
- 31 Fiebig, D. G. Seemingly unrelated regression. *A companion to theoretical econometrics*, 101-121 (2001).
- 32 Smith, M. & Kohn, R. Nonparametric seemingly unrelated regression. *Journal of Econometrics* **98**, 257-281 (2000).
- 33 Bastin, J.-F. *et al.* The global tree restoration potential. *Science* **365**, 76-79 (2019).
- 34 Gu, B. *et al.* Cleaning up nitrogen pollution may reduce future carbon sinks. *Global environmental change* **48**, 56-66 (2018).
- 35 Kunčič, A. Institutional quality dataset. *Journal of institutional economics* **10**, 135-161 (2014).
- 36 Eskander, S. M. S. U. & Fankhauser, S. Reduction in greenhouse gas emissions from national climate legislation. *Nature Climate Change*, doi:10.1038/s41558-020-0831-z (2020).
- 37 Lesiv, M. *et al.* Estimating the global distribution of field size using crowdsourcing. *Global change biology* **25**, 174-186 (2019).
- 38 Rockström, J. *et al.* A safe operating space for humanity. *nature* **461**, 472 (2009).
- 39 Mueller, N. D. *et al.* Declining spatial efficiency of global cropland nitrogen allocation. *Global Biogeochemical Cycles* **31**, 245-257 (2017).
- 40 Lassaletta, L. *et al.* Nitrogen use in the global food system: past trends and future trajectories of agronomic performance, pollution, trade, and dietary demand. *Environmental Research Letters* **11**, 095007, doi:10.1088/1748-9326/11/9/095007 (2016).
- 41 Pe'Er, G. *et al.* A greener path for the EU Common Agricultural Policy. *Science* **365**, 449-451 (2019).
- 42 Finger, R. Nitrogen use and the effects of nitrogen taxation under consideration of production and price risks. *Agricultural Systems* **107**, 13-20 (2012).
- 43 Pretty, J. Intensification for redesigned and sustainable agricultural systems. *Science* **362**, eaav0294 (2018).
- 44 Holden, S. T. Fertilizer and sustainable intensification in Sub-Saharan Africa. *Global food security* **18**, 20-26 (2018).
- 45 Finger, R., Swinton, S. M., Benni, N. E. & Walter, A. Precision farming at the nexus of agricultural production and the environment. *Annual Review of Resource Economics* **11**, 1-23 (2019).
- 46 Walter, A., Finger, R., Huber, R. & Buchmann, N. Opinion: Smart farming is key to developing sustainable agriculture. *Proceedings of the National Academy of Sciences* **114**, 6148-6150 (2017).
- 47 Keele, L. J. & Titiunik, R. Geographic boundaries as regression discontinuities. *Political Analysis* **23**, 127-155 (2014).
- 48 Cattaneo, M. D. & Vazquez-Bare, G. The choice of neighborhood in regression discontinuity designs. *Observational Studies* **2**, A146 (2016).
- 49 SEDAC. *Gridded Population of the World (GPW)*, <<https://sedac.ciesin.columbia.edu/data/collection/gpw-v4>> (2019).
- 50 Gilbert, M. *et al.* Global distribution data for cattle, buffaloes, horses, sheep, goats, pigs, chickens and ducks in 2010. *Scientific Data* **5**, 180227, doi:10.1038/sdata.2018.227 (2018).
- 51 Wing, C. & Bello-Gomez, R. A. Regression discontinuity and beyond: Options for studying external validity in an internally valid design. *American Journal of Evaluation* **39**, 91-108 (2018).

1 52 Hansen, M. C. *et al.* High-resolution global maps of 21st-century forest cover change. *science*
2 **342**, 850-853 (2013).

3 53 Borrelli, P. *et al.* An assessment of the global impact of 21st century land use change on soil
4 erosion. *Nature Communications* **8**, 2013 (2017).

5 54 Carlson, K. *et al.* Greenhouse gas emissions intensity of global croplands. *Nature Climate*
6 *Change* **7**, 63, doi:10.1038/nclimate3158 (2016).

7 55 Leach, A. M. *et al.* A nitrogen footprint model to help consumers understand their role in
8 nitrogen losses to the environment. *Environmental Development* **1**, 40-66 (2012).

9 56 Gu, B. *et al.* The role of industrial nitrogen in the global nitrogen biogeochemical cycle.
10 *Scientific reports* **3**, 2579 (2013).

11 57 FAO. Fertistat: On-Line Database on fertilizer use by crop (2012).

12 58 Heffer, P. Assessment of fertilizer use by crop at the global level. *International Fertilizer*
13 *Industry Association, Paris, www.fertilizer.org/ifa/Home-Page/LIBRARY/Publication-*
14 *database.html/Assessment-of-Fertilizer-Use-by-Crop-at-the-Global-Level-2006-07-2007-08.*
15 *html2* (2009).

16 59 International Fertilizer Industry Association. Fertilizer use by crop. (Rome, 2002).

17 60 Monfreda, C., Ramankutty, N. & Foley, J. A. Farming the planet: 2. Geographic distribution of
18 crop areas, yields, physiological types, and net primary production in the year 2000. *Global*
19 *biogeochemical cycles* **22** (2008).

20 61 Hijmans, R., Cameron, S., Parra, J., Jones, P. & Jarvis, A. Very high resolution interpolated global
21 terrestrial climate surfaces. *International Journal of Climatology* **25**, 1965-1978 (2005).

22 62 Licker, R. *et al.* Mind the gap: how do climate and agricultural management explain the 'yield
23 gap' of croplands around the world? *Global ecology and biogeography* **19**, 769-782 (2010).

24 63 FAO. *Global Livestock Densities*, <[http://www.fao.org/livestock-systems/global-](http://www.fao.org/livestock-systems/global-distributions/en/)
25 [distributions/en/](http://www.fao.org/livestock-systems/global-distributions/en/)> (2012).

26 64 Dentener, F. *et al.* The global atmospheric environment for the next generation. *Environmental*
27 *Science & Technology* **40**, 3586-3594 (2006).

28 65 Lesschen, J., Stoorvogel, J., Smaling, E., Heuvelink, G. & Veldkamp, A. A spatially explicit
29 methodology to quantify soil nutrient balances and their uncertainties at the national level.
30 *Nutrient Cycling in Agroecosystems* **78**, 111-131 (2007).

31 66 Liu, J. *et al.* A high-resolution assessment on global nitrogen flows in cropland. *Proceedings of*
32 *the National Academy of Sciences* **107**, 8035-8040 (2010).

33 67 Bouwman, A., Boumans, L. & Batjes, N. Estimation of global NH₃ volatilization loss from
34 synthetic fertilizers and animal manure applied to arable lands and grasslands. *Global*
35 *Biogeochemical Cycles* **16**, 8-1-8-14 (2002).

36 68 Bouwman, A., Boumans, L. & Batjes, N. Modeling global annual N₂O and NO emissions from
37 fertilized fields. *Global Biogeochemical Cycles* **16**, 28-21-28-29 (2002).

38 69 Batjes, N. H. ISRIC-WISE derived soil properties on a 5 by 5 arc-minutes global grid (ver. 1.2).
39 (ISRIC-World Soil Information, 2012).

40 70 Allen, R. G., Pereira, L. S., Raes, D. & Smith, M. Crop evapotranspiration-Guidelines for
41 computing crop water requirements-FAO Irrigation and drainage paper 56. *Fao, Rome* **300**,
42 D05109 (1998).

43 71 Mitchell, T. D. & Jones, P. D. An improved method of constructing a database of monthly
44 climate observations and associated high-resolution grids. *International Journal of*
45 *Climatology: A Journal of the Royal Meteorological Society* **25**, 693-712 (2005).

46 72 Potter, P., Ramankutty, N., Bennett, E. M. & Donner, S. D. Characterizing the spatial patterns
47 of global fertilizer application and manure production. *Earth interactions* **14**, 1-22 (2010).

48 73 Smil, V. Nitrogen in crop production: An account of global flows. *Global biogeochemical cycles*
49 **13**, 647-662 (1999).

50 74 Dentener, F. *et al.* Nitrogen and sulfur deposition on regional and global scales: A multimodel
51 evaluation. *Global biogeochemical cycles* **20** (2006).

- 75 USDA. Agricultural waste management field handbook. *US Department of Agriculture: Washington, DC, USA* (1999).
- 76 Breiman, L. Random Forests. *Machine Learning* **45**, 5-32, doi:10.1023/a:1010933404324 (2001).
- 77 Gorelick, N. *et al.* Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment* **202**, 18-27, doi:<https://doi.org/10.1016/j.rse.2017.06.031> (2017).
- 78 Wuepper, D., Le Clech, S., Zilberman, D., Mueller, N. & Finger, R. Data: Countries Influence the Trade-Off between Crop Yields and Nitrogen Pollution. doi:<https://doi.org/10.3929/ethz-b-000430354> (2020).
- 79 Wuepper, D., Le Clech, S., Mueller, N., Zilberman, D. & Finger, R. Countries Influence the Trade-Off between Crop Yields and Nitrogen Pollution. *Nature Food* (2020).
- 80 World Bank. *World Bank Open Data*, <<https://data.worldbank.org/>> (2020).
- 81 Fulginiti, L. E., Perrin, R. K. & Yu, B. Institutions and agricultural productivity in Sub-Saharan Africa. *Agricultural Economics* **31**, 169-180 (2004).
- 82 Lio, M. & Liu, M.-C. Governance and agricultural productivity: A cross-national analysis. *Food Policy* **33**, 504-512 (2008).
- 83 FAO. *FAOSTAT*, <<http://www.fao.org/faostat/en/#home>> (2020).
- 84 Ceddia, M. G. The super-rich and cropland expansion via direct investments in agriculture. *Nature sustainability* **3**, 312-318 (2020).
- 85 McArthur, J. W. & McCord, G. C. Fertilizing growth: Agricultural inputs and their effects in economic development. *Journal of development economics* **127**, 133-152 (2017).
- 86 Chaudhary, A., Gustafson, D. & Mathys, A. Multi-indicator sustainability assessment of global food systems. *Nature communications* **9**, 848 (2018).