Estimating mixing processes of sources contributing to baseflow in Alpine headwater catchments

Other Conference Item

Author(s):

Floriancic, Marius ; Roques, Clément; Jimenez-Martinez, Joaquin

Publication date:

2020-05

Permanent link:

<https://doi.org/10.3929/ethz-b-000463221>

Rights / license:

[Creative Commons Attribution 4.0 International](#)

Originally published in:

EGUsphere, <https://doi.org/10.5194/egusphere-egu2020-1721>

EGU2020-1721

<https://doi.org/10.5194/egusphere-egu2020-1721>

EGU General Assembly 2020

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.

Estimating mixing processes of sources contributing to baseflow in Alpine headwater catchments

Marius G. Floriancic^{1,2}, Clément Roques³, and Joaquin Jimenez-Martinez^{2,4}

¹ETH Zürich, Institute of Terrestrial Ecosystems, Zürich, Switzerland (floriancic@ifu.baug.ethz.ch)

²ETH Zürich, Institute of Environmental Engineering, Zürich, Switzerland

³ETH Zürich, Geological Institute, Zürich, Switzerland

⁴EAWAG, Department of Water Resources and Drinking Water, Dübendorf, Switzerland

Baseflow is fed by groundwater to a large fraction. Estimating water quantity and quality from groundwater stores is essential for water management. However, there are few datasets available that contain detailed water chemistry analysis on high spatial resolution across multiple headwater catchments in (high) Alpine environments. This information is essential to analyze mixing processes on catchment scale from distinct landscape features.

We use two data sets: i) water chemistry analysis snapshot sampling campaigns in 7 headwater streams during low-flow periods across Switzerland, and ii) a detailed chemical screening (every 25 m) in one selected catchment during baseflow, including electric conductivity and temperature. Major ions, stable isotopes, TOC, DOC, trace elements were analyzed for some of the samples (> 80). These data reveal the chemical fingerprint of the contributing groundwater sources. The chemical composition of these contributing sources to baseflow is largely influenced by weathering products depended on lithology and geomorphology. Using maximum likelihood calculations, we define the ion composition and the isotopic signature of the potential major endmembers (up to three), based on the mixed samples along the main stem.

The proposed methodology allows to i) reduce uncertainty of the endmembers, and ii) quantify the relative contribution of different lithology and geomorphological features to streamflow and shows iii) which spatial scale of input information is needed to analyze mixing processes from various groundwater sources. Our results show how the contribution of different lithologies, along with topography and geomorphological features, varies spatially throughout Alpine headwater catchments.