Semi-classical dynamics in quantum spin systems

Journal Article**Author(s):**

Froehlich, Juerg; Knowles, Antti; Lenzmann, Enno

Publication date:

2007-12

Permanent link:

<https://doi.org/10.3929/ethz-b-000058650>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Originally published in:

Letters in Mathematical Physics 82(2-3), <https://doi.org/10.1007/s11005-007-0202-y>

Semi-Classical Dynamics in Quantum Spin Systems

JÜRGEN FRÖHLICH¹, ANTTI KNOWLES¹ and ENNO LENZMANN²

¹*Institute of Theoretical Physics, ETH Hönggerberg, 8093 Zürich, Switzerland.*

e-mail: juerg@itp.phys.ethz.ch; aknowles@itp.phys.ethz.ch

²*Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139-4307, USA. e-mail: lenzmann@math.mit.edu*

Received: 2 October 2007 / Revised: 26 October 2007 / Accepted: 31 October 2007

Published online: 20 November 2007

Abstract. We consider two limiting regimes, the large-spin and the mean-field limit, for the dynamical evolution of quantum spin systems. We prove that, in these limits, the time evolution of a class of quantum spin systems is determined by a corresponding Hamiltonian dynamics of classical spins. This result can be viewed as a Egorov-type theorem. We extend our results to the thermodynamic limit of lattice spin systems and continuum domains of infinite size, and we study the time evolution of coherent spin states in these limiting regimes.

Mathematics Subject Classification (2000). 35Q40, 82C20.

Keywords. Mean-field limit, Quantum spin systems, Landau-Lifschitz equation, Egorov theorem.

1. Introduction

The purpose of this letter is to study classical limits of quantum spin systems. Work in this direction was undertaken already at the beginning of the seventies. Many of the mathematical results established so far only concern time-independent aspects, such as the classical limit of quantum partition functions for spin systems [4,6]. Here we consider the dynamical evolution of quantum spin systems in limiting regimes; see [5] as well as [8,9] for earlier results. In particular, we discuss (i) *the large-spin limit* and (ii) *the mean-field/continuum limit*. As our main results, we prove that the time evolution of a large class of quantum spin systems approaches the time evolution of classical spins. Our results can be regarded as *Egorov-type theorems*, asserting that quantization commutes with time evolution in the classical limit; see [1] for a similar result on classical and quantum Bose gases. Along the way, we discuss thermodynamic limits and the time evolution of coherent spin states, in the two limits mentioned above.

An example of an evolution equation for classical spins is the *Landau-Lifshitz equation*

$$\partial_t M = M \wedge H_{\text{ex}}(M), \tag{1}$$

which is widely used in the study of ferromagnetism. Here $M = M(t, x) \in \mathbb{S}^2$ denotes a classical spin field with values on the unit sphere, and $\wedge$ stands for the vector product in $\mathbb{R}^3$. A standard choice for the exchange field is $H_{\text{ex}}(M) = J \Delta M$, where Δ denotes the Laplacian and J is the exchange coupling constant. Equation (1) then becomes

$$\partial_t M = J M \wedge \Delta M. \quad (2)$$

This form of the Landau-Lifshitz equation has been studied in the mathematical literature; see for instance [2, 3] and references given there. In physical terms, (2) describes the dynamics of spin waves in a ferromagnet with nearest neighbor exchange interactions in a classical regime; see [7].

In this paper we consider the Landau-Lifshitz equation with an exchange field $H_{\text{ex}}(M)$ given by an integral operator applied to M , and generalizations thereof. Equation (1) then takes the form

$$\partial_t M(t, x) = M(t, x) \wedge \int J(x, y) M(t, y) dy. \quad (3)$$

The integral kernel $J(x, y)$ describes the exchange interactions between classical spins beyond the nearest-neighbor approximation in the continuum limit. A formal argument on how to derive (2) from (3) is given in a remark in Section 3.4.

Our paper is organized as follows. In Section 2, we study the dynamics of finite lattice systems of quantum spins in the limit where their spin s approaches ∞ . The main result of Section 2 is formulated in Theorem 1 below. In order to prepare the ground for this theorem and its proof, we first introduce a class of Hamilton functions for classical spins and define their quantization by means of a normal-ordering prescription. At the end of Section 2, we pass to the thermodynamic limit, and we discuss the time evolution of coherent spin states.

In Section 3, we present a similar analysis for the mean-field limit of quantum spin systems defined on a lattice with spacing $h > 0$ in the continuum limit, $h \rightarrow 0$. The main result of Section 3 is stated in Theorem 4.

2. Large-Spin Limit

2.1. A SYSTEM OF CLASSICAL SPINS

Let Λ be a finite subset of the lattice $\mathbb{Z}^d$ (or any other lattice). A classical spin system on Λ is described in terms of the finite-dimensional phase space

$$\Gamma_\Lambda := \prod_{x \in \Lambda} \mathbb{S}^2,$$

i.e. we associate an element $M(x)$ of the unit two-sphere $\mathbb{S}^2 \subset \mathbb{R}^3$ with each site $x \in \Lambda$. The phase space Γ_Λ is conveniently coordinatized as follows. For each site $x \in \Lambda$, let $(M_1(x), M_2(x), M_3(x))$ denote the three cartesian components of a

unit vector $M(x) \in \mathbb{S}^2$, and define the complex coordinate functions $(M_+(x), M_z(x), M_-(x))$ on $\mathbb{S}^2$ by

$$M_{\pm}(x) := \frac{M_1(x) \pm iM_2(x)}{\sqrt{2}}, \quad M_z(x) := M_3(x).$$

We define a Poisson structure¹ on Γ_{Λ} by setting

$$\{M_i(x), M_j(y)\} = i\tilde{\varepsilon}_{ijk} \delta(x, y) M_k(x). \quad (4)$$

Here $\delta(x, y)$ stands for the Kronecker delta, and the indices i, j, k run through the index set $I := \{+, z, -\}$, where the symbol $\tilde{\varepsilon}_{ijk}$ is defined as $\tilde{\varepsilon}_{\pm\mp z} = \pm 1$, $\tilde{\varepsilon}_{\pm z \pm} = \mp 1$, $\tilde{\varepsilon}_{z \pm \pm} = \pm 1$, and $\tilde{\varepsilon}_{ijk} = 0$ otherwise.

For our purposes it is convenient (but not necessary) to replace $\mathbb{S}^2$ with the closed unit ball $\overline{B_1(0)} \subset \mathbb{R}^3$. To this end, we introduce a larger “phase space” (a Poisson manifold)

$$\Xi_{\Lambda} := \prod_{x \in \Lambda} \overline{B_1(0)},$$

equipped with the l^{∞} -norm. The algebra

$$\mathfrak{P}_{\Lambda} := \mathbb{C}[\{M_i(x) : i \in I, x \in \Lambda\}]$$

of complex polynomials is a Poisson algebra with Poisson bracket determined by (4). We equip $\mathfrak{P}_{\Lambda}$ with the norm $\|A\|_{\infty} := \sup_{M \in \Xi_{\Lambda}} |A(M)|$, and we denote its norm closure by $\mathfrak{A}_{\Lambda}$. Note that, by the Stone-Weierstrass theorem, $\mathfrak{A}_{\Lambda}$ is the algebra of continuous complex-valued functions on Ξ_{Λ} .

A fairly general class of Hamilton functions H_{Λ} on Ξ_{Λ} may be described as follows: With each multi-index $\alpha \in \mathbb{N}^{I \times \mathbb{Z}^d}$ satisfying $|\alpha| := \sum_{x \in \mathbb{Z}^d} \sum_{i \in I} \alpha_i(x) < \infty$ we associate a complex number $V(\alpha)$. Using the trivial embedding $\mathbb{N}^{I \times \Lambda} \subset \mathbb{N}^{I \times \mathbb{Z}^d}$ defined by adjoining zeroes, we consider Hamilton functions of the form

$$H_{\Lambda} := \sum_{\alpha \in \mathbb{N}^{I \times \Lambda}} V(\alpha) M^{\alpha}. \quad (5)$$

In order to obtain a real-valued H_{Λ} , we require that $\overline{V(\alpha)} = V(\bar{\alpha})$, where the “conjugation” $\bar{\alpha}$ of a multi-index α is defined as $\bar{\alpha}_i(x) := \alpha_{\bar{i}}(x)$, with $\bar{\cdot} : (+, z, -) \mapsto (-, z, +)$. Furthermore, we impose the following bound on the interaction potential:²

$$\|V\| := \sum_{n \in \mathbb{N}} \sup_{x \in \mathbb{Z}^d} \sum_{\substack{\alpha \in \mathbb{N}^{I \times \mathbb{Z}^d} \\ |\alpha| = n}} |\alpha(x)| |V(\alpha)| e^n < \infty. \quad (6)$$

¹Actually Γ_{Λ} is symplectic, with symplectic structure determined by the usual one on $\mathbb{S}^2$.

²Note that this condition may be weakened by replacing e^n with e^{rn} , for any $r > 0$. It may be checked that this does not affect the following results.

It is then easy to see that the series (5) converges in norm and that the set of allowed interaction potentials V is a Banach space. The Hamiltonian equation of motion reads $\dot{A} = \{H_\Lambda, A\}$, for any observable $A \in \mathfrak{A}_\Lambda$. In particular, a straightforward calculation yields

$$\frac{d}{dt} M_i(t, x) = \sum_{\alpha \in \mathbb{N}^I \times \Lambda} V(\alpha) \sum_{j, k} i \tilde{\varepsilon}_{jik} \alpha_j(x) M^{\alpha - \delta_j(x) + \delta_k(x)}(t), \quad (7)$$

where the multi-index $\delta_i(x)$ is defined by $[\delta_i(x)]_j(y) := \delta_{ij} \delta(x, y)$.

We record the following well-posedness result for the dynamics generated by the class of Hamiltonians introduced above.

LEMMA 1. *Let Λ be a (possibly infinite) subset of $\mathbb{Z}^d$. Let $M_0 \in \Xi_\Lambda$. Then the Hamiltonian equation (7) has a unique global-in-time solution $M \in C^1(\mathbb{R}, \Xi_\Lambda)$ that satisfies $M(0) = M_0$. Moreover, the solution M depends continuously on the initial condition M_0 , and we have the pointwise conservation law $|M(t, x)| = |M(0, x)|$ for all t .*

Proof. Local-in-time existence and uniqueness follows from a simple contraction mapping argument for the integral equation associated with (7). We omit the details. Also, continuous dependence on M_0 follows from standard arguments. Finally, the claim that $|M(0, x)| = |M(t, x)|$ for all t can be easily verified by using (7), which implies that $\frac{d}{dt} M(t, x)$ is perpendicular to $M(t, x)$. $\square$

Remarks. 1. In what follows, we denote the flow map $M_0 \mapsto M(t)$ by ϕ_Λ^t . Note that, under our assumptions, (7) also makes sense for infinite $\Lambda \subset \mathbb{Z}^d$, whereas the Hamiltonian H_Λ does not have a limit when $|\Lambda| \rightarrow \infty$.

2. The last statement implies that the magnitude of each spin remains constant in time, i.e. the spins precess. In particular, if $M_0 \in \Gamma_\Lambda$, it follows that $M(t) \in \Gamma_\Lambda$ for all t . Mathematically, this is simply the statement that the symplectic leaves of the Poisson manifold Ξ_Λ remain invariant under the Hamiltonian flow.
3. Time-dependent potentials $V(t, \alpha)$ may be treated without additional complications, provided that the map $t \mapsto V(t)$ is continuous (in the above norm) and $\sup_x$ in (6) is replaced by $\sup_{x, t}$. The weaker assumption that $t \mapsto V(t, \alpha)$ is continuous for all α implies Lemma 1 with the slightly weaker statement that $M \in C(\mathbb{R}, \Xi_\Lambda)$ is a classical solution of (7).

EXAMPLE. Consider the Hamiltonian

$$H_\Lambda(t) = - \sum_{x \in \Lambda} h(t, x) \cdot M(x) - \frac{1}{2} \sum_{x, y \in \Lambda} J(x, y) M(x) \cdot M(y), \quad (8)$$

where $M(x) = (M_1(x), M_2(x), M_3(x))$. Here $h(t, x) \in \mathbb{R}^3$ is an “external magnetic field” satisfying $\sup_{t \in \mathbb{R}, x \in \mathbb{Z}^d} |h(t, x)| < \infty$. We also require the map $t \mapsto h(t, x)$ to

be continuous for all $x \in \mathbb{Z}^d$. The exchange coupling $J : \mathbb{Z}^d \times \mathbb{Z}^d \rightarrow \mathbb{R}$ is assumed to be symmetric and to satisfy $J(x, x) = 0$ for all x . Finally we assume, in accordance with condition (6), that $\sup_{x \in \mathbb{Z}^d} \sum_{y \in \mathbb{Z}^d} |J(x, y)| < \infty$. The corresponding equation of motion for $M(t, x)$ is given by

$$\frac{d}{dt} M(t, x) = M(t, x) \wedge \left[h(t, x) + \sum_{y \in \Lambda} J(x, y) M(t, y) \right], \quad (9)$$

i. e. the *Landau-Lifschitz equation* for a classical lattice spin system.

2.2. A SYSTEM OF QUANTUM SPINS

In this section we formulate the quantum analogue of the system of classical spins from the previous section. We associate with each point $x \in \mathbb{Z}^d$ a finite-dimensional Hilbert space $\mathcal{H}_x^s \equiv \mathcal{H}_x := \mathbb{C}^{2s+1}$ describing a quantum-mechanical spin of magnitude s . (Here, and in the following, we refrain from displaying the explicit s -dependence whenever it is not needed.) Furthermore, we associate with each finite set $\Lambda \subset \mathbb{Z}^d$ the product space $\mathcal{H}_\Lambda := \bigotimes_{x \in \Lambda} \mathcal{H}_x$, and we define the algebra $\widehat{\mathfrak{A}}_\Lambda$ as the algebra of (bounded) operators on $\mathcal{H}_\Lambda$, equipped with the operator norm $\|\cdot\|$.

The spins are represented on $\mathcal{H}_\Lambda$ by a family $\{\widehat{S}_i(x) : i = 1, 2, 3, x \in \Lambda\}$ of operators, where $\widehat{S}_i(x)$ is the i th generator of the spin- s -representation of $\mathfrak{su}(2)$ on $\mathcal{H}_x$, rescaled by s^{-1} . In analogy to the complex coordinatization of the classical phase space Γ_Λ in the previous section, we replace the operators $(\widehat{S}_1(x), \widehat{S}_2(x), \widehat{S}_3(x))$ with $(\widehat{S}_+(x), \widehat{S}_z(x), \widehat{S}_-(x))$ as follows:

$$\widehat{S}_\pm(x) := \frac{\widehat{S}_1(x) \pm i\widehat{S}_2(x)}{\sqrt{2}}, \quad \widehat{S}_z(x) := \widehat{S}_3(x), \quad \text{for all } x \in \mathbb{Z}^d.$$

An easy calculation yields $\|\widehat{S}_\pm(x)\| \leq 1$ and $\|\widehat{S}_z(x)\| = 1$ if $s \geq 1$, and $\|\widehat{S}_\pm(x)\| = \sqrt{2}$ and $\|\widehat{S}_z(x)\| = 1$ if $s = 1/2$. Furthermore one finds the fundamental commutation relations

$$[\widehat{S}_i(x), \widehat{S}_j(y)] = \frac{1}{s} \varepsilon_{ijk} \delta(x, y) \widehat{S}_k(x) \quad (10)$$

with $i, j, k \in I$.

2.3. QUANTIZATION

In order to quantize polynomials in $\mathfrak{P}_\Lambda$ we need a concept of normal ordering. We say that a monomial $\widehat{S}_{i_1}(x_1) \cdots \widehat{S}_{i_p}(x_p)$ is *normal-ordered* if $i_k < i_l \Rightarrow k < l$, where $<$ is defined on I through $+\prec z \prec -$. We then define normal-ordering by

$$:\widehat{S}_{i_1}(x_1) \cdots \widehat{S}_{i_p}(x_p): = \widehat{S}_{i_{\sigma(1)}}(x_{\sigma(1)}) \cdots \widehat{S}_{i_{\sigma(p)}}(x_{\sigma(p)}),$$

where $\sigma \in S_p$ is a permutation such that the monomial on the right side is normal-ordered. Next, we define *quantization* $\widehat{\cdot} : \mathfrak{P}_\Lambda \rightarrow \widehat{\mathfrak{A}}_\Lambda$ by setting

$$(M_{i_1}(x_1) \cdots M_{i_p}(x_p))^\widehat{\cdot} = :\widehat{S}_{i_1}(x_1) \cdots \widehat{S}_{i_p}(x_p):$$

and by linearity of $\widehat{\cdot}$. We set $\widehat{1} = \mathbb{1}$. Note that, by definition, $\widehat{\cdot}$ is a linear map (but, of course, not an algebra homomorphism) and satisfies $(\widehat{A})^* = \widehat{A}$. It is also easy to see that, for $A, B \in \mathfrak{P}_\Lambda$, we have

$$[\widehat{A}, \widehat{B}] = -\frac{i}{s} \widehat{\{A, B\}} + O(s^{-2}),$$

so that s^{-1} is the deformation parameter of $\widehat{\cdot}$.

2.4. DYNAMICS IN THE LARGE-SPIN LIMIT

For each finite $\Lambda \subset \mathbb{Z}^d$ we define the Hamiltonian $\widehat{H}_\Lambda$ as the quantization of H_Λ . More precisely, we quantize (5) term by term and note that the resulting series converges in operator norm. Because H_Λ is real, the operator $\widehat{H}_\Lambda$ is self-adjoint on the finite-dimensional Hilbert space $\mathcal{H}_\Lambda$ and generates a one-parameter group of unitary propagators $U_s(t; \widehat{H}_\Lambda)$ (equal to $e^{-is\widehat{H}_\Lambda t}$ if $\widehat{H}_\Lambda$ is time-independent).

We introduce the short-hand notation

$$\begin{aligned} \alpha_\Lambda^t A &:= A \circ \phi_\Lambda^t, & A &\in \mathfrak{A}_\Lambda, \\ \widehat{\alpha}_\Lambda^t A &:= U_s(t; \widehat{H}_\Lambda)^* A U_s(t; \widehat{H}_\Lambda), & A &\in \widehat{\mathfrak{A}}_\Lambda, \end{aligned}$$

where ϕ_Λ^t is the Hamiltonian flow on Ξ_Λ . Note that both α_Λ^t and $\widehat{\alpha}_\Lambda^t$ are norm-preserving.

We are now able to state and prove our main result for the case of a finite lattice Λ . Roughly it states that time evolution and quantization commute in the $s \rightarrow \infty$ limit. This is a *Egorov-type result*.

THEOREM 1. *Let $A \in \mathfrak{P}_\Lambda$ and $\varepsilon > 0$. Then there exists a function $A(t) \in \mathfrak{P}_\Lambda$ such that*

$$\sup_{t \in \mathbb{R}} \|\alpha_\Lambda^t A - A(t)\|_\infty \leq \varepsilon, \tag{11}$$

and, for any $t \in \mathbb{R}$,

$$\|\widehat{\alpha}_\Lambda^t \widehat{A} - \widehat{A(t)}\| \leq \varepsilon + \frac{C(\varepsilon, t, A)}{s}, \tag{12}$$

where $C(\varepsilon, t, A)$ is independent of Λ .

Proof. Without loss of generality we assume that $A = M^\beta$ for some $\beta \in \mathbb{N}^{I \times \Lambda}$. For simplicity of notation we also assume, here and in the following proofs, that

H_Λ is time-independent. Consider the Lie-Schwinger series for the time evolution of the classical spin system,

$$\sum_{l=0}^{\infty} \frac{t^l}{l!} \{H_\Lambda, A\}^{(l)}, \quad (13)$$

where $\{H_\Lambda, A\}^{(l)} = \{H_\Lambda, \{H_\Lambda, A\}^{(l-1)}\}$ and $\{H_\Lambda, A\}^{(0)} = A$. In order to compute the nested Poisson brackets we observe that

$$\{M^\alpha, M^\beta\} = \sum_{x \in \Lambda} \sum_{i, j, k \in I} i \tilde{\varepsilon}_{ijk} \alpha_i(x) \beta_j(x) M^{\alpha+\beta-\delta_i(x)-\delta_j(x)+\delta_k(x)}, \quad (14)$$

as can be seen after a short calculation. Iterating this identity yields

$$\begin{aligned} \{H_\Lambda, A\}^{(l)} = & i^l \sum_{\alpha^1, \dots, \alpha^l} \sum_{x_1, \dots, x_l} \sum_{i_1, \dots, i_l} \sum_{j_1, \dots, j_l} \sum_{k_1, \dots, k_l} \times \\ & \times \left[\prod_{q=1}^l \tilde{\varepsilon}_{i_q j_q k_q} V(\alpha^q) \alpha_{i_q}^q(x_q) \left[\beta + \sum_{r=1}^{q-1} (\alpha^r - \delta_{i_r}(x_r) - \delta_{j_r}(x_r) + \delta_{k_r}(x_r)) \right]_{j_q}(x_q) \right] \times \\ & \times M^{\beta + \sum_{r=1}^l (\alpha^r - \delta_{i_r}(x_r) - \delta_{j_r}(x_r) + \delta_{k_r}(x_r))}. \end{aligned} \quad (15)$$

In order to estimate this series, we recall that $\|M^\gamma\|_\infty \leq 1$ and rewrite it by using that

$$\sum_{\alpha^1, \dots, \alpha^l} = \sum_{n_1, \dots, n_l=1}^{\infty} \sum_{|\alpha^1|=n_1} \cdots \sum_{|\alpha^l|=n_l}.$$

We then proceed recursively, starting with the sum over $\alpha^l, x_l, i_l, j_l, k_l$ and, at each step, using that

$$\sum_{|\alpha|=n} \sum_x \sum_{i, j, k} |\tilde{\varepsilon}_{ijk}| \alpha_i(x) \gamma_j(x) |V(\alpha)| \leq |\gamma| \|V\|^{(n)},$$

where

$$\|V\|^{(n)} := \sup_{x \in \mathbb{Z}^d} \sum_{|\alpha|=n} |V(\alpha)| |\alpha(x)|.$$

In this manner we find that

$$\begin{aligned} & \left\| \{H_\Lambda, A\}^{(l)} \right\|_\infty \\ & \leq \sum_{n_1, \dots, n_l=1}^{\infty} |\beta| (|\beta| + n_1) \cdots (|\beta| + n_1 + \cdots + n_{l-1}) \|V\|^{(n_1)} \cdots \|V\|^{(n_l)} \\ & \leq \sum_{n_1, \dots, n_l=1}^{\infty} (|\beta| + n_1 + \cdots + n_l)^l \|V\|^{(n_1)} \cdots \|V\|^{(n_l)} \end{aligned}$$

$$\begin{aligned}
&\leq l! \sum_{n_1, \dots, n_l=1}^{\infty} e^{|\beta|+n_1+\dots+n_l} \|V\|^{(n_1)} \dots \|V\|^{(n_l)} \\
&= l! e^{|\beta|} \|V\|^l.
\end{aligned}$$

Thus, for $|t| < \|V\|^{-1}$, the series (13) converges in norm, and an analogous estimate of the remainder of the Lie-Schwinger expansion of $\alpha_\Lambda^t A$ shows that (13) equals $\alpha_\Lambda^t A$. As all estimates are independent of Λ , the convergence is uniform in Λ .

The quantum-mechanical case is similar. Consider the Lie-Schwinger series for the time evolution of the quantum spin system:

$$\sum_{l=0}^{\infty} \frac{t^l}{l!} (is)^l [\widehat{H}_\Lambda, \widehat{A}]^{(l)}, \quad (16)$$

where $[\widehat{H}_\Lambda, \widehat{A}]^{(l)} = [\widehat{H}_\Lambda, [\widehat{H}_\Lambda, \widehat{A}]^{(l-1)}]$ and $[\widehat{H}_\Lambda, \widehat{A}]^{(0)} = \widehat{A}$. In order to estimate the multiple commutators, we remark that, from (4) and (10) and since both $\{\cdot, \cdot\}$ and $is[\cdot, \cdot]$ are derivations in both arguments, we see that $(is)^l [\widehat{H}_\Lambda, \widehat{A}]^{(l)}$ is equal to the expression obtained from $\{H_\Lambda, A\}^{(l)}$ by reordering the terms appropriately and by replacing $M_i(x)$ with $\widehat{S}_i(x)$. In particular (assuming $s \geq 1$)

$$\|(is)^l [\widehat{H}_\Lambda, \widehat{A}]^{(l)}\| \leq l! e^\beta \|V\|^l,$$

and we deduce exactly as above that (16) equals $\widehat{\alpha}_\Lambda^t \widehat{A}$ for $t < \|V\|^{-1}$.

To show the claim of the theorem for $|t| < \|V\|^{-1}$ we first remark that $\widehat{\alpha}_\Lambda^t(A)$ is well-defined through its convergent power series expansion. Now as shown above, each term of $(is)^l [\widehat{H}_\Lambda, \widehat{A}]^{(l)}$, as a polynomial in $\widehat{\mathfrak{A}}_\Lambda$, is equal to a reordering of the corresponding term of $\{H_\Lambda, A\}^{(l)}$. If P is a monomial (with coefficient 1) of degree p in the generating variables $\{\widehat{S}_i(x)\}$ and $\tilde{P}$ a monomial obtained from P by any reordering of terms, the commutation relations (10) imply that

$$\|P - \tilde{P}\| \leq \frac{p^2}{s}.$$

Thus

$$\left(\{H_\Lambda, A\}^{(l)}\right)^\wedge = (is)^l [\widehat{H}_\Lambda, \widehat{A}]^{(l)} + R_l,$$

where, recalling the expression (15) and the estimates following it, we see that the “loop terms” R_l are bounded by

$$\begin{aligned}
\|R_l\| &\leq \frac{1}{s} \sum_{n_1, \dots, n_l=1}^{\infty} (|\beta| + n_1 + \dots + n_l)^{l+2} \|V\|^{(n_1)} \dots \|V\|^{(n_l)} \\
&\leq \frac{(l+2)!}{s} e^{|\beta|} \|V\|^l.
\end{aligned}$$

Therefore, if $|t| < \|V\|^{-1}$,

$$\|\widehat{\alpha}'_{\Lambda} \widehat{A} - \widehat{\alpha''}_{\Lambda} \widehat{A}\| \leq \frac{e^{|\beta|}}{s} \sum_{l=0}^{\infty} (l+2)(l+1)(t\|V\|)^l \leq \frac{C(t, A)}{s},$$

where $C(t, A)$ is independent of Λ .

In order to extend the result to arbitrary times we proceed by iteration. The crucial observations that enable this process are that the convergence radius $\|V\|^{-1}$ is independent of $|\beta|$ and $\alpha'_{\Lambda}, \widehat{\alpha}'_{\Lambda}$ are norm-preserving. Let $t \in \mathbb{R}$ and choose $v \in \mathbb{N}$ such that $\tau := t/v$ satisfies $|\tau| < \|V\|^{-1}$. In order to iterate we need to introduce a cutoff in the series (13) and (15). The series (13) consists of an infinite sum of terms in $\mathfrak{P}_{\Lambda}$ which are indexed by $(l, \alpha^1, \dots, \alpha^l, x_1, \dots, x_l)$. Now let $\varepsilon > 0$ be given. Since the series converges in norm there is a finite subset

$$B_1 = B_1(\varepsilon) \subset \{(l, \alpha^1, \dots, \alpha^l, x_1, \dots, x_l)\} = \bigcup_{l=0}^{\infty} (\mathbb{N}^{I \times \Lambda})^l \times \Lambda^l$$

such that the norm of the series restricted to the complement of B_1 is smaller than ε/v . This induces a splitting $\alpha'_{\Lambda} A = \alpha_{B_1} A + \alpha_{B_1^c} A$ (in self-explanatory notation), such that $\alpha_{B_1} A \in \mathfrak{P}_{\Lambda}$ and $\|\alpha_{B_1^c} A\|_{\infty} \leq \varepsilon/v$. Similarly, one splits $\widehat{\alpha}'_{\Lambda} \widehat{A} = \widehat{\alpha}_{B_1} \widehat{A} + \widehat{\alpha}_{B_1^c} \widehat{A}$ where, after an eventual increase of B_1 , $\|\widehat{\alpha}_{B_1^c} \widehat{A}\| \leq \varepsilon/v$.

Now we use the above result for $|\tau| < \|V\|^{-1}$:

$$\widehat{\alpha}_{\Lambda}^{\tau} \widehat{A} = \widehat{\alpha}_{B_1} \widehat{A} + \frac{R_1}{s} + \widehat{\alpha}_{B_1^c} \widehat{A}$$

where R_1 is some bounded operator. Since $\alpha_{B_1} A \in \mathfrak{P}_{\Lambda}$ we may repeat the process on the time interval $[\tau, 2\tau]$:

$$\begin{aligned} \widehat{\alpha}_{\Lambda}^{\tau} \widehat{\alpha}_{\Lambda}^{\tau} \widehat{A} &= \widehat{\alpha}_{\Lambda}^{\tau} \widehat{\alpha}_{B_1} \widehat{A} + \frac{\widehat{\alpha}_{\Lambda}^{\tau} R_1}{s} + \widehat{\alpha}_{\Lambda}^{\tau} \widehat{\alpha}_{B_1^c} \widehat{A} \\ &= (\alpha_{B_2} \alpha_{B_1} A)^{\widehat{}} + \widehat{\alpha}_{B_2^c} \widehat{\alpha}_{B_1} \widehat{A} + \widehat{\alpha}_{\Lambda}^{\tau} \widehat{\alpha}_{B_1^c} \widehat{A} + \frac{R_2 + \widehat{\alpha}_{\Lambda}^{\tau} R_1}{s} \end{aligned}$$

Continuing in this manner one sees that, since α'_{Λ} and $\widehat{\alpha}'_{\Lambda}$ are norm-preserving, $A(t) := \alpha_{B_v} \cdots \alpha_{B_1} A \in \mathfrak{P}_{\Lambda}$ satisfies

$$\|\alpha'_{\Lambda} A - A(t)\|_{\infty} \leq \varepsilon,$$

as well as

$$\|\widehat{\alpha}'_{\Lambda} \widehat{A} - \widehat{A}(t)\| \leq \varepsilon + \frac{C(\varepsilon, t, A)}{s}.$$

□

2.5. THE THERMODYNAMIC LIMIT

The above analysis was done for a finite subset Λ , but the observed uniformity in Λ allows for a statement of the result directly in limit $\Lambda = \mathbb{Z}^d$. We pause to describe how this works.

Concentrate first on the quantum case. If $\Lambda_1 \subset \Lambda_2$, an operator $A_1 \in \widehat{\mathfrak{A}}_{\Lambda_1}$ may be identified in the usual fashion with an operator $A_2 \in \widehat{\mathfrak{A}}_{\Lambda_2}$ by setting $A_2 = A_1 \otimes \mathbb{1}_{\Lambda_2 \setminus \Lambda_1}$. We shall tacitly make use of this identification in the following. It induces the norm-preserving mapping $\widehat{\mathfrak{A}}_{\Lambda_1} \rightarrow \widehat{\mathfrak{A}}_{\Lambda_2}$ of the abstract C^* -algebras and the isotony relation $\widehat{\mathfrak{A}}_{\Lambda_1} \subset \widehat{\mathfrak{A}}_{\Lambda_2}$. Observables of the quantum spin system in the thermodynamic limit are then elements of the *quasi-local algebra*

$$\widehat{\mathfrak{A}} := \overline{\bigvee_{\Lambda \subset \mathbb{Z}^d \text{ finite}} \widehat{\mathfrak{A}}_{\Lambda}},$$

which is the C^* -algebra defined as the closure of the normed algebra generated by the union of all $\widehat{\mathfrak{A}}_{\Lambda}$'s, where Λ is finite. The spins are represented on $\widehat{\mathfrak{A}}$ by a family $\{\widehat{S}_i(x) : i \in I, x \in \mathbb{Z}^v\}$ of operators.

The dynamics of the system is determined by a one-parameter group $(\widehat{\alpha}^t)_{t \in \mathbb{R}}$ of automorphisms of $\widehat{\mathfrak{A}}$. Its existence is a corollary of the proof of Theorem 1.

LEMMA 2. *Let $A \in \widehat{\mathfrak{A}}_{\Lambda_0}$ for some finite $\Lambda_0 \subset \mathbb{Z}^d$ and $t \in \mathbb{R}$. Then the following limit exists in the norm sense:*

$$\lim_{\Lambda \rightarrow \infty} \widehat{\alpha}_{\Lambda}^t A =: \widehat{\alpha}^t A,$$

where $\Lambda \rightarrow \infty$ means that Λ eventually contains every finite subset. By continuity this extends to a strongly continuous one-parameter group $(\widehat{\alpha}^t)_{t \in \mathbb{R}}$ of automorphisms of $\widehat{\mathfrak{A}}$.

Proof. For $|t| < \|V\|^{-1}$ the series (16) is bounded in norm, uniformly in Λ , so to show convergence of the series it suffices to show the convergence of $[\widehat{H}_{\Lambda}, \widehat{A}]^{(l)}$ for each $l \in \mathbb{N}$, which is an easy exercise.

Thus $\widehat{\alpha}^t A$ is well-defined for any polynomial A . By continuity, $\widehat{\alpha}^t$ extends to an automorphism of $\widehat{\mathfrak{A}}$. Since $\widehat{\alpha}^t A \in \widehat{\mathfrak{A}}$ and $\widehat{\alpha}^t$ is a one-parameter group, we may extend it to all times by iteration. Strong continuity follows since $\widehat{\alpha}^t A$, for small t and polynomial A , is defined through a convergent power series:

$$\lim_{t \rightarrow 0} \|\widehat{\alpha}^t A - A\| = 0.$$

By continuity, this remains true for all $A \in \widehat{\mathfrak{A}}$. □

For classical spin systems and finite Λ we recall that $\mathfrak{A}_{\Lambda} = C\left(\prod_{x \in \Lambda} \overline{B_1(0)}; \mathbb{C}\right)$, a C^* -algebra under $\|\cdot\|_{\infty}$. As above, for $\Lambda_1 \subset \Lambda_2$, we identify $A_1 \in \mathfrak{A}_{\Lambda_1}$ with a function $A_2 \in \mathfrak{A}_{\Lambda_2}$ by setting $A_2 = A_1 \otimes \mathbb{1}_{\Lambda_2 \setminus \Lambda_1}$. We thus get a norm-preserving mapping $\mathfrak{A}_{\Lambda_1} \rightarrow \mathfrak{A}_{\Lambda_2}$ of the abstract C^* -algebras and the relation $\mathfrak{A}_{\Lambda_1} \subset \mathfrak{A}_{\Lambda_2}$. Define the classical quasi-local algebra as

$$\mathfrak{A} := \overline{\bigvee_{\Lambda \subset \mathbb{Z}^d \text{ finite}} \mathfrak{A}_\Lambda}.$$

Note that $\mathfrak{A}$ is equal to the space of continuous complex functions on $\prod_{x \in \mathbb{Z}^d} \overline{B_1(0)}$, equipped with the product topology (this is an immediate consequence of the Tychonoff and Stone–Weierstrass theorems).

The spins are represented on $\mathfrak{A}$ by a family $\{M_i(x) : i \in I, x \in \mathbb{Z}^v\}$ of functions. Existence of the dynamics follows exactly as above.

LEMMA 3. *Let $A \in \mathfrak{A}_{\Lambda_0}$ for some finite $\Lambda_0 \subset \mathbb{Z}^d$ and $t \in \mathbb{R}$. Then the following limit exists in $\|\cdot\|_\infty$:*

$$\lim_{\Lambda \rightarrow \infty} \alpha_\Lambda^t A =: \alpha^t A.$$

By continuity this extends to a strongly continuous one-parameter group $(\alpha^t)_{t \in \mathbb{R}}$ of automorphisms of $\mathfrak{A}$. Furthermore, $\alpha^t A = A \circ \phi^t$, where $\phi^t = \phi_{\mathbb{Z}^d}^t$ is the Landau–Lifschitz flow defined in the remark after Lemma 1.

Now set $\mathfrak{P} := \mathbb{C}[\{M_i(x) : i \in I, x \in \mathbb{Z}^d\}]$. Then the proof of Theorem 1 yields the following

THEOREM 2. *Let $A \in \mathfrak{P}$ and $\varepsilon > 0$. Then there exists a function $A(t) \in \mathfrak{P}$ such that*

$$\sup_{t \in \mathbb{R}} \|\alpha^t A - A(t)\|_\infty \leq \varepsilon, \quad (17)$$

and, for any $t \in \mathbb{R}$,

$$\|\widehat{\alpha^t A} - \widehat{A(t)}\| \leq \varepsilon + \frac{C(\varepsilon, t, A)}{s}. \quad (18)$$

Remark. In particular, the result applies to classical equations of motion of the form (9) where the sum over y ranges over $\mathbb{Z}^d$.

2.6. EVOLUTION OF COHERENT STATES

Denote by $S_i := s\widehat{S}_i$ the unscaled spin operator in the spin- s -representation of $\mathfrak{su}(2)$. For the polar angles $(\theta, \varphi) \in [0, \pi] \times [0, 2\pi)$ corresponding to the unit vector $M \in \mathbb{S}^2$ we define the coherent state in $\mathbb{C}^{2s+1}$ as

$$|M\rangle := \exp \frac{\theta}{\sqrt{2}} [e^{i\varphi} S_- - e^{-i\varphi} S_+] |s\rangle,$$

where $|s\rangle$ is the highest-weight state, i.e. $S_z |s\rangle = s |s\rangle$. Note that $|M\rangle = e^{i\alpha \cdot S} |s\rangle$, where $\alpha = \theta n$ and n is the unit vector $(\sin \varphi, -\cos \varphi, 0)$.

Set $A := \frac{\theta}{\sqrt{2}} [e^{i\varphi} S_- - e^{-i\varphi} S_+]$ and $U := e^A$ so that $|M\rangle = U|s\rangle$. Then using

$$U^* S_i U = \sum_{k=0}^{\infty} \frac{1}{k!} [\dots [S_i, A], \dots, A],$$

we find

$$\begin{aligned} U^* S_1 U &= \sin \theta \cos \varphi S_z + \frac{1}{\sqrt{2}} \cos^2 \frac{\theta}{2} (S_+ + S_-) - \frac{1}{\sqrt{2}} \sin^2 \frac{\theta}{2} (e^{-i\varphi} S_+ + e^{2i\varphi} S_-), \\ U^* S_2 U &= \sin \theta \sin \varphi S_z + \frac{1}{\sqrt{2}i} \cos^2 \frac{\theta}{2} (S_+ - S_-) - \frac{1}{\sqrt{2}i} \sin^2 \frac{\theta}{2} (e^{2i\varphi} S_+ - e^{-2i\varphi} S_-), \\ U^* S_3 U &= \cos \theta S_z - \frac{1}{\sqrt{2}} \sin \theta (e^{-i\varphi} S_+ + e^{i\varphi} S_-). \end{aligned} \quad (19)$$

As a consequence note that

$$\langle M, \widehat{S} M \rangle = \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix} = M. \quad (20)$$

In order to derive our main result for coherent spins states we need the following lemma, which follows from direct calculations.

LEMMA 4. *For any unit vector $M \in \mathbb{S}^2$, we have that*

$$|\langle M, \widehat{S}_{i_1} \cdots \widehat{S}_{i_p} M \rangle - M_{i_1} \cdots M_{i_p}| \leq p \sqrt{\frac{2}{s}}.$$

Now let $M: \mathbb{Z}^d \rightarrow \mathbb{S}^2$ be a configuration of classical spins on the lattice. Then M defines a state ρ_M on $\widehat{\mathfrak{A}}$ as follows. For finite Λ , consider the product state

$$|M_\Lambda\rangle := \bigotimes_{x \in \Lambda} |M(x)\rangle \in \mathcal{H}_\Lambda.$$

Then, for $\mathbf{A} \in \widehat{\mathfrak{A}}_\Lambda$, we set

$$\rho_M(\mathbf{A}) := \langle M_\Lambda, \mathbf{A} M_\Lambda \rangle,$$

and extend the definition of ρ_M to arbitrary $\mathbf{A} \in \widehat{\mathfrak{A}}$ by continuity.

Let $M: \mathbb{R} \times \mathbb{Z}^d \rightarrow \mathbb{S}^2$ be the solution of the Hamiltonian equation of motion (7) with initial conditions $M(0, x) = M(x)$. The following result links the quantum time evolution for coherent spin states with the corresponding classical configuration in the large-spin limit.

THEOREM 3. *Let $t \in \mathbb{R}$, $A \in \mathfrak{P}$ and $M: \mathbb{Z}^d \rightarrow \mathbb{S}^2$. Then*

$$\lim_{s \rightarrow \infty} \rho_M(\widehat{\alpha}^t \widehat{A}) = A(M(t)),$$

uniformly in t on compact time intervals.

Proof. The proof is essentially a corollary of Lemma 4 and the proof of Theorem 1. First, let $|t| < \|V\|^{-1}$. We know from (13) and (16) that

$$A(M(t)) = \sum_{l=0}^{\infty} \frac{t^l}{l!} \lim_{\Lambda \rightarrow \infty} \{H_{\Lambda}, A\}^{(l)}(M),$$

as well as

$$\rho_M(\widehat{\alpha}_{\Lambda}^t \widehat{A}) = \sum_{l=0}^{\infty} \frac{t^l}{l!} (is)^l \rho_M \left(\lim_{\Lambda \rightarrow \infty} [\widehat{H}_{\Lambda}, \widehat{A}]^{(l)} \right).$$

Now Lemma 4 implies that

$$|\rho_M(\widehat{M}^{\alpha}) - M^{\alpha}| \leq |\alpha| \sqrt{\frac{2}{s}}.$$

Arguing exactly as in the proof of Theorem 1, we get the bound

$$|\rho_M(\widehat{\alpha}^t \widehat{A}) - A(M(t))| \leq \frac{C(t, A)}{\sqrt{s}}. \quad (21)$$

Arbitrary times are reached by iteration as in the proof of Theorem 1. $\square$

3. Mean-Field Limit

This section is devoted to the dynamics of a quantum spin system in the mean-field/continuum limit. More precisely, we consider a system of quantum spins on a lattice with spacing $h > 0$. The limit $h \rightarrow 0$ yields again a Egorov-type result: The quantum dynamics approaches the dynamics of a classical spin system defined on a continuum set. As in the previous section, we also discuss the thermodynamic limit and the time evolution of coherent states.

3.1. A SYSTEM OF QUANTUM SPINS ON A LATTICE

Let $\Lambda \subset \mathbb{R}^d$ be bounded and open. We associate with each spacing $h > 0$ the finite lattice

$$\Lambda^{(h)} := h\mathbb{Z}^d \cap \Lambda.$$

At each lattice site $x \in \Lambda^{(h)}$ there is a spin of (fixed) magnitude s . The Hilbert space of this quantum system is

$$\mathcal{H}_{\Lambda}^{(h)} := \bigotimes_{x \in \Lambda^{(h)}} \mathbb{C}^{2s+1}.$$

The algebra of bounded operators on $\mathcal{H}_{\Lambda}^{(h)}$ is denoted by $\widehat{\mathfrak{A}}_{\Lambda}^{(h)}$.

The spins are represented on $\mathcal{H}_{\Lambda}^{(h)}$ by a family $\{\widehat{S}_i(x) : i = 1, 2, 3, x \in \Lambda^{(h)}\}$ of operators, where $\widehat{S}_i(x)$ is the i th generator of the spin- s -representation of $\mathfrak{su}(2)$,

rescaled by h^d/s . As usual, we replace the operators $(\widehat{S}_1, \widehat{S}_2, \widehat{S}_3)$ with $(\widehat{S}_+, \widehat{S}_z, \widehat{S}_-)$. They satisfy the bounds $\|\widehat{S}_\pm\| \leq h^d$ and $\|\widehat{S}_z\| = h^d$ if $s \geq 1$, as well as $\|\widehat{S}_\pm\| = \sqrt{2}h^d$ and $\|\widehat{S}_z\| = h^d$ if $s = 1/2$. The commutation relations now read

$$[\widehat{S}_i(x), \widehat{S}_j(y)] = \frac{h^d}{s} \tilde{\varepsilon}_{ijk} \delta(x, y) \widehat{S}_k(x), \quad (22)$$

with $i, j, k \in I$.

3.2. A CONTINUUM THEORY OF SPINS

Let $\Lambda \subset \mathbb{R}^d$ be a bounded, open set. A system of classical spins on Λ is represented in terms of the Poisson “phase space”³

$$\Xi_\Lambda := \left\{ M \in L^\infty(\Lambda; \mathbb{R}^3) : \|M\|_\infty \leq 1 \right\},$$

which we equip with the L^∞ -norm. In analogy to Section 2, we use the complex coordinates (M_+, M_z, M_-) instead of (M_1, M_2, M_3) , so that the Poisson bracket on Ξ_Λ satisfies

$$\{M_i(x), M_j(y)\} = i \tilde{\varepsilon}_{ijk} \delta(x - y) M_k(x), \quad (23)$$

for $i, j, k \in I$.

In order to describe a useful class of observables on Ξ_Λ , we introduce the space $\mathcal{B}^{(p)}$, $p \in \mathbb{N}$, which consists of all functions f in $C(\mathbb{R}^{pd}; \mathbb{C}^{3p})$ that are symmetric in their arguments, in the sense that $Pf = f$, where

$$(Pf)_{i_1 \dots i_p}(x_1, \dots, x_p) := \frac{1}{p!} \sum_{\sigma \in S_p} f_{i_{\sigma(1)} \dots i_{\sigma(p)}}(x_{\sigma(1)}, \dots, x_{\sigma(p)}).$$

On the space $\mathcal{B}^{(p)}$ we introduce the norms

$$\begin{aligned} \|f\|_1^{(h)} &:= h^{pd} \sum_{i_1, \dots, i_p \in I} \sum_{x_1, \dots, x_p \in h\mathbb{Z}^d} |f_{i_1 \dots i_p}(x_1, \dots, x_p)|, \\ \|f\|_{\infty, 1}^{(h)} &:= \sup_x \sum_{i_1, \dots, i_p \in I} h^{(p-1)d} \sum_{x_2, \dots, x_p \in h\mathbb{Z}^d} |f_{i_1 \dots i_p}(x, x_2, \dots, x_p)|. \end{aligned}$$

We shall be interested in observables arising from $f \in \mathcal{B}^{(p)}$ satisfying

$$\limsup_{h \rightarrow 0} \|f\|_1^{(h)} < \infty. \quad (24)$$

Note that Fatou’s lemma implies that $\|f\|_1 \leq \limsup_{h \rightarrow 0} \|f\|_1^{(h)}$.

³As in the previous section, one may introduce a symplectic phase space Γ_Λ consisting of all $M \in \Xi_\Lambda$ such that $|M(x)| = 1$ a.e.

We define $\mathfrak{P}_\Lambda$ as the “polynomial” algebra of functions on Ξ_Λ generated by functions of the form

$$M_\Lambda(f) := \sum_{i_1, \dots, i_p \in \Lambda} \int dx_1 \cdots dx_p f_{i_1 \dots i_p}(x_1, \dots, x_p) M_{i_1}(x_1) \cdots M_{i_p}(x_p),$$

where $f \in \mathcal{B}^{(p)}$ satisfies (24). $\mathfrak{P}_\Lambda$ is clearly a Poisson algebra. We equip it with the norm $\|A\|_\infty := \sup_{M \in \Xi_\Lambda} |A(M)|$ so that

$$\|M_\Lambda(f)\|_\infty \leq \|f\|_1. \quad (25)$$

3.3. QUANTIZATION

For $f \in \mathcal{B}^{(p)}$ let us define

$$\widehat{S}_\Lambda(f) := \sum_{i_1, \dots, i_p} \sum_{x_1, \dots, x_p \in \Lambda^{(h)}} f_{i_1 \dots i_p}(x_1, \dots, x_p) \widehat{S}_{i_1}(x_1) \cdots \widehat{S}_{i_p}(x_p). \quad (26)$$

If f satisfies (24), we find that

$$\|\widehat{S}_\Lambda(f)\| \leq \|f\|_1^{(h)}. \quad (27)$$

As above, quantization $\widehat{\cdot} : \mathfrak{P}_\Lambda \rightarrow \widehat{\mathfrak{A}}_\Lambda^{(h)}$ is defined by $\widehat{M_\Lambda(f)} = \widehat{S}_\Lambda(f)$ and linearity. Here $\widehat{\cdot}$ denotes the normal-ordering of the spin operators introduced above. Also, we set $\widehat{1} = \mathbb{I}$. Again, $(\widehat{A})^* = \widehat{\bar{A}}$. Furthermore one sees, as in the previous section, that $\hbar^d/s$ is the deformation parameter of $\widehat{\cdot}$.

3.4. DYNAMICS IN THE MEAN-FIELD LIMIT

We consider a family $V = (V^{(n)})_{n=1}^\infty$ of functions, where $V^{(n)} \in \mathcal{B}^{(p)}$ satisfies

$$\overline{V_{i_1 \dots i_n}^{(n)}}(x_1, \dots, x_n) = V_{\bar{i}_1 \dots \bar{i}_n}^{(n)}(x_1, \dots, x_n),$$

where, we recall, $\bar{\cdot}$ on I maps $(+, z, -)$ to $(-, z, +)$. We define the Hamilton function on Ξ_Λ through

$$H_\Lambda := \sum_{n=1}^\infty M_\Lambda(V^{(n)}). \quad (28)$$

Set

$$\|V\|^{(h)} := \sum_{n=1}^\infty n e^n \|V^{(n)}\|_{\infty, 1}^{(h)}. \quad (29)$$

We impose the condition $\|V\| := \limsup_{h \rightarrow 0} \|V\|^{(h)} < \infty$. In the continuum limit, we observe that

$$\sum_{n=1}^\infty n e^n \sup_x \sum_{i_1, \dots, i_n} \int dx_2 \cdots dx_n |V_{i_1 \dots i_n}^{(n)}(x, x_2, \dots, x_n)| \leq \|V\|, \quad (30)$$

as can be seen using Fatou's lemma. It now follows easily that, for each bounded set Λ , the sum (28) converges in $\|\cdot\|_\infty$ on Ξ_Λ and yields a well-defined real Hamilton function H_Λ .

The Hamiltonian equation of motion reads

$$\begin{aligned} \frac{d}{dt} M_i(t, x) = i \sum_{n=1}^{\infty} n \sum_{i_1, \dots, i_n, j} \int dx_2 \cdots dx_n V_{i_1 \dots i_n}^{(n)}(x, x_2, \dots, x_n) \times \\ \times \tilde{\varepsilon}_{i_1 i j} M_j(t, x) M_{i_2}(t, x_2) \cdots M_{i_n}(t, x_n). \end{aligned} \quad (31)$$

By standard methods, we find the following global well-posedness result for (31).

LEMMA 5. *Let $\Lambda \subset \mathbb{R}^d$ by any open subset of $\mathbb{R}^d$ and $M_0 \in \Xi_\Lambda$. Then (31) has a unique solution $M \in C^1(\mathbb{R}, \Xi_\Lambda)$ that satisfies $M(0) = M_0$. Moreover, the solution M depends continuously on the initial condition M_0 . Finally, we have the pointwise conservation law $|M(t, x)| = |M(0, x)|$ for all t .*

Remarks. 1. As in Section 2, we denote the norm-preserving Hamiltonian flow by ϕ_Λ^t .

2. Time-dependent potentials $V(t)$ may be treated exactly as in the previous section.

EXAMPLE. Consider

$$H_\Lambda = - \int_\Lambda dx h(t, x) \cdot M(x) - \frac{1}{2} \int_{\Lambda \times \Lambda} dx dy J(x, y) M(x) \cdot M(y),$$

which yields the Landau-Lifshitz equation of motion

$$\frac{d}{dt} M(t, x) = M(t, x) \wedge \left[h(t, x) + \int_\Lambda dy J(x, y) M(t, y) \right].$$

Remark. In a formal way, the Landau-Lifshitz equation (2) mentioned in the introduction can be obtained from (3) with $J = J(|x - y|)$ by Taylor expanding $M(t, y)$ up to second order in $y - x$. This leads to (2) after rescaling time by $t \mapsto \alpha t$ where $\alpha = \frac{1}{2d} \int J(|x|) |x|^2 dx$.

The quantum dynamics is generated by the Hamiltonian $\widehat{H}_\Lambda \in \widehat{\mathfrak{A}}_\Lambda^{(h)}$ defined as the quantization of H_Λ . More precisely, each term of H_Λ is quantized and it may be easily verified that the resulting series converges in operator norm. The fact that H_Λ is real immediately implies that $\widehat{H}_\Lambda$ is self-adjoint. As above we introduce the short-hand notation

$$\begin{aligned} \alpha_\Lambda^t A &:= A \circ \phi_\Lambda^t, \quad A \in \mathfrak{A}_\Lambda, \\ \widehat{\alpha}_\Lambda^t A &:= U_h(t; \widehat{H}_\Lambda)^* A U_h(t; \widehat{H}_\Lambda), \quad A \in \widehat{\mathfrak{A}}_\Lambda^{(h)}. \end{aligned}$$

Here, $U_h(t; \widehat{H}_\Lambda)$ is the quantum mechanical propagator, equal to $e^{ish^{-d}\widehat{H}_\Lambda t}$ if $\widehat{H}_\Lambda$ is time-independent.

We are now in a position to state our main result on the mean-field dynamics of the quantum system on the finite lattice $\Lambda^{(h)}$ in the continuum limit, as $h \rightarrow 0$.

THEOREM 4. *Let $\Lambda \subset \mathbb{R}^d$ be open and bounded, $A \in \mathfrak{P}_\Lambda$ and $\varepsilon > 0$. Then there exists a function $A(t) \in \mathfrak{P}_\Lambda$ such that*

$$\sup_{t \in \mathbb{R}} \|\alpha_\Lambda^t A - A(t)\|_\infty \leq \varepsilon, \quad (32)$$

and, for any $t \in \mathbb{R}$,

$$\|\widehat{\alpha}_\Lambda^t \widehat{A} - \widehat{A(t)}\| \leq \varepsilon + C(\varepsilon, t, A) h^d, \quad (33)$$

where $C(\varepsilon, t, A)$ is independent of Λ .

Proof. One finds, for $f \in \mathcal{B}^{(p)}$ and $g \in \mathcal{B}^{(q)}$,

$$\{M_\Lambda(f), M_\Lambda(g)\} = pq M_\Lambda(f \rightharpoonup g) \quad (34)$$

where $f \rightharpoonup g \in \mathcal{B}^{(p+q-1)}$ is defined by

$$(f \rightharpoonup g)_{i_1 \dots i_{p+q-1}}(x_1, \dots, x_{p+q-1}) := i P \sum_{i,j} \tilde{\varepsilon}_{ij i_1} f_{i i_2 \dots i_p}(x_1, \dots, x_p) g_{j i_{p+1} \dots i_{p+q-1}}(x_1, x_{p+1}, \dots, x_{p+q-1}). \quad (35)$$

We have the estimate

$$\|f \rightharpoonup g\|_1 \leq \|f\|_{\infty,1} \|g\|_1, \quad (36)$$

where

$$\|f\|_{\infty,1} := \sup_x \sum_{i_1, \dots, i_p} \int dx_2 \dots dx_p |f_{i_1 \dots i_p}(x, x_2, \dots, x_p)|.$$

Without loss of generality, we assume that $A = M_\Lambda(f)$ for some $f \in \mathcal{B}^{(p)}$ satisfying the bound (24). Iterating

$$\{H_\Lambda, M_\Lambda(f)\} = \sum_{n=1}^{\infty} np M_\Lambda(V^{(n)} \rightharpoonup f)$$

we obtain that

$$\begin{aligned} \{H_\Lambda, M_\Lambda(f)\}^{(l)} &= \sum_{n_1, \dots, n_l=1}^{\infty} [pn_1] [(p+n_1-1)n_2] \cdots [(p+n_1+\dots+n_{l-1}-l+1)n_l] \times \\ &\quad \times M_\Lambda \left(V^{(n_l)} \rightharpoonup \left(V^{(n_{l-1})} \rightharpoonup \dots (V^{(n_1)} \rightharpoonup f) \right) \right), \end{aligned}$$

with norm

$$\begin{aligned}
 & \left\| \{H_\Lambda, M_\Lambda(f)\}^{(l)} \right\|_\infty \\
 & \leq \sum_{n_1, \dots, n_l} [pn_1] [(p+n_1-1)n_2] \cdots [(p+n_1+\cdots+n_{l-1}-l+1)n_l] \times \\
 & \quad \times \|V^{(n_l)}\|_{\infty,1} \cdots \|V^{(n_1)}\|_{\infty,1} \|f\|_1 \\
 & \leq l! \sum_{n_1, \dots, n_l} \frac{(p+n_1+\cdots+n_l)^l}{l!} n_1 \cdots n_l \|V^{(n_l)}\|_{\infty,1} \cdots \|V^{(n_1)}\|_{\infty,1} \|f\|_1 \\
 & \leq e^p \|f\|_1 l! \left[\sum_n n e^n \|V^{(n)}\|_{\infty,1} \right]^l \\
 & \leq e^p \|f\|_1 l! \|V\|^l,
 \end{aligned} \tag{37}$$

by (30). Therefore, for $|t| < \|V\|^{-1}$, the series

$$\sum_{l=0}^{\infty} \frac{t^l}{l!} \{H_\Lambda, A\}^{(l)} \tag{38}$$

converges in $\|\cdot\|_\infty$ to $\alpha_\Lambda^t A$.

The quantum case is dealt with in a similar fashion, with the additional complication caused by the ordering of the generators $\{\widehat{S}_i(x)\}$. This does not trouble us, however, as an exact knowledge of the ordering is not required. It is easy to see that, for f and g as above,

$$ish^{-d} [\widehat{S}_\Lambda(f), \widehat{S}_\Lambda(g)]$$

is equal, up to a reordering of the spin operators, to $pq\widehat{S}_\Lambda(f \rightarrow g)$. Iterating this shows that

$$(ish^{-d})^l [\widehat{H}_\Lambda, \widehat{A}]^{(l)}$$

is equal, up to a reordering of the spin operators, to

$$\begin{aligned}
 & \sum_{n_1, \dots, n_l=1}^{\infty} [pn_1] [(p+n_1-1)n_2] \cdots [(p+n_1+\cdots+n_{l-1}-l+1)n_l] \times \\
 & \quad \times \widehat{S}_\Lambda \left(V^{(n_l)} \rightarrow \left(V^{(n_{l-1})} \rightarrow \cdots (V^{(n_1)} \rightarrow f) \right) \right),
 \end{aligned}$$

Consequently an estimate analogous to (37) yields, for $s \geq 1$,

$$\left\| (ish^{-d})^l [\widehat{H}_\Lambda, \widehat{A}]^{(l)} \right\| \leq e^p \|f\|_1^{(h)} l! (\|V\|^{(h)})^l,$$

which readily implies the bound

$$\left\| \sum_{l=0}^{\infty} \frac{t^l}{l!} (ish^{-d})^l [\widehat{H}_\Lambda, \widehat{A}]^{(l)} \right\| \leq e^p \|f\|_1^{(h)} \sum_{l=0}^{\infty} (|t| \|V\|^{(h)})^l. \tag{39}$$

If $s=1/2$, the first line of (37) gets the additional factor $\sqrt{2}^{n_1+\dots+n_l+p}$. This may be dealt with by replacing the factor $(p+n_1+\dots+n_l)^l$ in the second line of (37) with $(rp+rn_1+\dots+rn_l)^l/r^l$. The desired bound then follows for $0 < r \leq 1 - \frac{1}{2} \log 2$. Note that in this case the convergence radius for t is reduced to $r\|V\|^{-1}$. For ease of notation, we restrict the following analysis to the case $s \geq 1$, while bearing in mind that the extension to $s=1/2$ follows by using the above rescaling trick.

Now, by definition of $\|V\|$, for any $|t| < \|V\|^{-1}$ there is an h_0 such that (39) converges in norm to $\widehat{\alpha}_\Lambda^t \widehat{A}$ for all $h \leq h_0$, uniformly in h and Λ .

In order to establish the statement of the theorem for short times $|t| < \|V\|^{-1}$, we remark that the commutation relations (22) imply the bound

$$\|\mathbf{A} - \mathbf{B}\| \leq \frac{h^d}{s} p^2 \|f\|_1^{(h)},$$

for arbitrary reorderings, $\mathbf{A}$ and $\mathbf{B}$, of the same operator $\widehat{S}_\Lambda(f)$, with $f \in \mathcal{B}^{(p)}$ for some $p < \infty$.

If we define $\widehat{\alpha}_\Lambda^t A$ through its norm-convergent power series, we therefore get

$$\begin{aligned} & \left\| \widehat{\alpha}_\Lambda^t \widehat{A} - \widehat{\alpha}_\Lambda^t A \right\| \\ & \leq \frac{h^d}{s} \sum_{l=0}^{\infty} \frac{|t|^l}{l!} \sum_{n_1, \dots, n_l} [pn_1] [(p+n_1-1)n_2] \cdots [(p+n_1+\dots+n_{l-1}-l+1)n_l] \times \\ & \quad \times (n_1 + \dots + n_l - l + 1)^2 \|V^{(n_l)}\|_{\infty,1}^{(h)} \cdots \|V^{(n_1)}\|_{\infty,1}^{(h)} \|f\|_1^{(h)} \\ & \leq \frac{h^d}{s} \sum_{l=0}^{\infty} |t|^l \sum_{n_1, \dots, n_l} \frac{(p+n_1+\dots+n_l)^{l+2}}{l!} n_1 \cdots n_l \|V^{(n_l)}\|_{\infty,1}^{(h)} \cdots \|V^{(n_1)}\|_{\infty,1}^{(h)} \|f\|_1^{(h)} \\ & \leq \frac{h^d}{s} \sum_{l=0}^{\infty} |t|^l e^p \|f\|_1^{(h)} (l+2)(l+1) \left[\sum_n n e^n \|V^{(n_1)}\|_{\infty,1}^{(h)} \right]^l \\ & \leq \frac{h^d}{s} e^p \|f\|_1^{(h)} \sum_{l=0}^{\infty} (l+2)(l+1) (|t| \|V\|^{(h)})^l \\ & = O(h^d), \end{aligned}$$

where in the last step we have used the fact that the sum converges uniformly in h , for h small enough, as seen above.

Arbitrary times are reached by iteration of the above result. $\square$

3.5. THE THERMODYNAMIC LIMIT

The above result may again be formulated in the thermodynamic limit as $\Lambda \rightarrow h\mathbb{Z}^d$. We only sketch the arguments, which are almost identical to those of Section 2.5.

The quantum quasi-local algebra is

$$\widehat{\mathfrak{A}}^{(h)} := \overline{\bigvee_{\Lambda \subset \subset \mathbb{R}^d} \widehat{\mathfrak{A}}_{\Lambda}^{(h)}},$$

The existence of dynamics is guaranteed by the following statement.

LEMMA 6. *Let $h > 0$ and suppose $\mathbf{A} \in \widehat{\mathfrak{A}}_{\Lambda_0}^{(h)}$ for some bounded and open $\Lambda_0 \subset \mathbb{R}^d$. Then, for any $t \in \mathbb{R}$, the following limit exists in the norm sense:*

$$\lim_{\Lambda \rightarrow \infty} \widehat{\alpha}_{\Lambda}^t \mathbf{A} =: \widehat{\alpha}^t \mathbf{A},$$

By continuity this extends to a strongly continuous one-parameter group $(\widehat{\alpha}^t)_{t \in \mathbb{R}}$ of automorphisms of $\widehat{\mathfrak{A}}^{(h)}$.

The classical quasi-local algebra is

$$\mathfrak{A} := \overline{\bigvee_{\Lambda \subset \subset \mathbb{R}^d} \mathfrak{P}_{\Lambda}}.$$

LEMMA 7. *Let $A \in \mathfrak{P}_{\Lambda_0}$ for some open and bounded $\Lambda_0 \subset \subset \mathbb{R}^d$. Then, for any $t \in \mathbb{R}$, the following limit exists in $\|\cdot\|_{\infty}$:*

$$\lim_{\Lambda \rightarrow \infty} \alpha_{\Lambda}^t A =: \alpha^t A,$$

By continuity this extends to a strongly continuous one-parameter group $(\alpha^t)_{t \in \mathbb{R}}$ of automorphisms of $\mathfrak{A}$. Furthermore, $\alpha^t A = A \circ \phi^t$, where $\phi^t = \phi_{\mathbb{R}^d}^t$ is the Landau-Lifschitz flow defined in Lemma 5.

Now, for $f \in \mathcal{B}^{(p)}$, $M(f)$ and $\widehat{S}(f)$ are well-defined in the obvious way. Define $\mathfrak{P}$ as the algebra generated by functions of the form $M(f)$, where f satisfies (24).

THEOREM 5. *Let $A \in \mathfrak{P}$ and $\varepsilon > 0$. Then there exists a function $A(t) \in \mathfrak{P}$ such that*

$$\sup_{t \in \mathbb{R}} \|\alpha^t A - A(t)\|_{\infty} \leq \varepsilon, \quad (40)$$

and, for any $t \in \mathbb{R}$,

$$\|\widehat{\alpha}^t \widehat{A} - \widehat{A}(t)\| \leq \varepsilon + C(\varepsilon, t, A) h^d. \quad (41)$$

3.6. EVOLUTION OF COHERENT STATES

In this section, our “smearing functions” f are assumed to have compact support, i. e. to belong to the space

$$\mathcal{B}_c^{(p)} := \mathcal{B}^{(p)} \cap C_c(\mathbb{R}^{pd}; \mathbb{C}^{3p}).$$

In addition, we require the interaction potential V to be of finite range in the sense that there exists a sequence $R_n > 0$ such that if $|x_i - x_j| > R_n$ for some pair (i, j) then $V_{i_1 \dots i_n}^{(n)}(x_1, \dots, x_n) = 0$.

Next, we take some initial classical spin configuration $M \in C(\mathbb{R}^d; \mathbb{S}^2)$, or, more generally, a function $M: \mathbb{R}^d \rightarrow \mathbb{S}^2$ whose points of discontinuity form a null set. We shall study the time evolution of product states ρ_M on $\widehat{\mathfrak{A}}^{(h)}$ that reproduce the given classical state M . For open and bounded $\Lambda \subset \mathbb{R}^d$, we define the product state

$$|M_\Lambda\rangle := \bigotimes_{x \in \Lambda^{(h)}} |M(x)\rangle,$$

where $|M(x)\rangle$ is the coherent spin state corresponding to the unit vector $M(x)$. For $\mathbf{A} \in \widehat{\mathfrak{A}}_\Lambda^{(h)}$, define

$$\rho_M(\mathbf{A}) := \langle M_\Lambda, \mathbf{A} M_\Lambda \rangle,$$

which we extend to arbitrary $\mathbf{A} \in \widehat{\mathfrak{A}}^{(h)}$ by continuity.

For our main result on the time evolution of coherent states, we first record the following auxiliary result whose elementary proof we omit.

LEMMA 8. *Let $f \in \mathcal{B}_c^{(p)}$ satisfy (24). Then*

$$\lim_{h \rightarrow 0} \rho_M(\widehat{S}(f)) = M(f). \quad (42)$$

The last result in this paper links the quantum time evolution of coherent spin states with the classical evolution in the mean-field/continuum limit when the lattice spacing h tends to 0.

THEOREM 6. *Let $t \in \mathbb{R}$, $A \in \mathfrak{P}$ and M be as described above. Let $M(t)$ be the solution of (31) on $\mathbb{R}^d$ with initial configuration M . Then*

$$\lim_{h \rightarrow 0} \rho_M(\widehat{\alpha}^t \widehat{A}) = A(M(t)),$$

uniformly in t on compact time intervals.

Proof. The proof is a corollary of the proof of Theorem 4. First, let $|t| < \|V\|^{-1}$ and pick an $\varepsilon > 0$. Choose a cutoff such that the tails of the thermodynamic limits of the series (38) and (39) are bounded by ε . We therefore have to estimate a finite sum of terms of the form

$$|\rho_M(\widehat{S}(g)) - M(g)|,$$

where $g \in \mathcal{B}_c^{(p(g))}$ because of our assumptions on V . By Lemma 8, for h small enough, these are all bounded by ε , and the claim for small times follows. Finally, by iteration, we extend the result to arbitrary times. $\square$

Acknowledgements

We thank Simon Schwarz for very useful discussions. We also thank a referee for calling our attention to some relevant references.

References

1. Fröhlich, J., Knowles, A., Pizzo, A.: Atomism and quantization. *J. Phys. A* **40**(12), 3033–3045 (2007)
2. Gustafson, S., Kang, K., Tsai, T.-P.: Schrödinger flow near harmonic maps. *Comm. Pure Appl. Math.* **60**(4), 463–499 (2007)
3. Kružík, M., Prohl, A.: Recent developments in the modeling, analysis, and numerics of ferromagnetism. *SIAM Rev.* **48**(3), 439–483 (2006)
4. Lieb, E.H.: The classical limit of quantum spin systems. *Comm. Math. Phys.* **31**, 327–340 (1973)
5. Moser, M., Prets, A., Spitzer, W.L.: Time evolution of spin waves. *Phys. Rev. Lett.* **83**(17), 3542–3545 (1999)
6. Simon, B.: The classical limit of quantum partition functions. *Comm. Math. Phys.* **71**, 247–276 (1980)
7. Sulem, P.-L., Sulem, C., Bardos, C.: On the continuous limit for a system of classical spins. *Comm. Math. Phys.* **107**(3), 431–454 (1986)
8. Werner, R.F.: The classical limit of quantum theory. [quant-ph/9504016v1](#)
9. Werner, R.F., Wolff, M.P.: Classical mechanics as quantum mechanics with infinitesimal $\hbar$. *Phys. Lett. A* **202**, 155–159 (1995)