Soft triple-real radiation for Higgs production at N³LO

Journal Article**Author(s):**

Anastasiou, Charalampos; Duhr, Claude; Dulat, Falko; Mistlberger, Bernhard

Publication date:

2013-07

Permanent link:

<https://doi.org/10.3929/ethz-b-000074393>

Rights / license:

[Creative Commons Attribution 4.0 International](#)

Originally published in:

Journal of High Energy Physics 2013(7), [https://doi.org/10.1007/JHEP07\(2013\)003](https://doi.org/10.1007/JHEP07(2013)003)

Soft triple-real radiation for Higgs production at N₃LO

Charalampos Anastasiou,^a Claude Duhr,^{a,b} Falko Dulat^a and Bernhard Mistlberger^a

^a*Institute for Theoretical Physics, ETH Zurich, 8093 Zurich, Switzerland*

^b*Institute for Particle Physics Phenomenology, University of Durham,
Durham, DH1 3LE, U.K.*

E-mail: babis@phys.ethz.ch, duhrc@itp.phys.ethz.ch,
dulatf@itp.phys.ethz.ch, bmistlbe@itp.phys.ethz.ch

ABSTRACT: We present the two first terms in the threshold expansion of Higgs production partonic cross-sections at hadron colliders for processes with three partons in the final state. These are contributions to the inclusive Higgs cross-section in gluon fusion at N₃LO. We have developed a new technique for the expansion of the squared matrix-elements around the soft limit and for the reduction of the required phase-space integrals to only ten single-scale master integrals. We compute the master integrals building upon modern techniques for the integration of multidimensional integrals in dimensional regularization. Our results constitute an important step towards a systematic computation of the Higgs boson cross-section as an expansion around the threshold limit.

KEYWORDS: Higgs Physics, QCD

ARXIV EPRINT: [1302.4379](https://arxiv.org/abs/1302.4379)

Contents

1	Introduction	1
2	Reverse unitarity, threshold expansion and soft integrals	4
3	Validation of the method and examples	6
4	Relations between phase space integrals in different dimensions	10
5	From soft integrals to angular integrals and Mellin-Barnes integrals	12
6	Triple real emission phase space integrals in the soft limit	14
6.1	Triple real soft master integrals for Higgs production	14
6.2	Dimensional recurrence relations	16
6.3	Analytic results for the soft master integrals	17
7	The threshold approximation to the partonic $2 \rightarrow H + 3$ parton cross-section	20
7.1	General setup	20
7.2	Calculation	21
7.3	Leading soft contribution	21
7.4	Next-to-leading soft contribution	24
8	Analytic computation of the master integrals	30
8.1	The master integral $\mathcal{F}_3$	31
8.2	The master integral $\mathcal{F}_2$	32
8.3	The master integral $\mathcal{F}_7$	32
8.4	The master integral $\mathcal{F}_4$	33
8.5	The master integral $\mathcal{F}_6$	34
8.6	The master integral $\mathcal{F}_{10}$	35
8.7	The master integral $\mathcal{F}_5$	35
8.8	The master integral $\mathcal{F}_8$	36
8.9	The master integral $\mathcal{F}_9$	43
9	Conclusions and outlook	47
A	Parametrisation of the $2 \rightarrow H + (N - 2)$ phase space	49
B	Derivation of the phase space volume	51
C	From Mellin-Barnes integrals to parametric integrals	53

D Symbolic integration of parametric integrals	55
D.1 Denominator reduction	56
D.2 Symbolic integration	58
D.3 Symbolic integration and denominator reduction	63
D.4 A toy example	65

1 Introduction

The first years of experiments at the Large Hadron Collider (LHC) have probed decisively the physics of electroweak symmetry breaking, demonstrating a great understanding of the theory at this energy scale. The success of particle theory must be attributed not only to the Standard Model (SM), but also to remarkable progress in perturbative calculations. Results in this field of research are outstanding and have resulted in very accurate comparisons with data. Indeed, partonic cross-sections for a large variety of processes are now routinely computed at next-to-leading order (NLO) in the strong coupling constant α_s .¹ In addition, modern Monte-Carlo programs have reached a high level of sophistication with the development of methods to match and merge parton showers with fixed-order calculations.² Finally, basic processes at the LHC with a small final-state multiplicity can now be computed fully differentially at next-to-next-to-leading order (NNLO).³

A classic example where perturbative calculations have been crucial for the interpretation of LHC data is the production of a Higgs boson. Perturbative corrections for Higgs boson signal processes are generically large, especially so for production via gluon fusion [53–59]. For this process, the total and fully differential cross-sections are known through NNLO with a typical uncertainty of about 10% due to variations of the factorization and renormalization scales [50, 60]. Indeed, without the knowledge of the NNLO corrections, predictions at LO or NLO would be assigned theory uncertainties which are larger than the already achieved experimental uncertainties. With future LHC data, a comparison with theory at a level of precision of a few percent will be revealing the fine details of the mechanism of electroweak symmetry breaking and potentially uncovering new fundamental laws of physics.

Besides being of importance to Higgs physics, NNLO predictions have promoted the Drell-Yan process to a standard candle for studies at hadron colliders, where observables in this process are computed with a typical precision of better than 2% as indicated by scale variations [61, 62]. Measurements of the mass of the W boson and the weak mixing angle from Drell-Yan data are one of the most stringent constraints on the Standard Model. Furthermore, Drell-Yan production data is an essential input for the extraction of parton distribution functions from hadron collider processes. Last but not least, the clean detector signatures of Drell-Yan events, combined with the excellent theoretical predictions of their

¹See, for example, refs. [1–18]

²See, for example, refs. [19–39]

³See, for example, refs. [40–52]

rates, allow for a precise determination of the luminosity. Indeed, luminosity determination from Drell-Yan is relatively insensitive to high pile-up conditions, a fact which will be even more important during the high-luminosity phase of the LHC.

We believe that it is possible to develop efficient methods for computing Higgs boson and Drell-Yan hadroproduction cross-sections at the next perturbative order, “N₃LO”. The benefits of such a program are potentially very important.

In Higgs production via gluon fusion the uncertainty due to scale variations at N₃LO is anticipated to be half of what is found at NNLO [63]. Scale variation estimates cannot be derived from first principles, but entail a certain degree of subjectiveness. With the knowledge of more orders in the perturbative expansion, the remaining perturbative uncertainty can be estimated more reliably, not only from scale variations but also from the progression of the series. Improvements of the Higgs boson production cross-section uncertainty will propagate into the determination of the couplings of the Higgs boson.

Although existing NNLO calculations for the Drell-Yan process indicate that the cross-section is already known very precisely, an explicit N₃LO computation will educate us further by challenging or verifying the traditional prescriptions for uncertainty estimates. Furthermore, high-luminosity runs at the LHC require the triggering of leptons at a higher transverse momentum, resulting in a deterioration of the NNLO scale uncertainty [49]. An excellent theoretical precision can then be recovered by including the N₃LO perturbative corrections.

So far, there has been no N₃LO calculation for hadron collider processes performed in the literature. In this paper, we attempt a first step towards computing inclusive hadroproduction cross-sections at this perturbative order. We focus on one of the most complicated N₃LO contributions to the inclusive Higgs boson cross-section in the gluon fusion channel, namely, the contributions coming from partonic cross-sections for tree-level radiative processes with three additional partons emitted in the final state. A direct integration of the corresponding matrix-elements over phase space is however tedious. To simplify the problem, we perform a threshold expansion and devise a method to compute the coefficients of the expansion analytically. One of the main results of our paper is the computation of the first two terms of the threshold expansion for triple real corrections to inclusive Higgs production.

Our expansion method builds upon *reverse unitarity* [59, 62, 64–66], a technique developed for the calculation of the inclusive Higgs cross-section and the rapidity distribution of electroweak gauge bosons at NNLO. The main idea is that phase-space integrals of matrix-elements are dual to loop integrals in their algebraic properties: recurrence identities in the powers of propagators and the number of dimensions as well as differential equations satisfied by loop integrals also apply to phase-space integrals. This is achieved by associating on-shell conditions or other phase-space constraints in the form of delta functions with differences of otherwise identical propagators with opposite infinitesimal imaginary parts, in the spirit of Cutkosky’s rules. In this paper, we observe and exploit that the duality rules of reverse unitarity can be expanded in kinematic parameters.

After performing a threshold expansion of loop integrals dual to the cross-sections for Higgs plus three partons processes, we apply automated reduction algorithms to reduce

the coefficients of the expansion to master integrals. We have performed the reduction using existing public programs and programs developed specifically for the purpose of this computation.

The reduction yields ten master integrals for the first two coefficients in the threshold expansion. The master integrals themselves are not specific to Higgs production and will appear in the threshold expansion of other hadroproduction processes, such as Drell-Yan. For processes with a single colorless final state our set of master integrals is complete and our expressions are universal for the partonic cross-sections with the same initial state at leading order in the threshold expansion.

For the computation of the *soft* master integrals we have employed a combination of methods. The master integrals are evaluated by using a parametrization of the phase space in terms of the energies and angles of the momenta of the soft partons. When needed, we perform the phase-space integrals by introducing Mellin-Barnes representations. We evaluate some of the Mellin-Barnes master integrals in a closed form as hypergeometric functions or directly as a Laurent-expansion in the dimensional regulator ϵ with infinite nested sums as coefficients, which naturally evaluate to multiple zeta values in all cases. Some master integrals cannot be evaluated easily from their Mellin-Barnes representations. We have developed a procedure to turn Mellin-Barnes integrals into integrals over positive real parameters, which are easy to expand in ϵ whenever the integral is finite. The resulting parametric integrals are then evaluated by integrating out the integration variables one-by-one in terms of multiple polylogarithms. The last integration then again produces multiple zeta values in a natural way.

We have restricted our computation of the partonic cross-sections to the first two terms of the threshold expansion. There are two obvious ways to extend this work in the future. We could continue with the computation of subleading terms in the threshold expansion. This requires a more intensive but not prohibitive computer algebra and the evaluation of *soft* master integrals from topologies which contribute only to higher orders in the soft expansion. The soft expansion is rather fast converging for Higgs production but rather slow for Drell-Yan, as has been noticed at NNLO. A second possibility is to perform the phase-space integrations for arbitrary kinematics, reducing them to a different set of master integrals. The *soft* master integrals which we present in this article can serve as boundary conditions for solving the differential equations satisfied by the master integrals in arbitrary kinematics.

This article is organized as follows: in section 2 we present our general strategy based on reverse-unitarity to perform the threshold expansion for real-emission cross sections in terms of soft integrals, and we illustrate the method on some simple examples in section 3. In sections 4 and 5 we study some properties of soft integrals in general, and we show that there is an easy and canonical way to derive dimensional recurrence relations and Mellin-Barnes integral representations for generic soft integrals. The soft master integrals contributing to the first two terms in the threshold expansion of the leading-order partonic cross sections for $pp \rightarrow H + 3$ partons are discussed in section 6, and the corresponding results for the cross sections are presented in section 7. Section 8 contains technical details about the computation of the soft master integrals, and in section 9 we draw our conclusions

and give an outlook for future work. The paper contains several appendices discussing phase parametrisations and a generic formula for the phase-space volume for $H + n$ partons, as well as a new method to derive a parametric integral representation from a Mellin-Barnes integral and a description of an algorithmic way to perform the analytic integration of certain classes of parametric integrals.

2 Reverse unitarity, threshold expansion and soft integrals

We consider the production of a Higgs boson in association with $j = 3 \dots N$ massless partons in the final state from two massless partons $i = 1, 2$ in the initial state,

$$1 + 2 \rightarrow H + 3 + \dots + N. \quad (2.1)$$

The inclusive cross-section for this process in dimensional regularization is given by a phase-space integral over the momenta q_j of the final-state partons,

$$\sigma = \int d\Phi_{N-1}(q_H, q_3, \dots, q_N; M^2; s; D) |\mathcal{A}|^2(\{q_j\}, q_1, q_2; D). \quad (2.2)$$

We work in $D = 4 - 2\epsilon$ dimensions and denote by q_1, q_2 the momenta of the initial-state partons and we also use the shorthand notation

$$q_{12} = q_1 + q_2, \quad q_{345} = q_3 + q_4 + q_5, \quad \text{etc.}$$

In the following we will often drop the functional dependence on the dimension D for clarity. The mass of the Higgs boson is denoted by M , and we denote the (squared) center-of-mass energy by $s = (q_1 + q_2)^2$. $|\mathcal{A}|^2$ represents the squared matrix-element multiplied with the appropriate flux and symmetry factors. The D -dimensional phase space measure is given by

$$\begin{aligned} & d\Phi_{N-1}(q_H, q_3, \dots, q_N; M^2; s; D) \\ &= (2\pi)^D \delta^{(D)}(q_{12} - q_H - q_{3\dots N}) \frac{d^D q_H}{(2\pi)^{D-1}} \delta_+(q_H^2 - M^2) \prod_{j=3}^N \frac{d^D q_j}{(2\pi)^{D-1}} \delta_+(q_j^2), \end{aligned} \quad (2.3)$$

with $\delta_+(q^2 - m^2) = \delta(q^2 - m^2)\Theta(q^0)$. Integrating out the momentum of the Higgs boson, we can rewrite eq. (2.2) as

$$\sigma = \frac{1}{2\pi} \int \left[\prod_{j=3}^N \frac{d^D q_j}{(2\pi)^{D-1}} \delta_+(q_j^2) \right] \delta_+([q_{3\dots N} - q_{12}]^2 - M^2) |\mathcal{A}|^2(\{q_j\}, q_1, q_2). \quad (2.4)$$

In this article, we restrict ourselves to the case of real-radiation matrix-elements without virtual corrections. We introduce the variables

$$z = \frac{M^2}{s} \quad \text{and} \quad \bar{z} = 1 - z. \quad (2.5)$$

We now rescale the momenta of all the partons,

$$q_i = \begin{cases} \sqrt{s} p_i, & \text{if } i = 1, 2, \\ \sqrt{s} \bar{z} p_i, & \text{if } i = 3 \dots N, \end{cases} \quad (2.6)$$

which captures the scaling of the partonic momenta in the final-state. We emphasize that this is not, as yet, an approximation, but rather a convenient change of integration variables which captures the correct asymptotic behavior at threshold as $\bar{z} \rightarrow 0$. In the following we assume $s = 1$, and we find

$$\begin{aligned} \sigma = \bar{z}^{(D-2)(N-2)-1} \frac{1}{2\pi} \int \left[\prod_{j=3}^N \frac{d^D p_j}{(2\pi)^{D-1}} \delta_+(p_j^2) \right] \delta_+ \left([p_{3\dots N} - p_{12}]^2 - z p_{3\dots N}^2 \right) \\ \times |\mathcal{A}|^2 (\{\bar{z} p_j\}, p_1, p_2). \end{aligned} \quad (2.7)$$

Note that the full s -dependence can easily be recovered from dimensional analysis.

The squared matrix-element $|\mathcal{A}|^2$ consists of a rapidly growing number of terms with N , yielding a correspondingly large number of phase-space integrals. The method of reverse unitarity, developed in refs. [59, 62, 64–66], allows the reduction of phase-space integrals to a basis of fewer master integrals by establishing a duality of phase-space and loop integrals, where the latter are amenable to algebraic methods [67, 68] based on integration by parts [69, 70].

Following the reverse unitarity approach, on-shell and other phase-space constraints are dual to propagators:

$$\delta_+(q^2) \rightarrow \left(\frac{1}{q^2} \right)_c \equiv \frac{1}{2\pi i} \text{Disc} \frac{1}{q^2} = \frac{1}{2\pi i} \left(\frac{1}{q^2 + i0} - \frac{1}{q^2 - i0} \right). \quad (2.8)$$

“Cut” propagators can be differentiated in a similar way to ordinary propagators with respect to their momenta,

$$\frac{\partial}{\partial q_\mu} \left[\left(\frac{1}{q^2} \right)_c \right]^\nu = -\nu \left[\left(\frac{1}{q^2} \right)_c \right]^{\nu+1} 2q^\mu, \quad (2.9)$$

leading to identical integration-by-parts (IBP) identities for phase-space integrals as for their dual loop integrals. Solving the system of IBP identities for phase-space integrals proceeds in the same way as for loop integrals, with the exception that for cut-propagators we can use the simplifying identity:

$$\left[\left(\frac{1}{q^2} \right)_c \right]^{-\nu} \rightarrow 0, \quad \forall \nu = 0, 1, 2, \dots \quad (2.10)$$

According to the reverse unitarity method, we find a dual forward scattering loop-amplitude with $N - 1$ cut-propagators for the real radiation contribution of eq. (2.7), namely,

$$\begin{aligned} \sigma = \bar{z}^{(D-2)(N-2)-1} \frac{1}{2\pi} \int \left[\prod_{j=3}^N \frac{d^D p_j}{(2\pi)^{D-1}} \left(\frac{1}{p_j^2} \right)_c \right] \left[\frac{1}{[p_{3\dots N} - p_{12}]^2 - z p_{3\dots N}^2} \right]_c \\ \times |\mathcal{A}|^2 (\{\bar{z} p_j\}, p_1, p_2). \end{aligned} \quad (2.11)$$

In this article, we take one further step and expand cut-propagators and the squared matrix-elements around $z = 1$,

$$|\mathcal{A}|^2(\{\bar{z} p_j\}, p_1, p_2) = \bar{z}^{-2(N-2)} \sum_{k=0}^{\infty} |\mathcal{A}|_k^2(\{p_j\}, p_1, p_2) \bar{z}^k, \quad (2.12)$$

and

$$\frac{1}{[p_{3\dots N} - p_{12}]^2 - z p_{3\dots N}^2} = \sum_{k=0}^{\infty} \bar{z}^k \frac{(-p_{3\dots N}^2)^k}{[p_{12} \cdot (p_{12} - 2p_{3\dots N})]^{k+1}}. \quad (2.13)$$

In this approximation, the cross section can be expanded into a power series in $\bar{z}$,

$$\sigma = \bar{z}^{(D-4)(N-2)-1} \sum_{k=0}^{\infty} \bar{z}^k \sigma^{S(k)}. \quad (2.14)$$

The coefficients of the power series are given by

$$\sigma^{S(k)} = \sum_{l=0}^k (-1)^l \int d\Phi_{N-1}^S \left[\frac{1}{p_{12} \cdot (p_{12} - 2p_{3\dots N})} \right]_c^l (p_{3\dots N}^2)^l |\mathcal{A}|_{k-l}^2(\{p_j\}, p_1, p_2), \quad (2.15)$$

where $d\Phi_{N-1}^S$ denotes the “soft” phase space measure

$$\begin{aligned} d\Phi_{N-1}^S &\equiv \frac{1}{2\pi} \left[\frac{1}{p_{12} \cdot (p_{12} - 2p_{3\dots N})} \right]_c \prod_{j=3}^N \frac{d^D p_j}{(2\pi)^{D-1}} \left(\frac{1}{p_j^2} \right)_c \\ &= \frac{1}{2\pi} \delta_+(p_{12}^2 - 2p_{12} \cdot p_{3\dots N}) \prod_{j=3}^N \frac{d^D p_j}{(2\pi)^{D-1}} \delta_+(p_j^2). \end{aligned} \quad (2.16)$$

The integrals which emerge after the $\bar{z}$ expansion depend trivially on one dimensionful parameter $p_{12}^2 = s$. If we put $s = 1$, the integrals are number integrals whose only functional dependence is through the space-time dimension $D = 4 - 2\epsilon$. We will refer to such integrals as *soft (phase space) integrals*, and they are the main subject of this paper. We note that, apart from the cut Higgs boson propagator, the integrands of soft phase space integrals are homogeneous functions under a simultaneous rescaling of the final-state momenta. In addition, a soft integral can be reduced to a set of “soft” master integrals using IBP identities by exploiting the duality to loop integrals via reverse-unitarity. We will illustrate this property in the next section where we check our method on several examples.

3 Validation of the method and examples

In this section, we study the validity of the method described in the previous section at NLO and NNLO — two perturbative orders that are well studied in the literature and so we can compare our results readily with known results. In particular, we show that our method reproduces the correct results for the leading behavior of NLO and NNLO real emission amplitudes in the soft limit, as well as for the subleading terms in the expansion of the phase space volume up to N₃LO and for a non-trivial double real emission master integral at NNLO.

At NLO, all phase space integrals that contribute to the real emission amplitude in general kinematics can be reduced to the phase space volume for $H + 1$ parton,

$$\Phi_2(\bar{z}; \epsilon) = \frac{1}{2(4\pi)^{1-\epsilon}} \frac{\Gamma(1-\epsilon)}{\Gamma(2-2\epsilon)} \bar{z}^{1-2\epsilon}. \quad (3.1)$$

As there is only one master integral which is a monomial in $\bar{z}$, our method trivially gives the correct answer at NLO.

At NNLO all double real emission phase space integral can be reduced in general kinematics to a linear combination of 18 master integrals [59]. The leading contribution of all master integrals in the soft limit to all orders in ϵ was computed in ref. [71], and it was observed that in this limit 17 master integrals are proportional to the soft limit of the phase space volume for $H + 2$ partons,

$$\begin{aligned} \Phi_3(\bar{z}; \epsilon) &= \frac{1}{2(4\pi)^{3-2\epsilon}} \bar{z}^{3-4\epsilon} \frac{\Gamma(1-\epsilon)^2}{\Gamma(4-4\epsilon)} {}_2F_1(1-\epsilon, 2-2\epsilon; 4-4\epsilon; \bar{z}) \\ &= \bar{z}^{3-4\epsilon} \Phi_3^S(\epsilon) + \mathcal{O}(\bar{z}^4), \end{aligned} \quad (3.2)$$

where we defined

$$\Phi_3^S(\epsilon) = \frac{1}{2(4\pi)^{3-2\epsilon}} \frac{\Gamma(1-\epsilon)^2}{\Gamma(4-4\epsilon)}. \quad (3.3)$$

More precisely, it was shown in ref. [71] that if $\mathbf{X}_i^S(\bar{z}; \epsilon)$ denotes the leading term in the soft limit of the double real emission master integrals, then we can write⁴

$$\begin{aligned} \mathbf{X}_i^S(\bar{z}; \epsilon) &= \mathbf{S}_i(\bar{z}; \epsilon) \Phi_3^S(\epsilon), \quad 1 \leq i \leq 17, \\ \mathbf{X}_{18}^S(\bar{z}; \epsilon) &= -4 \bar{z}^{-1-4\epsilon} \frac{(1-2\epsilon)(3-4\epsilon)(1-4\epsilon)}{\epsilon^3} {}_3F_2(1, 1, -\epsilon; 1-\epsilon, 1-2\epsilon; 1) \Phi_3^S(\epsilon), \end{aligned} \quad (3.4)$$

where $\mathbf{S}_i(\bar{z}; \epsilon)$ are monomials in $\bar{z}$ and rational functions of ϵ . Using the method described in the previous section, we can easily explain the structure of eq. (3.4). Indeed, we observe that in the soft limit all the double real emission phase space integrals can be reduced to only two master integrals. In particular, the IBP identities in the soft limit allow us to express all but one of the $\mathbf{X}_i^S$ in terms of the phase space volume, and the coefficients appearing in the reduction are precisely the functions $\mathbf{S}_i$. In other words, in the soft limit all double real emission phase space integrals can be reduced to linear combinations of the following two soft master integrals

$$\begin{array}{c} \text{Diagram 1: A circle with two horizontal lines passing through its center. The left line has arrows pointing right towards the circle, labeled 1 and 2. The right line has arrows pointing left towards the circle, labeled 1 and 2. A vertical dashed line passes through the center of the circle.} \end{array} = \int d\Phi_3^S, \quad (3.5)$$

$$\begin{array}{c} \text{Diagram 2: A triangle with vertices labeled 1, 2, and 3. Solid lines connect the vertices. Dashed lines extend from each vertex. Arrows point towards the vertices: from the left towards vertex 1 (labeled 1, 2), from the bottom towards vertex 2 (labeled 1, 2), and from the right towards vertex 3 (labeled 1, 2).} \end{array} = \int \frac{d\Phi_3^S}{s_{14}s_{23}s_{34}}. \quad (3.6)$$

⁴Note that the normalization differs slightly from the normalization of ref. [71].

Our method thus provides the correct leading soft behavior of the double real emission contribution at NNLO. We emphasize that all the diagrams in this paper represent soft phase space integrals, i.e., all the diagrams represent integrals with respect to the soft phase space measure of eq. (2.16). In addition, the invariants appearing in the integrands of the soft integrals are defined with respect to the rescaled momenta defined in eq. (2.6),

$$s_{ij} = (\tau_i p_i + \tau_j p_j)^2, \quad \tau_i = \begin{cases} -1, & \text{if } i = 1, 2, \\ +1, & \text{if } i = 3 \dots N. \end{cases} \quad (3.7)$$

Our method does not only allow us to compute the leading soft behavior, but we can consistently expand around the soft limit $\bar{z} = 0$. In the following we show that we can correctly reproduce the first few terms in the soft expansion of double and triple emission phase space volumes, as well as for the NNLO master integral $\mathbf{X}_{18}$ of refs. [59, 71, 72].

Let us start with the phase space volume for $H + 2$ partons in the limit where the two partons are soft. On the one hand, from eq. (3.2) we immediately see that Φ_3 admits the expansion

$$\begin{aligned} \Phi_3(\bar{z}; \epsilon) &= \bar{z}^{3-4\epsilon} \Phi_3^S(\epsilon) \sum_{n=0}^{\infty} \frac{(1-\epsilon)_n (2-2\epsilon)_n}{(4-4\epsilon)_n} \bar{z}^n \\ &= \bar{z}^{3-4\epsilon} \Phi_3^S(\epsilon) \left[1 + \frac{1-\epsilon}{2} \bar{z} + \frac{(1-\epsilon)(2-\epsilon)(3-2\epsilon)}{4(5-4\epsilon)} \bar{z}^2 + \mathcal{O}(\bar{z}^3) \right]. \end{aligned} \quad (3.8)$$

On the other hand, using eq. (2.13) we obtain the diagrammatic expansion

$$\Phi_3(\bar{z}; \epsilon) = \bar{z}^{3-4\epsilon} \left[\text{Diagram 1} - \bar{z} \text{Diagram 2} + \bar{z}^2 \text{Diagram 3} + \mathcal{O}(\bar{z}^3) \right], \quad (3.9)$$

where the dashed lines indicate numerator factors and dots represent additional powers of the propagators or the numerators. The diagrams appearing in eq. (3.9) are in one-to-one correspondence with the terms in the expansion (3.8). Indeed, IBP reduction of the integrals in eq. (3.9) reveals

$$\text{Diagram 2} = -\frac{1-\epsilon}{2} \text{Diagram 1}, \quad (3.10)$$

$$\text{Diagram 3} = \frac{(1-\epsilon)(2-\epsilon)(3-2\epsilon)}{4(5-4\epsilon)} \text{Diagram 1}, \quad (3.11)$$

in perfect agreement with eq. (3.8). We checked explicitly that our method reproduces correctly the first ten terms of the soft expansion of the phase space volume for $H + 2$ partons.

As a second example we derive the subleading terms in the soft expansion of the double real emission master integral $\mathbf{X}_{18}$. Unlike for the phase space volume, no result is known for

$\mathbf{X}_{18}$ valid to all orders in ϵ in general kinematics, but the integral was evaluated explicitly up to $\mathcal{O}(\epsilon)$ in terms of harmonic polylogarithms [73] in ref. [59, 71, 72]. We can thus compare the result of our method order by order in ϵ to the expansion of the harmonic polylogarithms around $z = 1$. Using eq. (2.13) we obtain

$$\int \frac{d\Phi_3}{q_{14}^2 q_{23}^2 q_{34}^2} = \bar{z}^{-1-4\epsilon} \left[\text{triangle diagram} - \bar{z} \text{triangle diagram} + \bar{z}^2 \text{triangle diagram} + \mathcal{O}(\bar{z}^3) \right], \quad (3.12)$$

IBP reduction of the diagrams appearing in the subleading terms gives

$$\text{triangle diagram} = -\frac{2(1-4\epsilon)(3-4\epsilon)(1-2\epsilon)}{\epsilon^2} \text{circular diagram}, \quad (3.13)$$

$$\text{triangle diagram} = \frac{(3-4\epsilon)(1-2\epsilon)(2\epsilon^2-2\epsilon+1)}{\epsilon^2} \text{circular diagram}. \quad (3.14)$$

We checked that using these identities we can reproduce correctly the first five terms in the soft expansion of $\mathbf{X}_{18}$.

The aim of this paper is to compute the leading terms in the soft expansion of the triple real emission amplitude for inclusive Higgs production. In order to test our method at N₃LO, we verified that we can reproduce the correct soft expansion of the phase space volume for $H + 3$ partons. The phase space volume for $H + 3$ partons in general kinematics can be written in the form (see appendix B)

$$\begin{aligned} \Phi_4(\bar{z}; \epsilon) &= \frac{1}{2(4\pi)^{5-3\epsilon}} \bar{z}^{5-6\epsilon} \frac{\Gamma(1-\epsilon)^3}{\Gamma(6-6\epsilon)} {}_2F_1(2-2\epsilon, 3-3\epsilon; 6-6\epsilon; \bar{z}) \\ &= \bar{z}^{5-6\epsilon} \Phi_4^S(\epsilon) \left[1 + (1-\epsilon)\bar{z} + \frac{(1-\epsilon)(3-2\epsilon)(4-3\epsilon)}{2(7-6\epsilon)} \bar{z}^2 + \mathcal{O}(\bar{z}^3) \right], \end{aligned} \quad (3.15)$$

where we defined

$$\Phi_4^S(\epsilon) = \frac{1}{2(4\pi)^{5-3\epsilon}} \frac{\Gamma(1-\epsilon)^3}{\Gamma(6-6\epsilon)}. \quad (3.16)$$

Using our method, we obtain the following diagrammatic expansion

$$\Phi_4(\bar{z}; \epsilon) = \bar{z}^{5-6\epsilon} \left[\text{circular diagram} - \bar{z} \text{triangle diagram} + \bar{z}^2 \text{triangle diagram} + \mathcal{O}(\bar{z}^3) \right]. \quad (3.17)$$

All the diagrams in the expansion can be reduced to the soft phase space volume, as expected,

$$= -(1 - \epsilon) \text{ (diagram) }, \quad (3.18)$$

$$= \frac{(1 - \epsilon)(3 - 2\epsilon)(4 - 3\epsilon)}{2(7 - 6\epsilon)} \text{ (diagram) }. \quad (3.19)$$

To summarize, our method provides a systematic way to perform the threshold expansion of phase space integrals for the production of a heavy colorless state. Every term in the expansion corresponds to a soft integral, as defined in section 2, which can be reduced to a small set of soft master integral using IBP reduction. In the next two sections we study some additional properties of soft integrals in general, before applying our method to compute the threshold expansion of the triple real emission contribution to inclusive Higgs production.

4 Relations between phase space integrals in different dimensions

It is well known that loop integrals in different space-time dimensions are related by so-called dimensional shift identities [74]. After reduction to master integrals, the dimensional shift identities reduce to recurrence relations in the space-time dimension D for the master integrals themselves [75–79]. Reverse unitarity rules show that the same dimensional shift identities hold for the dual phase-space integrals (see also ref. [80]).

In this section we present an easy way to derive the dimensional shift identities for phase space integrals. We stress that the results of this section are generic and apply to phase space integrals in general kinematics. To start, let us consider a phase space integral in D dimensions,

$$F(D; \nu_1, \dots, \nu_n) = \int d\Phi_{N-1}(D) f(\nu_1, \dots, \nu_n), \quad (4.1)$$

where we explicitly indicate the dependence on the space-time dimension D . The integrand f can be written as a product,

$$f(\nu_1, \dots, \nu_N) = \prod_{l=1}^N P_l^{-\nu_l}, \quad (4.2)$$

where the P_l are polynomials in the rescaled kinematic invariants s_{ij} , raised to some power ν_l . In appendix A we show that the phase space measure $d\Phi_{N-1}(D)$ for parton + parton $\rightarrow H + (N - 2)$ partons can be parametrized solely in terms of kinematic invariants,

$$d\Phi_{N-1}(D) = \mathcal{N}_{N-2}(D) \bar{z}^{(N-2)(D-2)-1} \left(\prod_{\substack{1 \leq i, j \leq N \\ i \neq j, (i, j) \neq (1, 2)}} ds_{ij} \right) \times \delta \left(1 - \sum_{i=3}^N (s_{1i} + s_{2i}) + \bar{z} \sum_{i=3}^N \sum_{j=3}^{i-1} s_{ij} \right) G_N^{\frac{D-N-1}{2}}(\{s_{ij}\}) \Theta[G_N(\{s_{ij}\})], \quad (4.3)$$

where we have defined the normalisation factor

$$\mathcal{N}_{N-2}(D) = (-1)^{\frac{(N-2)(N-3)}{2}} 2^{-(N-2)\frac{D}{2}} (2\pi)^{(N-1)-(N-2)D} \prod_{i=3}^N \Omega_{D-i+1}, \quad (4.4)$$

$$\Omega_D = \frac{2\pi^{\frac{D}{2}}}{\Gamma(\frac{D}{2})},$$

and the Gram determinant G_N

$$G_N(\{s_{ij}\}) = \det(s_{1i}s_{2j} + s_{1j}s_{2i} - s_{ij})_{3 \leq i,j \leq n}. \quad (4.5)$$

Obviously, the D -dependent constants factor out of the integral, and so the actual integral depends on D only through the exponent of the Gram determinant. It is then easy to see that the phase space measures in shifted dimensions are related by

$$d\Phi_{N-1}(D+2) = \frac{\mathcal{N}_{N-2}(D+2)}{\mathcal{N}_{N-2}(D)} \bar{z}^{2(N-2)} d\Phi_{N-1}(D) G_N(\{s_{ij}\}). \quad (4.6)$$

We can thus express a phase space integral in $D+2$ dimensions as

$$F(D+2; \nu_1, \dots, \nu_n) = \frac{\mathcal{N}_{N-2}(D+2)}{\mathcal{N}_{N-2}(D)} \bar{z}^{2(N-2)} \int d\Phi_{N-1}(D) G_N(\{s_{ij}\}) f(\nu_1, \dots, \nu_n). \quad (4.7)$$

For a given set of polynomials $\{P_l\}$, the form of the integrand f depends only on the exponents $\{\nu_i\}$. One therefore introduces the operators I_i^+ and I_i^- acting on f as

$$\begin{aligned} (I_i^+ f)(\nu_1, \dots, \nu_i, \dots, \nu_n) &= f(\nu_1, \dots, \nu_i + 1, \dots, \nu_n), \\ (I_i^- f)(\nu_1, \dots, \nu_i, \dots, \nu_n) &= f(\nu_1, \dots, \nu_i - 1, \dots, \nu_n). \end{aligned} \quad (4.8)$$

If we assume that the $\{P_l\}$ are linearly independent, we can express the invariants s_{ij} as linear combinations of the I_i^+ and I_i^- . This allows us to rewrite the extra power of the Gram determinant in eq. (4.6) as a polynomial of degree N in the I_i^-

$$G_N(\{s_{ij}\}) f(\nu_1, \dots, \nu_n) = G_N(\{I_i^-\}) f(\nu_1, \dots, \nu_n). \quad (4.9)$$

We thus obtain the following compact formula relating phase space integrals in different dimensions,

$$F(D+2; \nu_1, \dots, \nu_n) = \frac{\mathcal{N}_{N-2}(D+2)}{\mathcal{N}_{N-2}(D)} \bar{z}^{2(N-2)} G_N(\{I_i^-\}) F(D; \nu_1, \dots, \nu_n). \quad (4.10)$$

Every term in the polynomial can be evaluated according to the action of the I_i^- operators, yielding a superposition of modified integrals in D dimensions. By applying this method to a master integral, we can express the master integral in $D+2$ dimensions as a linear combination of integrals in D dimensions. Using IBP identities, we can reduce the integrals in D dimensions to master integrals and thus we find a relation between the master integral in $D+2$ dimensions and D dimensions. This dimensional recurrence relation can formally be written as

$$F_i(D+2) = \sum_j c_{ij}(D) F_j(D), \quad (4.11)$$

with coefficients $c_{ij}(D)$ that are determined from the IBP reduction.

5 From soft integrals to angular integrals and Mellin-Barnes integrals

In this section we show that there is a canonical way to derive a Mellin-Barnes (MB) representation for the soft integrals that appear in the threshold expansion of phase space integrals. The soft integrals we need to consider have the form

$$F(\epsilon) = \int d\Phi_{N-1}^S f(\{p_j\}; p_1, p_2), \quad (5.1)$$

where $f(\{p_j\}; p_1, p_2)$ is a ratio of products of multi-particle invariants that is homogeneous under a simultaneous rescaling of the final-state momenta, i.e.,

$$f(\{\lambda p_j\}; p_1, p_2) = f(\{p_j\}; p_1, p_2) \lambda^a, \quad (5.2)$$

for some a . Note that this is true only if the (cut) Higgs boson propagator is not raised to an additional power. In that case, however, we can always reduce the integrand to a homogeneous function using IBP identities.

Next, we note that there is a subclass of soft integrals that have an additional property: they are homogeneous with respect to *individual* rescalings of the final-state momenta, i.e.,

$$f(\{\lambda_j p_j\}; p_1, p_2) = f(\{p_j\}; p_1, p_2) \prod_{j=3}^N \lambda_j^{a_j}, \quad (5.3)$$

for some a_j . This subclass of soft integrals is precisely the one where the integrand consists of products of power of *two*-particle invariants,

$$f(\{p_j\}; p_1, p_2) = \prod_{k=1}^m s_{i_k j_k}^{-\alpha_k} = \prod_{k=1}^m (2p_{i_k} \cdot p_{j_k})^{-\alpha_k}, \quad (5.4)$$

where the index k runs over all the two-particle invariants appearing in f . Every soft integral can be converted into an integral of this type, to the price of introducing additional MB integrations. Indeed, if we write every multi-particle invariant as a sum of two-particle invariants, then we can convert sums into product by using the usual formula

$$\frac{1}{(A+B)^\lambda} = \frac{1}{\Gamma(\lambda)} \int_{-i\infty}^{+i\infty} \frac{dz}{2\pi i} \Gamma(-z) \Gamma(\lambda+z) \frac{A^z}{B^{z+\lambda}}, \quad (5.5)$$

where the contour separates the poles at $z = n$ from those in $z = \lambda - n$, $n \in \mathbb{N}$. Without loss of generality we can thus assume that our soft integral is homogeneous with respect to individual rescalings of the final state momenta.

If we concentrate on soft integral that satisfy eq. (5.3), it is natural to choose a parametrization of the soft phase space that makes the homogeneity explicit. One possible parametrization with this property is the so-called ‘energies and angles’ parametrization, where the final-state momenta are parametrized as⁵

$$\begin{aligned} p_1 &= \frac{1}{2}(1, 1, 0, \dots), \\ p_2 &= \frac{1}{2}(1, -1, 0, \dots), \\ p_i &= \frac{1}{2} E_i \beta_i, \quad 3 \leq i \leq N. \end{aligned} \quad (5.6)$$

⁵We work with the rescaled momenta, eq. (2.6).

The E_i parametrize the energies of the final-state partons and β_i is the D -velocity in the direction p_i . In this parametrization the phase space measure for each final-state parton takes the form

$$d^D p_i \delta_+(p_i^2) = 2^{-(D-1)} \Theta(E_i) E_i^{D-3} dE_i d\Omega_i^{(D-1)}, \quad (5.7)$$

where $d\Omega_i^{(D-1)}$ is the measure on the unit sphere parametrizing the solid angle of particle i . Furthermore, the on-shell condition for the Higgs boson, coming from the cut propagator in eq. (2.16), can be rewritten as

$$\left[\frac{1}{p_{12} \cdot (p_{12} - 2p_{3\dots N})} \right]_c = \delta_+(p_{12} \cdot (p_{12} - 2p_{3\dots N})) = \delta \left(1 - \sum_{i=3}^N E_i \right). \quad (5.8)$$

Thus, the soft phase space measure can be parametrized as

$$d\Phi_{N-1}^S = (2\pi)^{N-1-(N-2)D} 2^{-(N-2)(D-1)} \delta \left(1 - \sum_{i=3}^N E_i \right) \prod_{i=3}^N E_i^{D-3} dE_i d\Omega_i^{(D-1)}. \quad (5.9)$$

Using this parametrization, we see that every soft integral with an integrand of the form (5.4) can be written as

$$F(\epsilon) = \int d\Phi_{N-1}^S \prod_{k=1}^m s_{i_k j_k}^{-\alpha_k} = 2^\alpha \int d\Phi_{N-1}^S \prod_{k=1}^m (E_{i_k} E_{j_k})^{-\alpha_k} (\beta_{i_k} \cdot \beta_{j_k})^{\alpha_k}, \quad \alpha = \sum_{k=1}^m \alpha_k. \quad (5.10)$$

Both the measure and the integrand can be written in a factorized form, and so we can integrate out the energies in terms of a generalized Beta function,

$$\int_0^1 \left(\prod_{k=3}^N dE_k E_k^{a_k-1} \right) \delta \left(1 - \sum_{k=3}^N E_k \right) = \frac{\Gamma(a_1) \dots \Gamma(a_m)}{\Gamma(a_1 + \dots + a_m)}. \quad (5.11)$$

Hence, the only non-trivial integration is a multiple angular integration over the solid angles of the final-state partons. Angular integrals can be written in the general form

$$\Omega_{D-1}^{(\alpha_1, \dots, \alpha_m)}(\{\beta_{j_a} \cdot \beta_{j_b}\}) = \int \frac{d\Omega_i^{(D-1)}}{(\beta_{j_1} \cdot \beta_i)^{\alpha_1} \dots (\beta_{j_m} \cdot \beta_i)^{\alpha_m}}. \quad (5.12)$$

In ref. [81] it was shown that such integrals fall into a class of generalized hypergeometric functions known as **H** functions, and an MB representation for the most general angular integral of this type was derived. We have thus a general recipe to derive MB representations for generic soft integrals.

Although the previous technique allows us to derive a multifold MB representation for every soft integral we need to consider, it can sometimes be useful to insert, if existent, explicit closed expressions for the angular integrals. Indeed, for small values of m the integrals (5.12) are very simple and can be evaluated in closed form. In the following we briefly review some results for angular integrals which will be useful in our case.

The case $m = 0$ corresponds to the volume of the solid angle

$$\Omega_{D-1} = \int d\Omega_i^{(D-1)} = \frac{2\pi^{(D-1)/2}}{\Gamma(\frac{D-1}{2})}. \quad (5.13)$$

Note that this integral is sufficient to compute the soft phase space volume, which corresponds to putting $m = 0$ in eq. (5.10): we simply obtain a factor Ω_{D-1} for each final-state parton. Thus, we obtain

$$\Phi_{N-1}^S(\epsilon) = (2\pi)^{-2N+5+(N-2)\epsilon} 2^{-(N-2)(2-\epsilon)} \frac{\Gamma(1-\epsilon)^{N-2}}{\Gamma(2(N-2)(1-\epsilon))}, \quad (5.14)$$

in agreement with the results of appendix B.

As we are only interested in massless momenta, $\beta_j^2 = 0$, Lorentz invariance implies that the angular integral with one propagator must evaluate to a constant. Indeed, we have

$$\Omega_{D-1}^{(\alpha)} = \int \frac{d\Omega_i^{(D-1)}}{(\beta_j \cdot \beta_i)^\alpha} = 2^{2-\alpha-2\epsilon} \pi^{1-\epsilon} \frac{\Gamma(1-\epsilon-\alpha)}{\Gamma(2-2\epsilon-\alpha)}. \quad (5.15)$$

For angular integrals with two massless propagators one obtains [82]

$$\begin{aligned} \Omega_{D-1}^{(\alpha_{j_1}, \alpha_{j_2})}(\beta_{j_1} \cdot \beta_{j_2}) &= \int \frac{d\Omega_i^{(D-1)}}{(\beta_{j_1} \cdot \beta_i)^{\alpha_1} (\beta_{j_2} \cdot \beta_i)^{\alpha_2}} \\ &= 2^{2-\alpha_1-\alpha_2-2\epsilon} \pi^{1-\epsilon} \frac{\Gamma(1-\epsilon-\alpha_1)\Gamma(1-\epsilon-\alpha_2)}{\Gamma(1-\epsilon)\Gamma(2-2\epsilon-\alpha_1-\alpha_2)} \\ &\quad \times {}_2F_1\left(\alpha_1, \alpha_2; 1-\epsilon; 1 - \frac{\beta_{j_1} \cdot \beta_{j_2}}{2}\right). \end{aligned} \quad (5.16)$$

To our knowledge, there are no closed formulas for angular integrals with three or more massless propagators. We will however need later on the angular integral with three massless propagators, which admits the MB representation [81],

$$\begin{aligned} \Omega_{D-1}^{(\alpha_{j_1}, \alpha_{j_2}, \alpha_{j_3})}(\beta_{j_1} \cdot \beta_{j_2}, \beta_{j_2} \cdot \beta_{j_3}, \beta_{j_1} \cdot \beta_{j_3}) &= \frac{2^{2-\alpha_1-\alpha_2-\alpha_3-2\epsilon} \pi^{1-\epsilon}}{\Gamma(\alpha_1)\Gamma(\alpha_2)\Gamma(\alpha_3)\Gamma(2-\alpha_1-\alpha_2-\alpha_3-2\epsilon)} \\ &\times \int_{-i\infty}^{+i\infty} \frac{dz_{12}dz_{13}dz_{23}}{(2\pi i)^3} \Gamma(-z_{12})\Gamma(-z_{13})\Gamma(-z_{23})\Gamma(\alpha_1+z_{12}+z_{13})\Gamma(\alpha_2+z_{12}+z_{23}) \\ &\quad \times \Gamma(\alpha_3+z_{13}+z_{23})\Gamma(1-\alpha_1-\alpha_2-\alpha_3-\epsilon-z_{12}-z_{13}-z_{23}) \\ &\quad \times \left(\frac{\beta_{j_1} \cdot \beta_{j_2}}{2}\right)^{z_{12}} \left(\frac{\beta_{j_1} \cdot \beta_{j_3}}{2}\right)^{z_{13}} \left(\frac{\beta_{j_2} \cdot \beta_{j_3}}{2}\right)^{z_{23}}, \end{aligned} \quad (5.17)$$

with $z_{ij} = z_i + z_j$, and where the contours separate the poles coming from Γ functions of the form $\Gamma(\dots - z_{ij})$ from those coming from $\Gamma(\dots + z_{ij})$.

6 Triple real emission phase space integrals in the soft limit

6.1 Triple real soft master integrals for Higgs production

After having studied some properties of soft integral in the previous sections, we will use the technology developed in the previous sections to compute the threshold expansion of the

leading-order cross sections for H plus five partons. More details about the construction of the amplitude in this limit will be given in section 7. Here it suffices to say that we have computed the squared amplitude and we have checked that in the limit where we only keep the first two terms in the threshold expansion, all the phase space integrals can be reduced to linear combinations of the following ten soft master integrals,

$$= \int d\Phi_4^S = \Phi_4^S(\epsilon), \quad (6.1)$$

$$= \int \frac{d\Phi_4^S}{(s_{13} + s_{15})s_{34}} = \Phi_4^S(\epsilon) \mathcal{F}_2(\epsilon), \quad (6.2)$$

$$= \int \frac{d\Phi_4^S}{s_{14}s_{23}s_{34}} = \Phi_4^S(\epsilon) \mathcal{F}_3(\epsilon), \quad (6.3)$$

$$= \int \frac{d\Phi_4^S}{s_{13}s_{15}s_{34}s_{45}} = \Phi_4^S(\epsilon) \mathcal{F}_4(\epsilon), \quad (6.4)$$

$$= \int \frac{d\Phi_4^S}{(s_{14} + s_{15})s_{23}s_{345}} = \Phi_4^S(\epsilon) \mathcal{F}_5(\epsilon), \quad (6.5)$$

$$= \int \frac{d\Phi_4^S}{(s_{13} + s_{14})(s_{14} + s_{15})s_{23}s_{34}} = \Phi_4^S(\epsilon) \mathcal{F}_6(\epsilon), \quad (6.6)$$

$$= \int \frac{d\Phi_4^S}{s_{15}s_{24}s_{34}s_{35}} = \Phi_4^S(\epsilon) \mathcal{F}_7(\epsilon), \quad (6.7)$$

$$= \int \frac{d\Phi_4^S}{(s_{13} + s_{15})(s_{23} + s_{24})s_{34}s_{35}} = \Phi_4^S(\epsilon) \mathcal{F}_8(\epsilon), \quad (6.8)$$

$$= \int \frac{d\Phi_4^S}{s_{15}(s_{14} + s_{15})s_{23}s_{34}s_{345}} = \Phi_4^S(\epsilon) \mathcal{F}_9(\epsilon), \quad (6.9)$$

$$= \int \frac{d\Phi_4^S}{(s_{23} + s_{24})(s_{24} + s_{25})s_{34}s_{45}} = \Phi_4^S(\epsilon) \mathcal{F}_{10}(\epsilon). \quad (6.10)$$

We have normalized all the integrals to the soft phase space volume for $H + 3g$ defined in eq. (3.16). In the remainder of this section we give the dimensional recurrence relations satisfied by the master integrals and present the analytic results for each master integral as a Laurent expansion in the dimensional regulator ϵ . Technical details about how to compute the master integrals analytically will be given in section 8.

6.2 Dimensional recurrence relations

Using the technique described in section 4, we can derive dimensional recurrence relations for all the master integrals defined in the previous section. The knowledge of these recurrence relations provides us with a strong check on our results. In addition, it turns out that the master integral $\mathcal{F}_9(D)$ is easier to compute in $D = 6 - 2\epsilon$ dimensions, where it is finite, and the dimensional recurrence relations allow us to relate the six-dimensional and four-dimensional results in an easy way.

The recurrence relation for the soft phase space volume is trivial to obtain from the recurrence relation for the Γ function,

$$\Phi_4^S(D+2) = \frac{(D-4)(D-3)(D-2)^3}{72(D-1)(3D-5)(3D-4)(3D-2)(3D-1)} \frac{\Gamma(D-4)}{64\pi^3\Gamma(D-1)} \Phi_4^S(D). \quad (6.11)$$

As we have defined all our master integrals relative to the phase space volume Φ_4^S , we can simplify their recurrence relations by factoring out the above result. We therefore define the ratio

$$\mathcal{R} = \frac{\mathcal{N}_3^D}{\mathcal{N}_3^{D+2}} \frac{\Phi_4^S(D+2)}{\Phi_4^S(D)} = \frac{(D-4)(D-3)(D-2)^3}{72(D-1)(3D-5)(3D-4)(3D-2)(3D-1)}, \quad (6.12)$$

where $\mathcal{N}$ was defined in eq. (4.4). We give the results for the remaining master integrals relative to $\mathcal{R}$. The dimensional recurrence relations for the non-trivial master integrals are

$$\mathcal{F}_2(D+2)\mathcal{R} = -\frac{(D-4)(7D-18)}{3(3D-5)(3D-4)} - \frac{(D-4)^2(3D-10)}{24(3D-7)(3D-5)(3D-4)} \mathcal{F}_2(D), \quad (6.13)$$

$$\begin{aligned} \mathcal{F}_3(D+2)\mathcal{R} &= \frac{(38-28D+5D^2)}{3(D-4)(3D-5)} \\ &\quad - \frac{(D-4)^3(D-3)}{18(3D-10)(3D-8)(3D-7)(3D-5)} \mathcal{F}_3(D), \end{aligned} \quad (6.14)$$

$$\begin{aligned} \mathcal{F}_4(D+2)\mathcal{R} &= -\frac{4(386-387D+128D^2-14D^3)}{(D-4)^2(D-3)} \\ &\quad - \frac{(D-4)^2(3D-14)}{24(3D-11)(3D-8)(3D-7)} \mathcal{F}_4(D), \end{aligned} \quad (6.15)$$

$$\begin{aligned}\mathcal{F}_5(D+2)\mathcal{R} = & -\frac{(D-4)(4752-9636D+6706D^2-1962D^3+207D^4)}{72(D-3)(D-1)(3D-10)(3D-8)(3D-5)} \\ & +\frac{(D-4)^2(D-2)}{96(D-1)(3D-7)(3D-5)}\mathcal{F}_5(D),\end{aligned}\quad (6.16)$$

$$\begin{aligned}\mathcal{F}_6(D+2)\mathcal{R} = & \frac{(4256-6684D+4224D^2-1345D^3+216D^4-14D^5)}{3(D-4)^2(D-3)(D-2)^2} \\ & +\frac{(D-4)(3D-10)}{9(D-2)^2(3D-7)}\mathcal{F}_2(D) \\ & -\frac{(D-4)^3}{24(D-2)(3D-11)(3D-7)}\mathcal{F}_6(D),\end{aligned}\quad (6.17)$$

$$\begin{aligned}\mathcal{F}_7(D+2)\mathcal{R} = & -\frac{4(2D-7)}{(D-4)(D-3)} \\ & +\frac{(D-4)^4}{72(3D-11)(3D-10)(3D-8)(3D-7)}\mathcal{F}_7(D),\end{aligned}\quad (6.18)$$

$$\begin{aligned}\mathcal{F}_8(D+2)\mathcal{R} = & \frac{2(231-114D+14D^2)}{3(D-4)(D-3)} + \frac{2(D-4)^2(7D-24)}{9(D-3)(3D-8)(3D-7)}\mathcal{F}_2(D) \\ & +\frac{(D-4)^4}{72(3D-11)(3D-10)(3D-8)(3D-7)}\mathcal{F}_8(D),\end{aligned}\quad (6.19)$$

$$\begin{aligned}\mathcal{F}_9(D+2)\mathcal{R} = & \frac{2(3D-7)(6672-7824D+3460D^2-684D^3+51D^4)}{3(D-4)^2(D-3)^2(3D-10)} \\ & +\frac{(D-4)(3D-10)(5D-17)}{12(D-3)^2(3D-8)}\mathcal{F}_2(D) + \frac{(D-4)}{6(D-3)(3D-8)}\mathcal{F}_5(D) \\ & +\frac{(D-4)^3(3D-14)}{96(D-3)(3D-13)(3D-11)(3D-8)}\mathcal{F}_9(D),\end{aligned}\quad (6.20)$$

$$\begin{aligned}\mathcal{F}_{10}(D+2)\mathcal{R} = & -\frac{4(26-39D+16D^2-2D^3)}{(D-4)^2(D-3)} - \frac{(D-4)^2(3D-10)}{3(D-3)(3D-8)(3D-7)}\mathcal{F}_2(D) \\ & -\frac{(D-4)^2(3D-14)}{24(3D-11)(3D-8)(3D-7)}\mathcal{F}_{10}(D).\end{aligned}\quad (6.21)$$

6.3 Analytic results for the soft master integrals

In this section we present the analytical results for the master integrals contributing to hadronic Higgs production in the leading and next-to-leading soft approximation. As the explicit evaluation of the master integrals is rather long and technical, we delay all details about the computation to section 8 and only summarize the results at this point. The first master integral, the soft phase space volume, was already given in eq. (3.16) and will not be repeated here. All the remaining master integrals have been evaluated as a Laurent series in the dimensional regulator up to terms involving zeta values of weight at most six. We have checked that our results agree numerically with the MB integral representation for soft integrals derived in section 5. In addition, the results satisfy the dimensional recurrence relations for the master integrals given in the previous section (integrals in the shifted dimension have been evaluated numerically using the MB representation). Finally, we make an intriguing observation in our results: if we express all the zeta values up to

weight six in the basis $\{\zeta_2, \zeta_3, \zeta_4, \zeta_2 \zeta_3, \zeta_5, \zeta_3^2, \zeta_6\}$, the coefficients in front of the values are integers in all cases. We note that this statement is only true in the specific basis of zeta values that we chose. The results for the master integrals are listed in the rest of this section.

$$\begin{aligned}
 \mathcal{F}_2(\epsilon) &= \frac{1}{\Phi_4^S(\epsilon)} \int \frac{d\Phi_4^S}{(s_{13} + s_{15})s_{34}} \\
 &= \frac{\Gamma(6-6\epsilon)\Gamma(1-2\epsilon)^2}{\epsilon\Gamma(3-6\epsilon)\Gamma(2-2\epsilon)^2} {}_3F_2(1, 1, 1-\epsilon; 2-2\epsilon, 2-2\epsilon; 1) \\
 &= \frac{60}{\epsilon} \zeta_2 + 420 \zeta_3 - 282 \zeta_2 + \epsilon \left(1800 \zeta_4 - 1974 \zeta_3 + 432 \zeta_2 \right) \\
 &\quad + \epsilon^2 \left(5580 \zeta_5 + 480 \zeta_2 \zeta_3 - 8460 \zeta_4 + 3024 \zeta_3 - 216 \zeta_2 \right) \\
 &\quad + \epsilon^3 \left(19260 \zeta_6 + 1680 \zeta_3^2 - 26226 \zeta_5 - 2256 \zeta_2 \zeta_3 + 12960 \zeta_4 - 1512 \zeta_3 \right) + \mathcal{O}(\epsilon^4). \tag{6.22}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{F}_3(\epsilon) &= \frac{1}{\Phi_4^S(\epsilon)} \int \frac{d\Phi_4^S}{s_{14}s_{23}s_{34}} \\
 &= \frac{90}{\epsilon^4} - \frac{693}{\epsilon^3} + \frac{1}{\epsilon^2} \left(-60 \zeta_2 + 1917 \right) + \frac{1}{\epsilon} \left(-300 \zeta_3 + 462 \zeta_2 - 2268 \right) \\
 &\quad - 930 \zeta_4 + 2310 \zeta_3 - 1278 \zeta_2 + 972 + \epsilon \left(-2220 \zeta_5 - 120 \zeta_2 \zeta_3 + 7161 \zeta_4 \right. \\
 &\quad \left. - 6390 \zeta_3 + 1512 \zeta_2 \right) + \epsilon^2 \left(-5555 \zeta_6 - 300 \zeta_3^2 + 17094 \zeta_5 + 924 \zeta_2 \zeta_3 \right. \\
 &\quad \left. - 19809 \zeta_4 + 7560 \zeta_3 - 648 \zeta_2 \right) + \mathcal{O}(\epsilon^3). \tag{6.23}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{F}_4(\epsilon) &= \frac{1}{\Phi_4^S(\epsilon)} \int \frac{d\Phi_4^S}{s_{13}s_{15}s_{34}s_{45}} \\
 &= -\frac{3\Gamma(6-6\epsilon)\Gamma(1-2\epsilon)}{2\epsilon^4\Gamma(1-6\epsilon)} \\
 &\quad \times \left[\frac{3\Gamma(1-2\epsilon)\Gamma(\epsilon+1)}{(1+3\epsilon)\Gamma(1-3\epsilon)} {}_3F_2(-3\epsilon-1, -2\epsilon, -\epsilon; -3\epsilon, -3\epsilon; 1) \right. \\
 &\quad \left. + \frac{1}{(1+\epsilon)\Gamma(1-2\epsilon)} {}_4F_3(1, 1, 1-\epsilon, -2\epsilon; 1-2\epsilon, 1-2\epsilon, 2+\epsilon; 1) \right] \\
 &= -\frac{600}{\epsilon^4} + \frac{10020}{\epsilon^3} - \frac{70560}{\epsilon^2} - \frac{1}{\epsilon} \left(480 \zeta_3 - 303480 \right) - 3600 \zeta_4 + 8016 \zeta_3 \\
 &\quad - 1007640 - \epsilon \left(17280 \zeta_5 - 60120 \zeta_4 + 56448 \zeta_3 - 3061800 \right) - \epsilon^2 \left(66000 \zeta_6 \right. \\
 &\quad \left. + 1920 \zeta_3^2 - 288576 \zeta_5 + 423360 \zeta_4 - 242784 \zeta_3 + 9185400 \right) + \mathcal{O}(\epsilon^3). \tag{6.24}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{F}_5(\epsilon) &= \frac{1}{\Phi_4^S(\epsilon)} \int \frac{d\Phi_4^S}{(s_{14} + s_{15})s_{23}s_{345}} \\
 &= -\frac{120}{\epsilon} \zeta_2 - 960 \zeta_3 + 684 \zeta_2 + \epsilon \left(-4620 \zeta_4 + 5472 \zeta_3 - 1188 \zeta_2 \right) \\
 &\quad + \epsilon^2 \left(-17160 \zeta_5 - 720 \zeta_2 \zeta_3 + 26334 \zeta_4 - 9504 \zeta_3 + 648 \zeta_2 \right) \\
 &\quad + \epsilon^3 \left(-64110 \zeta_6 - 2880 \zeta_3^2 + 97812 \zeta_5 + 4104 \zeta_2 \zeta_3 - 45738 \zeta_4 + 5184 \zeta_3 \right) + \mathcal{O}(\epsilon^4). \tag{6.25}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{F}_6(\epsilon) &= \frac{1}{\Phi_4^S(\epsilon)} \int \frac{d\Phi_4^S}{(s_{13} + s_{14})(s_{14} + s_{15})s_{23}s_{34}} \\
 &= \frac{10}{\epsilon^5} - \frac{137}{\epsilon^4} + \frac{1}{\epsilon^3} \left(40\zeta_2 + 675 \right) + \frac{1}{\epsilon^2} \left(320\zeta_3 - 548\zeta_2 - 1530 \right) \\
 &\quad + \frac{1}{\epsilon} \left(1500\zeta_4 - 4384\zeta_3 + 2700\zeta_2 + 1620 \right) + 5160\zeta_5 + 320\zeta_2\zeta_3 - 20550\zeta_4 \\
 &\quad + 21600\zeta_3 - 6120\zeta_2 - 648 + \epsilon \left(18340\zeta_6 + 1280\zeta_3^2 - 70692\zeta_5 - 4384\zeta_2\zeta_3 \right. \\
 &\quad \left. + 101250\zeta_4 - 48960\zeta_3 + 6480\zeta_2 \right) + \mathcal{O}(\epsilon^2). \tag{6.26}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{F}_7(\epsilon) &= \frac{1}{\Phi_4^S(\epsilon)} \int \frac{d\Phi_4^S}{s_{15}s_{24}s_{34}s_{35}} \\
 &= -\frac{3}{2} \frac{\Gamma(6-6\epsilon)}{\epsilon^5 \Gamma(1-6\epsilon)} {}_3F_2(1, 1, -2\epsilon; 1-2\epsilon, 1-2\epsilon; 1) \\
 &= -\frac{360}{\epsilon^5} + \frac{4932}{\epsilon^4} + \frac{1}{\epsilon^3} \left(720\zeta_2 - 24300 \right) + \frac{1}{\epsilon^2} \left(4320\zeta_3 - 9864\zeta_2 + 55080 \right) \\
 &\quad + \frac{1}{\epsilon} \left(15120\zeta_4 - 59184\zeta_3 + 48600\zeta_2 - 58320 \right) + 43200\zeta_5 - 207144\zeta_4 \\
 &\quad + 291600\zeta_3 - 110160\zeta_2 + 23328 + \epsilon \left(111600\zeta_6 - 591840\zeta_5 + 1020600\zeta_4 \right. \\
 &\quad \left. - 660960\zeta_3 + 116640\zeta_2 \right) + \mathcal{O}(\epsilon^2). \tag{6.27}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{F}_8(\epsilon) &= \frac{1}{\Phi_4^S(\epsilon)} \int \frac{d\Phi_4^S}{(s_{13} + s_{15})(s_{23} + s_{24})s_{34}s_{35}} \\
 &= -\frac{60}{\epsilon^5} + \frac{822}{\epsilon^4} + \frac{1}{\epsilon^3} \left(240\zeta_2 - 4050 \right) + \frac{1}{\epsilon^2} \left(2400\zeta_3 - 3288\zeta_2 + 9180 \right) \\
 &\quad + \frac{1}{\epsilon} \left(13320\zeta_4 - 32880\zeta_3 + 16200\zeta_2 - 9720 \right) + 51840\zeta_5 + 3360\zeta_2\zeta_3 \\
 &\quad - 182484\zeta_4 + 162000\zeta_3 - 36720\zeta_2 + 3888 + \epsilon \left(207600\zeta_6 + 11760\zeta_3^2 \right. \\
 &\quad \left. - 710208\zeta_5 - 46032\zeta_2\zeta_3 + 899100\zeta_4 - 367200\zeta_3 + 38880\zeta_2 \right) + \mathcal{O}(\epsilon^2). \tag{6.28}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{F}_9(\epsilon) &= \frac{1}{\Phi_4^S(\epsilon)} \int \frac{d\Phi_4^S}{s_{15}(s_{14} + s_{15})s_{23}s_{34}s_{345}} \\
 &= \frac{160}{\epsilon^5} - \frac{1712}{\epsilon^4} + \frac{1}{\epsilon^3} \left(-120\zeta_2 + 2784 \right) + \frac{1}{\epsilon^2} \left(-120\zeta_3 + 1284\zeta_2 + 31968 \right) \\
 &\quad + \frac{1}{\epsilon} \left(2520\zeta_4 + 1284\zeta_3 - 2088\zeta_2 - 216864 \right) + 15720\zeta_5 + 1920\zeta_2\zeta_3 \\
 &\quad - 26964\zeta_4 - 2088\zeta_3 - 23976\zeta_2 + 795744 + \epsilon \left(82520\zeta_6 + 9600\zeta_3^2 \right. \\
 &\quad \left. - 168204\zeta_5 - 20544\zeta_2\zeta_3 + 43848\zeta_4 - 23976\zeta_3 + 162648\zeta_2 - 2449440 \right) \\
 &\quad + \mathcal{O}(\epsilon^2). \tag{6.29}
 \end{aligned}$$

$$\begin{aligned}
\mathcal{F}_{10}(\epsilon) &= \frac{1}{\Phi_4^S(\epsilon)} \int \frac{d\Phi_4^S}{(s_{23} + s_{24})(s_{24} + s_{25})s_{34}s_{45}} \\
&= -\frac{120}{\epsilon^4} + \frac{2004}{\epsilon^3} - \frac{14112}{\epsilon^2} + \frac{1}{\epsilon} \left(240 \zeta_3 + 60696 \right) + 1980 \zeta_4 - 4008 \zeta_3 \\
&\quad - 201528 + \epsilon \left(6960 \zeta_5 + 1680 \zeta_2 \zeta_3 - 33066 \zeta_4 + 28224 \zeta_3 + 612360 \right) \\
&\quad + \epsilon^2 \left(32700 \zeta_6 + 6840 \zeta_3^2 - 116232 \zeta_5 - 28056 \zeta_2 \zeta_3 + 232848 \zeta_4 - 121392 \zeta_3 \right. \\
&\quad \left. - 1837080 \right) + \mathcal{O}(\epsilon^3).
\end{aligned} \tag{6.30}$$

7 The threshold approximation to the partonic $2 \rightarrow H + 3$ parton cross-section

7.1 General setup

The production of a Higgs boson in the collision of two hadrons h_1, h_2 is dominated by QCD processes. The hadronic cross-section is related to the partonic cross-section by the general factorisation formula

$$\sigma_{h_1+h_2 \rightarrow H+X} = \sum_{i,j} \int_0^1 dx_1 dx_2 f_i^{h_1}(x_1) f_j^{h_2}(x_2) \sigma_{i+j \rightarrow H}(M^2, x_1 x_2 S). \tag{7.1}$$

Here $f_i^h(x)$ are the parton-distribution functions for the parton i inside of the hadron h , $\sigma_{i+j \rightarrow H}$ is the partonic cross-section and S is the square of the total centre-of-mass energy of the hadronic system. The centre-of-mass energy squared of the partonic system is consequently given by $s = x_i x_j S$. The partonic cross-section is expanded perturbatively in the strong coupling constant α_S . The leading QCD contribution arises in the SM via a top-quark loop at $\mathcal{O}(\alpha_S^2)$.

We consider the Higgs boson to be relatively light compared to the top quark. This justifies to work in the limit of an infinite top quark mass and consider N_f light quarks. We describe the interaction of the Higgs boson with gluons by introducing the effective Lagrangian

$$\mathcal{L}_{\text{eff}} = -\frac{1}{4} c_H G_{\mu\nu}^a G^{a\mu\nu} H. \tag{7.2}$$

Here $G_{\mu\nu}^a$ denotes the gluon field strength tensor and H is the Higgs field. The Wilson coefficient c_H can be found, e.g. in ref. [83–85]. The next-to-next-to-leading (NNLO) order correction to the inclusive Higgs cross-section has been computed in the past by employing this effective theory. In this work we present a part of the next term in the perturbative series (N₃LO) in the effective theory.

At every order in perturbative QCD, the cross-section receives contributions from various real and virtual radiation processes. We consider only tree-level processes with three real-emission partons in the final state. A direct integration over the phase-space of the corresponding matrix elements is challenging. As we have discussed in the introduction, we pave the way towards the full computation by performing an expansion of the phase-space integrals around the kinematic limit where the Higgs boson is produced at threshold,

$s \sim M^2$. In this limit, all partons emitted in the final state are soft and their momenta vanish as $\bar{z} \rightarrow 0$. We expand the cross section in the small parameter $\bar{z}$ defined in (2.5). In the following, we shall present the leading and subleading terms in the threshold expansion.

$$\sigma_{ij \rightarrow H+X}(s, \bar{z}) = \bar{z}^{-1-6\epsilon} s^{3\epsilon} \sum_{k=0}^{\infty} \bar{z}^k \sigma_{ij \rightarrow H+X}^{S(k)}. \quad (7.3)$$

Although we considered in our calculation the Higgs boson H as a final state, we would like to stress the universality of our result for the leading term in the soft expansion for any other colorless final state produced by gluons in the initial state [86]. The subleading term in the soft expansion is no longer universal.

7.2 Calculation

To obtain the real-emission cross-section we generate Feynman diagrams using QGRAF [87] and compute squared matrix-elements using programs based on GiNAC [88] or FORM [89] and our own C++ code for color and spin algebra. We perform our calculation in Feynman gauge with D gluon polarizations in order to maintain a simple structure for the denominators of the squared amplitudes. To recover the result for physical gluon polarizations, we add matrix-elements with Faddeev-Popov ghosts as external states. We compare our results with a set of numeric cross-sections for different phase-space points obtained with MADGRAPH [90] and find perfect agreement.

In a next step we use reverse-unitarity and interpret the phase-space integrals as three-loop integrals as described in section 2 and we expand them into a Laurent series in $\bar{z}$. The leading and next-to-leading terms in this series are then referred to as the soft and next-to-soft limit of the real emission cross-section. We then derive integration-by-parts (IBP) identities for the scalar soft phase-space integrals. We reduce all integrals to a set of 10 master integrals by employing the Laporta algorithm [67]. We implemented this algorithm in a C++ code that was developed by us specifically for this project, as well as the program AIR [91]. The calculation of the remaining 10 integrals is discussed in section 8.

7.3 Leading soft contribution

In this section we present the leading soft contributions to the $ij \rightarrow H + X$ real-emission cross-sections, where X where represents the three final-state partons. To obtain a $\overline{\text{MS}}$ -renormalised quantity we redefine the strong coupling constant

$$\alpha_S = \alpha_S^R(\mu_R) e^{\epsilon \gamma_E} (4\pi \mu_R)^{-\epsilon}, \quad (7.4)$$

where $\gamma_E = \Gamma'(1)$ is the Euler-Mascheroni constant and μ_R is the renormalisation scale. In our calculation we consider N_f massless quark flavours. We include a flux factor of $\frac{1}{2s}$ and average over incoming colours and spins, leading to a factor of $\frac{1}{2(1-\epsilon)(N_c^2-1)}$ for gluons and $\frac{1}{2N_c}$ for (anti-) quarks. We abbreviate $C_F = \frac{N_c^2-1}{2N_c}$ and $C_A = N_c$. In the case that the phase-space integration contains n identical particles we include a symmetry factor of $\frac{1}{n!}$. For cross-sections with 4 (anti-) quarks we distinguish between the case of only one or two quark flavours. The two-flavour case is indicated by explicitly labeling the (anti-) quarks q_i

or $\bar{q}_i$. If the (anti-) quarks do not carry any label they are considered all to be of the same flavour. We present our results first in terms of the soft master integrals of section 6. The resulting expressions are valid to all orders in the dimensional regulator ϵ . Next, we insert the results for the soft master integrals and write the cross-section as a Laurent series in ϵ up to terms containing zeta values of weight six, keeping a factor corresponding to the volume of the soft phase-space $\Phi_4^S(\epsilon)$, eq. (3.16), unexpanded.

The cross-section containing only gluons is given by

$$\begin{aligned}
 \sigma_{gg \rightarrow H+ggg}^{S(0)} &= \frac{1}{3!} \frac{1}{8(1-\epsilon)^2(N_c^2-1)^2} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_A^4 C_F c_H^2 \\
 &\times \left\{ \mathcal{F}_1(\epsilon) 32 \frac{1}{(3-2\epsilon)^2 \epsilon^5 (2\epsilon^2 + \epsilon - 1)} \right. \\
 &\quad \times (33696\epsilon^{11} - 270000\epsilon^{10} + 1341936\epsilon^9 - 3650136\epsilon^8 \\
 &\quad + 4913370\epsilon^7 - 2001211\epsilon^6 - 3045896\epsilon^5 + 5040807\epsilon^4 \\
 &\quad - 3323131\epsilon^3 + 1144330\epsilon^2 - 197535\epsilon + 13050) \\
 &\quad + \mathcal{F}_2(\epsilon) \frac{64}{3} \frac{432\epsilon^7 + 288\epsilon^6 - 372\epsilon^5 - 188\epsilon^4 + 10\epsilon^3 + 317\epsilon^2 - 142\epsilon + 15}{\epsilon^2(4\epsilon^3 - 4\epsilon^2 - 5\epsilon + 3)} \\
 &\quad - \mathcal{F}_3(\epsilon) 288 \left(6\epsilon + \frac{1}{\epsilon} - 7 \right) \\
 &\quad - \mathcal{F}_5(\epsilon) 32 \frac{12\epsilon^5 + 10\epsilon^4 + 4\epsilon^3 - 49\epsilon^2 + 26\epsilon - 3}{\epsilon^2(\epsilon+1)(2\epsilon-3)} \\
 &\quad - \mathcal{F}_7(\epsilon) \frac{64}{3} (\epsilon-1) \\
 &\quad - \mathcal{F}_8(\epsilon) \frac{64}{3} (\epsilon-1) \\
 &\quad \left. + \mathcal{F}_9(\epsilon) 32 \frac{3\epsilon^2 - 2\epsilon - 1}{6\epsilon + 1} \right\} \\
 &= \frac{2^5}{3^4} \frac{1}{3!} \frac{1}{8(N_c^2-1)^2} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_A^4 C_F c_H^2 \\
 &\times \left\{ -\frac{218700}{\epsilon^5} + \frac{2554740}{\epsilon^4} + \frac{1}{\epsilon^3} (131220\zeta_2 - 9709605) \right. \\
 &\quad + \frac{1}{\epsilon^2} (782460\zeta_3 - 1630854\zeta_2 + 14950359) + \frac{1}{\epsilon} (2869830\zeta_4 - 9687762\zeta_3 \\
 &\quad + 6810588\zeta_2 - 8547924) + 8373780\zeta_5 + 301320\zeta_2\zeta_3 - 35377641\zeta_4 \\
 &\quad + 40216932\zeta_3 - 11741904\zeta_2 + 107996 + \epsilon(24995385\zeta_6 + 763020\zeta_3^2 \\
 &\quad - 103032486\zeta_5 - 3541644\zeta_2\zeta_3 + 145858644\zeta_4 - 68849712\zeta_3 \\
 &\quad \left. + 7687776\zeta_2 - 455984) + \mathcal{O}(\epsilon^2) \right\}.
 \end{aligned}
 \tag{7.5}$$

Note that the only colour coefficient of the tree-level matrix element for a Higgs plus five gluons is $C_A^4 C_F$ (see, e.g., ref. [92]). Obviously this fact is left unchanged by the soft limit.

In the case that two final-state partons are a quark anti-quark pair we sum over all possible quark flavours and obtain a factor N_f . The corresponding cross-section is then given by

$$\begin{aligned}
 \sigma_{gg \rightarrow H+gq\bar{q}}^{S(0)} &= \frac{1}{8(1-\epsilon)^2(N_c^2-1)^2} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_A C_F c_H^2 N_f \\
 &\times \left\{ C_A^2 \left[\mathcal{F}_1(\epsilon) 32 \frac{1}{(3-2\epsilon)^2(\epsilon-1)\epsilon^4(\epsilon+1)(2\epsilon-1)} \right. \right. \\
 &\quad \times (288\epsilon^{11} - 18288\epsilon^{10} + 129712\epsilon^9 - 347128\epsilon^8 + 408738\epsilon^7 - 107919\epsilon^6 \\
 &\quad - 271140\epsilon^5 + 359705\epsilon^4 - 210605\epsilon^3 + 66680\epsilon^2 - 10853\epsilon + 690) \\
 &\quad + \mathcal{F}_2(\epsilon) \frac{32}{3} \left(-\frac{108\epsilon^6 + 180\epsilon^5 - 201\epsilon^4 - 14\epsilon^3 + 67\epsilon^2 - 22\epsilon + 2}{-4\epsilon^5 + 8\epsilon^4 + \epsilon^3 - 8\epsilon^2 + 3\epsilon} \right) \\
 &\quad - \mathcal{F}_5(\epsilon) 32 \frac{\epsilon - 6\epsilon^2}{-2\epsilon^2 + \epsilon + 3} \left. \right] \\
 &\quad - \mathcal{F}_1(\epsilon) 64 \frac{72\epsilon^7 - 396\epsilon^6 + 982\epsilon^5 - 1377\epsilon^4 + 1134\epsilon^3 - 527\epsilon^2 + 122\epsilon - 10}{\epsilon^4(2\epsilon-3)} \Big\} \\
 &= \frac{2^5}{3^7} \frac{1}{8(N_c^2-1)^2} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_A C_F c_H^2 N_f \\
 &\times \left\{ \frac{153090}{\epsilon^4} - \frac{1604043}{\epsilon^3} + \frac{1}{\epsilon^2} (-29160\zeta_2 + 4903902) \right. \\
 &\quad + \frac{1}{\epsilon} (-204120\zeta_3 + 321732\zeta_2 - 4833675) - 874800\zeta_4 + 2252124\zeta_3 - 911088\zeta_2 \\
 &\quad + 203535 + \epsilon (-2711880\zeta_5 - 233280\zeta_2\zeta_3 + 9651960\zeta_4 - 6290136\zeta_3 - 492210\zeta_2 \\
 &\quad + 1667109) + \epsilon^2 (-9360360\zeta_6 - 816480\zeta_3^2 + 29921076\zeta_5 + 2573856\zeta_2\zeta_3 \\
 &\quad - 26589060\zeta_4 - 4323186\zeta_3 + 4693212\zeta_2 + 1294731) \\
 &\quad + 2C_A C_F \left[\frac{167670}{\epsilon^4} - \frac{1743039}{\epsilon^3} + \frac{1}{\epsilon^2} (-29160\zeta_2 + 5267592) + \frac{1}{\epsilon} (-204120\zeta_3 \right. \\
 &\quad + 321732\zeta_2 - 5183163) - 874800\zeta_4 + 2252124\zeta_3 - 911088\zeta_2 + 337959 \\
 &\quad + \epsilon (-2711880\zeta_5 - 233280\zeta_2\zeta_3 + 9651960\zeta_4 - 6290136\zeta_3 \\
 &\quad - 492210\zeta_2 + 1651749) \\
 &\quad + \epsilon^2 (-9360360\zeta_6 - 816480\zeta_3^2 + 29921076\zeta_5 + 2573856\zeta_2\zeta_3 - 26589060\zeta_4 \\
 &\quad \left. - 4323186\zeta_3 + 4693212\zeta_2 + 1284491) \right] + \mathcal{O}(\epsilon^3) \Big\}.
 \end{aligned}
 \tag{7.6}$$

In the soft limit only the processes with gluons in the initial state contribute. All other contributions containing at least one (anti-) quark in the initial state vanish in this limit. This can be understood in terms of soft factorisation: there is no born-level process with a massless initial state fermion in Higgs production.

7.4 Next-to-leading soft contribution

In this section we present our results for the next-to-leading term in the soft expansion of the real-emission cross sections. Only partonic subprocesses that have at least one gluon in the initial state give a non-zero contribution at this order. The results are

$$\begin{aligned}
 \sigma_{gg \rightarrow H+gg}^{S(1)} &= \frac{1}{3!} \frac{1}{8(1-\epsilon)^2(N_c^2-1)^2} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_A^4 C_F c_H^2 \quad (7.7) \\
 &\times \left\{ -\mathcal{F}_1(\epsilon) 16 \frac{1}{(3-2\epsilon)^2 \epsilon^5 (6\epsilon^3 + \epsilon^2 - 4\epsilon + 1)} \right. \\
 &\quad \times (202176\epsilon^{13} - 2464992\epsilon^{12} + 12587184\epsilon^{11} - 25936632\epsilon^{10} + 9778008\epsilon^9 \\
 &\quad + 44147940\epsilon^8 - 70980864\epsilon^7 + 27845080\epsilon^6 + 29415875\epsilon^5 - 41617041\epsilon^4 \\
 &\quad + 22513771\epsilon^3 - 6360715\epsilon^2 + 915690\epsilon - 52200) \\
 &\quad - \mathcal{F}_2(\epsilon) \frac{64}{3} \frac{1}{\epsilon^2(\epsilon+1)(2\epsilon-1)(6\epsilon^2-11\epsilon+3)} \\
 &\quad \times (432\epsilon^9 + 4824\epsilon^8 - 1008\epsilon^7 - 5758\epsilon^6 + 2213\epsilon^5 \\
 &\quad + 968\epsilon^4 + 872\epsilon^3 - 1180\epsilon^2 + 347\epsilon - 30) \\
 &\quad + \mathcal{F}_3(\epsilon) 192 \left(6\epsilon^2 + 11\epsilon + \frac{3}{\epsilon} - 20 \right) \\
 &\quad + \mathcal{F}_5(\epsilon) 64 \frac{72\epsilon^8 - 168\epsilon^7 + 74\epsilon^6 + 208\epsilon^5 - 132\epsilon^4 - 170\epsilon^3 + 151\epsilon^2 - 38\epsilon + 3}{\epsilon^2(\epsilon+1)(6\epsilon^2-11\epsilon+3)} \\
 &\quad + \mathcal{F}_7(\epsilon) \frac{8}{3} \frac{24\epsilon^4 + 119\epsilon^3 - 242\epsilon^2 + 115\epsilon - 16}{(1-3\epsilon)^2} \\
 &\quad + \mathcal{F}_8(\epsilon) \frac{8}{3} \frac{24\epsilon^4 + 119\epsilon^3 - 242\epsilon^2 + 115\epsilon - 16}{(1-3\epsilon)^2} \\
 &\quad \left. - \mathcal{F}_9(\epsilon) 32 \frac{3\epsilon^4 + 16\epsilon^3 - 19\epsilon^2 - 2\epsilon + 2}{18\epsilon^2 - 3\epsilon - 1} \right\} \\
 &= \frac{2^5}{3^4} \frac{1}{3!} \frac{1}{8(N_c^2-1)^2} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_A^4 C_F c_H^2 \\
 &\quad \left\{ \frac{437400}{\epsilon^5} - \frac{4934520}{\epsilon^4} + \frac{1}{\epsilon^3} (-262440\zeta_2 + 17287938) \right. \\
 &\quad + \frac{1}{\epsilon^2} (-1564920\zeta_3 + 3242268\zeta_2 - 21475314) + \frac{1}{\epsilon} (-5739660\zeta_4 \\
 &\quad + 19317204\zeta_3 - 13267368\zeta_2 + 4182774) - 16747560\zeta_5 - 602640\zeta_2\zeta_3 \\
 &\quad + 70752852\zeta_4 - 78931800\zeta_3 + 21368412\zeta_2 + 7340216 + \epsilon(-49990770\zeta_6 \\
 &\quad - 1526040\zeta_3^2 + 206978652\zeta_5 + 6976368\zeta_2\zeta_3 - 288414837\zeta_4 + 126945108\zeta_3 \\
 &\quad \left. - 10614588\zeta_2 + 1548816) + \mathcal{O}(\epsilon^2) \right\}.
 \end{aligned}$$

$$\begin{aligned}
 \sigma_{gg \rightarrow H+gq\bar{q}}^{S(1)} &= \frac{1}{8(1-\epsilon)^2(N_c^2-1)^2} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_F C_A c_H^2 N_f \\
 &\times \left\{ C_A^2 \left[-\mathcal{F}_1(\epsilon) 16 \frac{1}{(3-2\epsilon)^2(\epsilon-1)\epsilon^4(\epsilon+1)(2\epsilon-1)(3\epsilon-1)} \right. \right. \\
 &\quad \times (1728\epsilon^{13} - 117792\epsilon^{12} + 595824\epsilon^{11} - 489880\epsilon^{10} - 2308824\epsilon^9 \\
 &\quad + 5971472\epsilon^8 - 5007932\epsilon^7 - 440720\epsilon^6 + 4236833\epsilon^5 - 3757178\epsilon^4 \\
 &\quad + 1686250\epsilon^3 - 424272\epsilon^2 + 56321\epsilon - 3030) \\
 &\quad + \mathcal{F}_2(\epsilon) \frac{4}{9} \left(\frac{1}{-4\epsilon^5 + 8\epsilon^4 + \epsilon^3 - 8\epsilon^2 + 3\epsilon} \right) \\
 &\quad \times (720\epsilon^7 + 11784\epsilon^6 + 5072\epsilon^5 - 16002\epsilon^4 + 3569\epsilon^3 + 3603\epsilon^2 - 1729\epsilon + 183) \\
 &\quad + \mathcal{F}_5(\epsilon) \frac{4}{3} \frac{144\epsilon^6 + 96\epsilon^5 + 640\epsilon^4 - 368\epsilon^3 + 199\epsilon^2 - 80\epsilon + 9}{6\epsilon^4 - 5\epsilon^3 - 8\epsilon^2 + 3\epsilon} \\
 &\quad + \mathcal{F}_7(\epsilon) 8 \frac{(\epsilon-1)\epsilon}{9-27\epsilon} \\
 &\quad + \mathcal{F}_8(\epsilon) 8 \frac{(\epsilon-1)\epsilon}{9-27\epsilon} \\
 &\quad + \mathcal{F}_9(\epsilon) 2 \frac{\epsilon(-3\epsilon^2+2\epsilon+1)}{-54\epsilon^2+9\epsilon+3} \Big] \\
 &\quad + \left[\mathcal{F}_1(\epsilon) 32 \frac{1}{\epsilon^4(2\epsilon-3)(2\epsilon-1)(3\epsilon-1)} \right. \\
 &\quad \times (864\epsilon^{10} - 2736\epsilon^9 + 192\epsilon^8 + 14320\epsilon^7 - 37150\epsilon^6 + 49403\epsilon^5 \\
 &\quad - 38954\epsilon^4 + 18385\epsilon^3 - 4992\epsilon^2 + 708\epsilon - 40) \\
 &\quad \left. + \mathcal{F}_2(\epsilon) 16 \frac{12\epsilon^3 - 8\epsilon^2 - 5\epsilon + 1}{6\epsilon - 3} \right] \Big\} \\
 &= \frac{2^3}{3^7} \frac{1}{8(N_c^2-1)^2} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_F C_A c_H^2 N_f \\
 &\times \left\{ -\frac{1224720}{\epsilon^4} + \frac{12657384}{\epsilon^3} + \frac{1}{\epsilon^2} (58320\zeta_2 - 35969184) + \frac{1}{\epsilon} (408240\zeta_3 \right. \\
 &\quad - 526824\zeta_2 + 21935124) + 1837080\zeta_4 - 3658608\zeta_3 - 1156032\zeta_2 + 20102076 \\
 &\quad + \epsilon (2361960\zeta_5 + 2303640\zeta_2\zeta_3 - 16405416\zeta_4 - 8880516\zeta_3 + 21218868\zeta_2 \\
 &\quad - 16893276) + \epsilon^2 (10114875\zeta_6 + 8193960\zeta_3^2 - 17837172\zeta_5 - 22092588\zeta_2\zeta_3 \\
 &\quad - 39133206\zeta_4 + 154780416\zeta_3 - 51706116\zeta_2 - 19146300) \\
 &\quad + 2C_A C_F \left[-\frac{1341360}{\epsilon^4} + \frac{13827672}{\epsilon^3} + \frac{1}{\epsilon^2} (58320\zeta_2 - 39026448) + \frac{1}{\epsilon} (408240\zeta_3 \right. \\
 &\quad - 439344\zeta_2 + 23445720) + 1837080\zeta_4 - 3046248\zeta_3 - 1654668\zeta_2 + 19514088 \\
 &\quad + \epsilon (2361960\zeta_5 + 2303640\zeta_2\zeta_3 - 13781016\zeta_4 - 12370968\zeta_3 + 20772720\zeta_2 \\
 &\quad - 17445648) + \epsilon^2 (10114875\zeta_6 + 8193960\zeta_3^2 - 9701532\zeta_5 - 21392748\zeta_2\zeta_3 \\
 &\quad \left. - 54092286\zeta_4 + 151657380\zeta_3 - 49947768\zeta_2 - 18496864) \right] + \mathcal{O}(\epsilon^3) \Big\}.
 \end{aligned}
 \tag{7.8}$$

$$\begin{aligned}
 \sigma_{g\bar{q} \rightarrow H+gg\bar{q}}^{S(1)} &= \sigma_{g\bar{q} \rightarrow H+gg\bar{q}}^{S(1)} = \frac{1}{2!} \frac{1}{8(1-\epsilon)N_c(N_c^2-1)} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_F c_H^2 (C_A - 2C_F) \\
 &\times \left\{ C_A^4 \left[\mathcal{F}_1(\epsilon) \frac{2}{3} \frac{1}{\epsilon^5 (4\epsilon^2 - 8\epsilon + 3)} \right. \right. \\
 &\quad \times (864\epsilon^{10} - 28512\epsilon^9 + 240744\epsilon^8 - 952620\epsilon^7 + 2144508\epsilon^6 \\
 &\quad - 2951235\epsilon^5 + 2534353\epsilon^4 - 1341935\epsilon^3 + 417659\epsilon^2 - 68326\epsilon + 4380) \\
 &\quad + \mathcal{F}_2(\epsilon) \frac{2}{9} \frac{324\epsilon^5 - 72\epsilon^4 - 567\epsilon^3 + 617\epsilon^2 - 201\epsilon + 19}{\epsilon^2(2\epsilon - 1)} \\
 &\quad + \mathcal{F}_3(\epsilon) 12 \left(6\epsilon + \frac{1}{\epsilon} - 7 \right) \\
 &\quad + \mathcal{F}_5(\epsilon) \frac{24\epsilon^5 - 112\epsilon^4 + 162\epsilon^3 - 48\epsilon^2 - 32\epsilon + 6}{9\epsilon^2 - 6\epsilon^3} \\
 &\quad + \mathcal{F}_7(\epsilon) \frac{4}{9} (\epsilon - 1) \\
 &\quad + \mathcal{F}_8(\epsilon) \frac{4}{9} (\epsilon - 1) \\
 &\quad \left. + \mathcal{F}_9(\epsilon) \frac{-3\epsilon^2 + 2\epsilon + 1}{6\epsilon + 1} \right] \\
 &+ C_A^2 \left[- \mathcal{F}_1(\epsilon) \frac{2}{3} \frac{1}{\epsilon^5 (4\epsilon^2 - 8\epsilon + 3)} \right. \\
 &\quad \times (1296\epsilon^{10} - 68256\epsilon^9 + 531432\epsilon^8 - 1871976\epsilon^7 + 3715893\epsilon^6 \\
 &\quad - 4517352\epsilon^5 + 3465908\epsilon^4 - 1668835\epsilon^3 + 482104\epsilon^2 - 74714\epsilon + 4620) \\
 &\quad + \mathcal{F}_2(\epsilon) \frac{2}{3} \frac{108\epsilon^5 - 129\epsilon^3 + 75\epsilon^2 - 15\epsilon + 1}{\epsilon^2(2\epsilon - 1)} \\
 &\quad + \mathcal{F}_4(\epsilon) \left(\epsilon - \frac{1}{3\epsilon} - \frac{2}{3} \right) \\
 &\quad + \mathcal{F}_5(\epsilon) \frac{2}{3} \left(-6\epsilon^2 - \frac{3}{\epsilon^2} + 43\epsilon + \frac{27}{\epsilon} - 61 \right) \\
 &\quad - \mathcal{F}_7(\epsilon) \frac{4}{9} (\epsilon - 1) \\
 &\quad + \mathcal{F}_9(\epsilon) \frac{3\epsilon^2 - 2\epsilon - 1}{18\epsilon + 3} \\
 &\quad + \left[\mathcal{F}_1(\epsilon) 2 \frac{1}{\epsilon^5(2\epsilon - 1)} \right. \\
 &\quad \times (72\epsilon^9 - 1332\epsilon^8 - 3014\epsilon^7 + 22217\epsilon^6 - 39799\epsilon^5 \\
 &\quad + 34159\epsilon^4 - 15710\epsilon^3 + 3841\epsilon^2 - 454\epsilon + 20) \\
 &\quad \left. \left. + \mathcal{F}_4(\epsilon) \left(\epsilon - \frac{1}{3\epsilon} - \frac{2}{3} \right) \right] \right\}
 \end{aligned} \tag{7.9}$$

$$\begin{aligned}
 &= \frac{2^2}{3^6} \frac{1}{2!} \frac{1}{8N_c(N_c^2 - 1)} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_F c_H^2 (C_A - 2C_F) \\
 &\times \left\{ \frac{277020}{\epsilon^5} - \frac{3661524}{\epsilon^4} + \frac{1}{\epsilon^3} (-189540\zeta_2 + 17139195) \right. \\
 &\quad + \frac{1}{\epsilon^2} (-947700\zeta_3 + 2489778\zeta_2 - 36505620) + \frac{1}{\epsilon} (-2850390\zeta_4 + 12171870\zeta_3 \\
 &\quad - 11487906\zeta_2 + 36053523) - 5846580\zeta_5 - 554040\zeta_2\zeta_3 + 35305713\zeta_4 - 54062316\zeta_3 \\
 &\quad + 23579532\zeta_2 - 12809922 + \epsilon (-12501135\zeta_6 - 1006020\zeta_3^2 + 66995586\zeta_5 \\
 &\quad + 6861348\zeta_2\zeta_3 - 146716434\zeta_4 + 103965822\zeta_3 - 20700180\zeta_2 - 893224) \\
 &\quad + C_A C_F \left[\frac{1370520}{\epsilon^5} - \frac{17969364}{\epsilon^4} + \frac{1}{\epsilon^3} (-962280\zeta_2 + 83256498) + \frac{1}{\epsilon^2} (-5277960\zeta_3 \right. \\
 &\quad + 12711816\zeta_2 - 175936887) + \frac{1}{\epsilon} (-17787600\zeta_4 + 68755392\zeta_3 - 59276286\zeta_2 \\
 &\quad + 174582603) - 45927000\zeta_5 - 2391120\zeta_2\zeta_3 + 227399400\zeta_4 - 313487658\zeta_3 \\
 &\quad + 124360218\zeta_2 - 65975079 + \epsilon (-124170570\zeta_6 - 5248800\zeta_3^2 + 572721840\zeta_5 \\
 &\quad \left. + 29492424\zeta_2\zeta_3 - 1004568156\zeta_4 + 635422050\zeta_3 - 116229744\zeta_2 - 466460) \right] \\
 &\quad + 6C_A^2 C_F^2 \left[\frac{291600}{\epsilon^5} - \frac{3785940}{\epsilon^4} + \frac{1}{\epsilon^3} (-194400\zeta_2 + 17329869) + \frac{1}{\epsilon^2} (-1108080\zeta_3 \right. \\
 &\quad + 2577420\zeta_2 - 36309654) + \frac{1}{\epsilon} (-3883140\zeta_4 + 14537556\zeta_3 - 12100158\zeta_2 + 36217701) \\
 &\quad - 10711440\zeta_5 - 427680\zeta_2\zeta_3 + 50265198\zeta_4 - 67142142\zeta_3 + 25733718\zeta_2 - 14421072 \\
 &\quad + \epsilon (-30383100\zeta_6 - 1001160\zeta_3^2 + 136655748\zeta_5 + 5256576\zeta_2\zeta_3 - 227203596\zeta_4 \\
 &\quad \left. + 139522482\zeta_3 - 24943128\zeta_2 + 612040) \right] + \mathcal{O}(\epsilon^2) \Big\}.
 \end{aligned}$$

$$\begin{aligned}
 \sigma_{gq \rightarrow H+q q \bar{q}}^{S(1)} &= \sigma_{g\bar{q} \rightarrow H+q \bar{q} \bar{q}}^{S(1)} = \frac{1}{2!} \frac{1}{8(1-\epsilon)N_c(N_c^2-1)} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_F c_H^2 (C_A - 2C_F) \\
 &\times \left\{ C_A^3 \left[-\mathcal{F}_1(\epsilon) 8 \frac{36\epsilon^7 - 1908\epsilon^6 + 7349\epsilon^5 - 11636\epsilon^4 + 9364\epsilon^3 - 3952\epsilon^2 + 807\epsilon - 60}{\epsilon^4(2\epsilon-3)} \right. \right. \\
 &\quad \left. \left. - \mathcal{F}_5(\epsilon) \frac{8}{3} \frac{6\epsilon^3 - 13\epsilon^2 + 8\epsilon - 1}{\epsilon(2\epsilon-3)} \right] \right. \\
 &\quad + C_A^2 \left[-\mathcal{F}_1(\epsilon) \frac{2}{3} \frac{1}{\epsilon^5(2\epsilon-1)} \right. \\
 &\quad \times (216\epsilon^9 - 1188\epsilon^8 + 6834\epsilon^7 - 17631\epsilon^6 + 22878\epsilon^5 \\
 &\quad \left. - 17064\epsilon^4 + 7853\epsilon^3 - 2212\epsilon^2 + 334\epsilon - 20) \right. \\
 &\quad + \mathcal{F}_2(\epsilon) \frac{2}{9} \frac{216\epsilon^5 - 306\epsilon^4 + 165\epsilon^3 - 110\epsilon^2 + 39\epsilon - 4}{\epsilon^2(2\epsilon-1)} \\
 &\quad + \mathcal{F}_5(\epsilon) \frac{2}{3} \left(12\epsilon^2 + \frac{2}{\epsilon^2} - 20\epsilon - \frac{15}{\epsilon} + 21 \right) \\
 &\quad - \mathcal{F}_8(\epsilon) \frac{4}{9} (\epsilon - 1) \\
 &\quad \left. + \mathcal{F}_{10}(\epsilon) \left(\epsilon - \frac{1}{3\epsilon} - \frac{2}{3} \right) \right] \\
 &\quad + C_A \left[\mathcal{F}_1(\epsilon) 8 \frac{36\epsilon^7 - 288\epsilon^6 + 869\epsilon^5 - 1331\epsilon^4 + 1129\epsilon^3 - 527\epsilon^2 + 122\epsilon - 10}{\epsilon^4(2\epsilon-3)} \right] \\
 &\quad + \left[\mathcal{F}_1(\epsilon) 2 \frac{1}{\epsilon^5(2\epsilon-1)} \right. \\
 &\quad \times (72\epsilon^9 - 396\epsilon^8 - 1178\epsilon^7 + 7947\epsilon^6 - 15222\epsilon^5 \\
 &\quad \left. + 14472\epsilon^4 - 7593\epsilon^3 + 2212\epsilon^2 - 334\epsilon + 20) \right. \\
 &\quad + \mathcal{F}_2(\epsilon) 2 \frac{18\epsilon^3 - 27\epsilon^2 + 10\epsilon - 1}{\epsilon(2\epsilon-1)} \\
 &\quad \left. \left. + \mathcal{F}_{10}(\epsilon) \left(\epsilon - \frac{1}{3\epsilon} - \frac{2}{3} \right) \right] \right\} \quad (7.10)
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{2}{3^6} \frac{1}{2!} \frac{1}{8N_c(N_c^2 - 1)} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_F c_H^2 \\
 &\times \left\{ \begin{aligned}
 &\frac{1}{\epsilon} (14580\zeta_4 - 87480\zeta_3) + 174960\zeta_5 - 58320\zeta_2\zeta_3 \\
 &- 943326\zeta_4 + 965196\zeta_3 + \epsilon(767880\zeta_6 - 612360\zeta_3^2 - 6770952\zeta_5 + 973944\zeta_2\zeta_3 \\
 &+ 9188316\zeta_4 - 3175524\zeta_3) \\
 &+ 4C_A \left[-\frac{12150}{\epsilon^4} + \frac{146205}{\epsilon^3} + \frac{1}{\epsilon^2} (9720\zeta_2 - 580500) + \frac{1}{\epsilon} (77760\zeta_3 - 116964\zeta_2 + 936135) \right. \\
 &+ 374220\zeta_4 - 935712\zeta_3 + 464400\zeta_2 - 556890 + \epsilon(1389960\zeta_5 + 58320\zeta_2\zeta_3 - 4503114\zeta_4 \\
 &\left. + 3715200\zeta_3 - 748908\zeta_2 + 22400) \right] \\
 &+ 2C_A C_F \left[\frac{1}{\epsilon^2} (29160\zeta_3 - 43740\zeta_2 + 47385) + \frac{1}{\epsilon} (255150\zeta_4 - 793152\zeta_3 + 511758\zeta_2 \right. \\
 &- 330237) + 1020600\zeta_5 + 145800\zeta_2\zeta_3 - 5551335\zeta_4 + 6515802\zeta_3 - 1928934\zeta_2 + 309825 \\
 &+ \epsilon(4740930\zeta_6 + 218700\zeta_3^2 - 22424040\zeta_5 - 2172420\zeta_2\zeta_3 + 40779531\zeta_4 - 21139542\zeta_3 \\
 &\left. + 2834352\zeta_2 - 174960) \right] \\
 &+ 24C_A^2 C_F \left[-\frac{4860}{\epsilon^4} + \frac{57267}{\epsilon^3} + \frac{1}{\epsilon^2} (3240\zeta_2 - 221427) + \frac{1}{\epsilon} (25920\zeta_3 - 38988\zeta_2 + 351261) \right. \\
 &+ 124740\zeta_4 - 311904\zeta_3 + 154800\zeta_2 - 209463 + \epsilon(463320\zeta_5 + 19440\zeta_2\zeta_3 - 1501038\zeta_4 \\
 &\left. + 1238400\zeta_3 - 249636\zeta_2 + 11990) \right] + \mathcal{O}(\epsilon^2) \Big\}.
 \end{aligned}
 \right.
 \end{aligned}$$

$$\begin{aligned}
 \sigma_{g \, q_1 \rightarrow H+q_1 \, q_2 \, \bar{q}_2}^{S(1)} &= \sigma_{g \, \bar{q}_1 \rightarrow H+q_2 \, \bar{q}_2 \, \bar{q}_1}^{S(1)} = \frac{1}{8(1-\epsilon)N_c(N_c^2-1)} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_F c_H^2 (N_f-1) \\
 &\times \left\{ C_A^2 \left[-\mathcal{F}_1(\epsilon) 4 \frac{36\epsilon^7 - 1908\epsilon^6 + 7349\epsilon^5 - 11636\epsilon^4 + 9364\epsilon^3 - 3952\epsilon^2 + 807\epsilon - 60}{\epsilon^4(2\epsilon-3)} \right. \right. \\
 &\quad \left. \left. - \mathcal{F}_5(\epsilon) \frac{4}{3} \frac{6\epsilon^3 - 13\epsilon^2 + 8\epsilon - 1}{\epsilon(2\epsilon-3)} \right] \right. \\
 &\quad \left. + \left[\mathcal{F}_1(\epsilon) 4 \frac{36\epsilon^7 - 288\epsilon^6 + 869\epsilon^5 - 1331\epsilon^4 + 1129\epsilon^3 - 527\epsilon^2 + 122\epsilon - 10}{\epsilon^4(2\epsilon-3)} \right] \right\} \\
 &= \frac{2^2}{3^7} \frac{1}{8N_c(N_c^2-1)} (4\pi\alpha_S)^3 \Phi_4^S(\epsilon) C_F c_H^2 (N_f-1) \\
 &\times \left\{ -\frac{36450}{\epsilon^4} + \frac{438615}{\epsilon^3} + \frac{1}{\epsilon^2} (29160\zeta_2 - 1741500) \right. \\
 &\quad + \frac{1}{\epsilon} (233280\zeta_3 - 350892\zeta_2 + 2808405) + 1122660\zeta_4 - 2807136\zeta_3 + 1393200\zeta_2 \\
 &\quad - 1670670 + \epsilon (4169880\zeta_5 + 174960\zeta_2\zeta_3 - 13509342\zeta_4 + 11145600\zeta_3 \\
 &\quad - 2246724\zeta_2 + 67200) + \epsilon^2 (15578730\zeta_6 + 699840\zeta_3^2 \\
 &\quad - 50177556\zeta_5 - 2105352\zeta_2\zeta_3 + 53638200\zeta_4 - 17973792\zeta_3 + 1336536\zeta_2 + 44800) \\
 &\quad + 6C_A C_F \left[-\frac{14580}{\epsilon^4} + \frac{171801}{\epsilon^3} + \frac{1}{\epsilon^2} (9720\zeta_2 - 664281) + \frac{1}{\epsilon} (77760\zeta_3 - 116964\zeta_2 \right. \\
 &\quad + 1053783) + 374220\zeta_4 - 935712\zeta_3 + 464400\zeta_2 - 628389 + \epsilon (1389960\zeta_5 \\
 &\quad + 58320\zeta_2\zeta_3 - 4503114\zeta_4 + 3715200\zeta_3 - 748908\zeta_2 + 35970) \\
 &\quad + \epsilon^2 (5192910\zeta_6 + 233280\zeta_3^2 - 16725852\zeta_5 - 701784\zeta_2\zeta_3 \\
 &\quad \left. \left. + 17879400\zeta_4 - 5991264\zeta_3 + 445512\zeta_2 + 15232) \right] + \mathcal{O}(\epsilon^2) \right\}.
 \end{aligned} \tag{7.11}$$

8 Analytic computation of the master integrals

In this section we present some details on how to evaluate analytically all the master integrals $\mathcal{F}_i$ defined in section 6. The analytic result for the soft phase space volume can easily be obtained from the expression for the phase space volume in general kinematics and will not be discussed here (see appendix B for the derivation).

In general, our strategy is to follow the steps outlined in section 5: we use the ‘energies and angles’ parametrization to obtain a representation for the integral where the energies and angles appear in a factorized form. This may require the introduction of MB integrations in order to factorize sums of invariants in a denominator. We then integrate out the energies and angles to obtain a multifold MB representation for each master integral. Whenever we are able to do so, we evaluate the remaining MB integrals to all orders in ϵ in terms of hypergeometric functions that can easily be expanded into a Laurent series ϵ using the `HypExp` package [93]. In those cases where we did not manage to perform the

MB integral in closed form for finite values of ϵ , we only compute the Laurent expansion of the integral around $\epsilon = 0$, e.g., by resolving singularities in ϵ and summing up harmonic sums or by converting the MB integral to a parametric integral which can be computed more easily. Details on how to perform these steps for the different master integrals will be given in the rest of this section.

8.1 The master integral $\mathcal{F}_3$

In this section we compute the master integral $\mathcal{F}_3$ defined by

$$\Phi_4^S(\epsilon) \mathcal{F}_3(\epsilon) = \int \frac{d\Phi_4^S}{s_{14}s_{23}s_{34}}. \quad (8.1)$$

The integrand only involves two-particle invariants, so following the discussion in section 5 we can immediately insert the energies and angles parametrization and integrate out the energies in terms of Γ functions. The remaining angular integrations can easily be carried out using the formulas given in section 5. Note that in the present case all the angular integrals can be performed in closed form, so there is no need to introduce MB representations for the angular integrals. We obtain

$$\begin{aligned} \mathcal{F}_3(\epsilon) &= 2^{6\epsilon-7} \pi^{3\epsilon-3} \frac{\Gamma(6-6\epsilon)\Gamma(2-2\epsilon)\Gamma(1-2\epsilon)^2}{\epsilon^2 \Gamma(2-6\epsilon)\Gamma(1-\epsilon)^3} \int \frac{d\Omega_3^{(D-1)} d\Omega_4^{(D-1)} d\Omega_5^{(D-1)}}{(\beta_1 \cdot \beta_4)(\beta_2 \cdot \beta_3)(\beta_3 \cdot \beta_4)} \\ &= 2^{6\epsilon-7} \pi^{3\epsilon-3} \frac{\Gamma(6-6\epsilon)\Gamma(2-2\epsilon)\Gamma(1-2\epsilon)^2}{\epsilon^2 \Gamma(2-6\epsilon)\Gamma(1-\epsilon)^3} \Omega_{3-2\epsilon} \int \frac{d\Omega_3^{(D-1)}}{(\beta_2 \cdot \beta_3)} \Omega_{D-1}^{(1,1)}(\beta_1 \cdot \beta_3) \\ &= -2^{2\epsilon-1} \frac{\Gamma(6-6\epsilon)\Gamma(1-2\epsilon)}{\epsilon^3 \Gamma(2-6\epsilon)\Gamma(1-\epsilon)^2} \int_{-1}^1 \frac{d \cos \theta_3}{(1 + \cos \theta_3)^{1+\epsilon}(1 - \cos \theta_3)^\epsilon} \\ &\quad \times {}_2F_1\left(1, 1; 1 - \epsilon; \frac{1 + \cos \theta_3}{2}\right). \end{aligned} \quad (8.2)$$

The remaining integral can be brought into a more standard form by the change of variable $\cos \theta_3 = 2y - 1$,

$$\begin{aligned} &\int_{-1}^1 \frac{d \cos \theta_3}{(1 + \cos \theta_3)^{1+\epsilon}(1 - \cos \theta_3)^\epsilon} {}_2F_1\left(1, 1; 1 - \epsilon; \frac{1 + \cos \theta_3}{2}\right) \\ &= 2^{-2\epsilon} \int_0^1 dy y^{-1-\epsilon} (1-y)^{-\epsilon} {}_2F_1(1, 1; 1 - \epsilon; y). \end{aligned} \quad (8.3)$$

The integral over y is now easily performed using the recursive definition of the hypergeometric function

$$\begin{aligned} {}_{p+1}F_p(a_1, \dots, a_{p+1}; b_1, \dots, b_p; z) &= \frac{\Gamma(b_p)}{\Gamma(a_{p+1})\Gamma(b_p - a_{p+1})} \\ &\quad \times \int_0^1 dt t^{a_{p+1}-1} (1-t)^{b_p-a_{p+1}-1} {}_pF_{p-1}(a_1, \dots, a_p; b_1, \dots, b_{p-1}; zt). \end{aligned} \quad (8.4)$$

We immediately get

$$\mathcal{F}_3(\epsilon) = \frac{\Gamma(6-6\epsilon)}{2\epsilon^4 \Gamma(2-6\epsilon)} {}_3F_2(1, 1, -\epsilon; 1-2\epsilon, 1-\epsilon; 1). \quad (8.5)$$

The ${}_3F_2$ function can be expanded to the desired order in ϵ using the **HypExp** package [93], and we arrive immediately at the Laurent series of eq. (6.23).

8.2 The master integral $\mathcal{F}_2$

The integrand of the master integral $\mathcal{F}_2$ involves a sum of two-particle invariants in the denominator. We replace the sum by a product to the price of introducing a MB integration via eq. (5.5),

$$\Phi_4^S(\epsilon) \mathcal{F}_2(\epsilon) = \int \frac{d\Phi_4^S}{(s_{13} + s_{15})s_{34}} = \int_{-i\infty}^{+i\infty} \frac{dz_1}{2\pi i} \Gamma(-z_1) \Gamma(z_1 + 1) \int \frac{d\Phi_4^S}{s_{13}^{z_1+1} s_{15}^{-z_1} s_{34}}. \quad (8.6)$$

The phase space integral is now in the form (5.10), and so we can introduce the energies and angles parametrization and integrate out all the energy and the angular variables. This results in the following two-fold integral representation for $\mathcal{F}_2$, which is of mixed MB and Euler-type,

$$\begin{aligned} \mathcal{F}_2(\epsilon) &= \frac{\Gamma(6-6\epsilon)\Gamma(1-2\epsilon)}{\epsilon\Gamma(3-6\epsilon)\Gamma(1-\epsilon)^4} \int_{-i\infty}^{+i\infty} \frac{dz_1}{2\pi i} \Gamma(-z_1) \Gamma(z_1 + 1) \Gamma(-\epsilon - z_1) \\ &\quad \times \Gamma(1 - \epsilon + z_1) \int_0^1 dy y^{-\epsilon} (1-y)^{-\epsilon} {}_2F_1(1, z_1 + 1; 1 - \epsilon; y). \end{aligned} \quad (8.7)$$

The Euler integral over y could immediately be performed in terms of a ${}_3F_2$ function, but after that we still need to integrate over the MB parameter z_1 . We therefore prefer not to perform the integration over y , but we rather insert the MB representation for the hypergeometric function in the integrand

$$\begin{aligned} &{}_pF_q(a_1, \dots, a_p; b_1, \dots, b_q; x) \\ &= \int_{-i\infty}^{+i\infty} \frac{dz}{2\pi i} (-x)^z \Gamma(-z) \left[\prod_{i=1}^p \frac{\Gamma(a_i + z)}{\Gamma(a_i)} \right] \left[\prod_{i=1}^q \frac{\Gamma(b_i)}{\Gamma(b_i + z)} \right]. \end{aligned} \quad (8.8)$$

The integral over y evaluates to a Beta function, and we are left with the following two-dimensional MB integral

$$\begin{aligned} \mathcal{F}_2(\epsilon) &= \frac{\Gamma(6-6\epsilon)\Gamma(1-2\epsilon)}{\epsilon\Gamma(3-6\epsilon)\Gamma(1-\epsilon)^2} \int_{-i\infty}^{+i\infty} \frac{dz_1 dz_2}{(2\pi i)^2} (-1)^{z_2} \Gamma(-z_1) \Gamma(-z_2) \\ &\quad \times \frac{\Gamma(z_2 + 1) \Gamma(z_1 + z_2 + 1) \Gamma(-\epsilon - z_1) \Gamma(1 - \epsilon + z_1)}{\Gamma(2 - 2\epsilon + z_2)}. \end{aligned} \quad (8.9)$$

The integral over z_1 is easily performed using Barnes' first lemma, and the remaining one-fold MB integral can immediately be recognized as a ${}_3F_2$ function (see eq. (8.8)). We finally obtain the following result for the master integral $\mathcal{F}_2$, in agreement with eq. (6.22),

$$\mathcal{F}_2(\epsilon) = \frac{\Gamma(6-6\epsilon)\Gamma(1-2\epsilon)^2}{\epsilon\Gamma(3-6\epsilon)\Gamma(2-2\epsilon)^2} {}_3F_2(1, 1, 1 - \epsilon; 2 - 2\epsilon, 2 - 2\epsilon; 1). \quad (8.10)$$

8.3 The master integral $\mathcal{F}_7$

The integrand of $\mathcal{F}_7$ only contains two-particle invariants,

$$\Phi_4^S(\epsilon) \mathcal{F}_7(\epsilon) = \int \frac{d\Phi_4^S}{s_{15} s_{24} s_{34} s_{35}}. \quad (8.11)$$

We can therefore immediately integrate out the energy and the angular variables. We obtain

$$\mathcal{F}_7(\epsilon) = 3 \frac{\Gamma(6-6\epsilon)\Gamma(1-2\epsilon)}{\epsilon^4 \Gamma(1-\epsilon)^2 \Gamma(1-6\epsilon)} \int_0^1 dy y^{-\epsilon} (1-y)^{-\epsilon} \times {}_2F_1(1, 1; 1-\epsilon; 1-y) {}_2F_1(1, 1; 1-\epsilon; y). \quad (8.12)$$

In order to perform the integral over y , we introduce an MB representation for each ${}_2F_1$ function in the integrand and perform the y -integration. This leaves us with the following two-fold MB representation for $\mathcal{F}_7$,

$$\mathcal{F}_7(\epsilon) = 3 \frac{\Gamma(6-6\epsilon)\Gamma(1-2\epsilon)}{\epsilon^4 \Gamma(1-\epsilon)^2 \Gamma(1-6\epsilon)} \times \int_{-i\infty}^{+i\infty} \frac{dz_1 dz_2}{(2\pi i)^2} (-1)^{z_1+z_2} \frac{\Gamma(-z_1) \Gamma(z_1+1)^2 \Gamma(-z_2) \Gamma(z_2+1)^2}{\Gamma(-2\epsilon+z_1+z_2+2)}. \quad (8.13)$$

To proceed, we notice that one of the two integrations evaluates to a ${}_2F_1$, which can be reduced to Γ functions using Gauss' identity,

$$\begin{aligned} \int_{-i\infty}^{+i\infty} \frac{dz_1}{2\pi i} (-1)^{z_1} \frac{\Gamma(-z_1) \Gamma(z_1+1)^2}{\Gamma(-2\epsilon+z_1+z_2+2)} &= \frac{1}{\Gamma(2-2\epsilon+z_2)} {}_2F_1(1, 1; 2-2\epsilon+z_2; 1) \\ &= \frac{\Gamma(-2\epsilon+z_2)}{\Gamma(1-2\epsilon+z_2)^2}. \end{aligned} \quad (8.14)$$

Inserting this result into the two-fold MB integral, we immediately see that the remaining MB integral evaluates to a ${}_3F_2$ function, and we get

$$\mathcal{F}_7(\epsilon) = -\frac{3}{2} \frac{\Gamma(6-6\epsilon)}{\epsilon^5 \Gamma(1-6\epsilon)} {}_3F_2(1, 1, -2\epsilon; 1-2\epsilon, 1-2\epsilon; 1). \quad (8.15)$$

8.4 The master integral $\mathcal{F}_4$

The integral $\mathcal{F}_4$ is defined by

$$\Phi_4^S(\epsilon) \mathcal{F}_4(\epsilon) = \int \frac{d\Phi_4^S}{s_{13}s_{15}s_{34}s_{45}}. \quad (8.16)$$

The integrand only contains two-particle invariants and we can immediately integrate out the energy and angular variables in the usual way. We obtain a one-fold MB representation,

$$\begin{aligned} \mathcal{F}_4(\epsilon) &= \frac{\Gamma(6-6\epsilon)\Gamma(-2\epsilon)}{\epsilon^4 \Gamma(-6\epsilon)\Gamma(-\epsilon)^2} \int_{-i\infty}^{+i\infty} \frac{dz_1}{2\pi i} \Gamma(-z_1) \\ &\times \frac{\Gamma(z_1+1)^2 \Gamma(z_1-2\epsilon) \Gamma(-z_1-\epsilon-1) \Gamma(z_1-\epsilon+1)}{\Gamma(z_1-2\epsilon+1)^2}. \end{aligned} \quad (8.17)$$

Closing the integration contour to the right and summing up residues at $z_1 = n$ and $z_1 = -1-\epsilon+n$, $n \in \mathbb{N}^\times$, we immediately see that $\mathcal{F}_4$ can be expressed as a combination

of hypergeometric functions,

$$\begin{aligned} \mathcal{F}_4(\epsilon) = & -\frac{3\Gamma(6-6\epsilon)\Gamma(1-2\epsilon)}{2\epsilon^4\Gamma(1-6\epsilon)} \\ & \times \left[\frac{3\Gamma(1-2\epsilon)\Gamma(\epsilon+1)}{(1+3\epsilon)\Gamma(1-3\epsilon)} {}_3F_2(-3\epsilon-1, -2\epsilon, -\epsilon; -3\epsilon, -3\epsilon; 1) \right. \\ & \left. + \frac{1}{(1+\epsilon)\Gamma(1-2\epsilon)} {}_4F_3(1, 1, 1-\epsilon, -2\epsilon; 1-2\epsilon, 1-2\epsilon, 2+\epsilon; 1) \right]. \end{aligned} \quad (8.18)$$

8.5 The master integral $\mathcal{F}_6$

The integral $\mathcal{F}_6$ contains two sums in the denominator, which we can replace by products to the price of introducing two MB integrations,

$$\begin{aligned} \Phi_4^S(\epsilon) \mathcal{F}_6(\epsilon) &= \int \frac{d\Phi_4^S}{(s_{13}+s_{14})(s_{14}+s_{15})s_{23}s_{34}} \\ &= \int_{-i\infty}^{+i\infty} \frac{dz_1 dz_2}{(2\pi i)^2} \Gamma(-z_1) \Gamma(z_1+1) \Gamma(-z_2) \Gamma(z_2+1) \int \frac{d\Phi_4^S}{s_{13}^{-z_1} s_{14}^{z_1+z_2+2} s_{15}^{-z_2} s_{23}s_{34}}. \end{aligned} \quad (8.19)$$

We then proceed in the by now familiar way and integrate out the energies and the angles, and we arrive at the following two-fold MB representation for $\mathcal{F}_6$,

$$\begin{aligned} \mathcal{F}_6(\epsilon) &= \frac{\Gamma(6-6\epsilon)}{\epsilon\Gamma(1-6\epsilon)\Gamma(1-\epsilon)^2} \int_{-i\infty}^{+i\infty} \frac{dz_1 dz_2}{(2\pi i)^2} \Gamma(-z_1) \Gamma(z_1+1) \Gamma(-z_2) \Gamma(z_2+1) \\ &\times \frac{\Gamma(-\epsilon+z_1-z_2) \Gamma(z_2-\epsilon) \Gamma(-2\epsilon-z_1+z_2) \Gamma(-\epsilon-z_1+z_2)}{\Gamma(-\epsilon+z_2+1) \Gamma(-2\epsilon-z_1+z_2+1)}. \end{aligned} \quad (8.20)$$

Unlike in the previous cases, we were not able to reduce this integral for generic ϵ to simple hypergeometric functions. We therefore only compute the Laurent expansion of the integral. We proceed in the standard way: we apply the packages **MB** [94], **MBresolve** [95] and **barnesroutines** [96] to resolve singularities in ϵ and to expand the resulting integrals under the integration sign and apply Barnes' lemmas in an automated way. The resulting MB integrals are at most two-fold, and all of them can easily be done by closing the contours to the right and summing up residues in terms of nested harmonic sums defined recursively by [97]

$$S_i(n) = \sum_{k=1}^n \frac{1}{k^i} \quad \text{and} \quad S_{i\bar{j}}(n) = \sum_{k=1}^n \frac{S_{\bar{j}}(k)}{k^i}. \quad (8.21)$$

Note that in the limit $n \rightarrow \infty$ harmonic sums immediately reduce to combinations of multiple zeta values. The result for $\mathcal{F}_6$ reads

$$\begin{aligned} \mathcal{F}_6(\epsilon) &= \frac{10}{\epsilon^5} - \frac{137}{\epsilon^4} + \frac{1}{\epsilon^3} \left(40\zeta_2 + 675 \right) + \frac{1}{\epsilon^2} \left(320\zeta_3 - 548\zeta_2 - 1530 \right) \\ &+ \frac{1}{\epsilon} \left(1500\zeta_4 - 4384\zeta_3 + 2700\zeta_2 + 1620 \right) + 5160\zeta_5 + 320\zeta_2\zeta_3 - 20550\zeta_4 \\ &+ 21600\zeta_3 - 6120\zeta_2 - 648 + \epsilon \left(18340\zeta_6 + 1280\zeta_3^2 - 70692\zeta_5 - 4384\zeta_2\zeta_3 \right. \\ &\left. + 101250\zeta_4 - 48960\zeta_3 + 6480\zeta_2 \right) + \mathcal{O}(\epsilon^2). \end{aligned} \quad (8.22)$$

8.6 The master integral $\mathcal{F}_{10}$

The integrand of $\mathcal{F}_{10}$ involves two sums in the denominator, so we start by introducing two MB representations,

$$\begin{aligned}\Phi_4^S(\epsilon) \mathcal{F}_{10}(\epsilon) &= \int \frac{d\Phi_4^S}{(s_{23} + s_{24})(s_{24} + s_{25})s_{34}s_{45}} \\ &= \int_{-i\infty}^{+i\infty} \frac{dz_1 dz_2}{(2\pi i)^2} \Gamma(-z_1) \Gamma(z_1 + 1) \Gamma(-z_2) \Gamma(z_2 + 1) \int \frac{d\Phi_4^S}{s_{23}^{-z_1} s_{24}^{2+z_1+z_2} s_{25}^{-z_2} s_{34}s_{45}}.\end{aligned}\quad (8.23)$$

Integrating over the angles of the particles three and four yields two hypergeometric functions, and we introduce an MB representation for each of them. Performing the integration over the last angle, we obtain a four-fold MB representation for $\mathcal{F}_{10}$. Two integrations can immediately be performed using Barnes' lemmas, and we obtain

$$\begin{aligned}\mathcal{F}_{10}(\epsilon) &= 6 \frac{\Gamma(6-6\epsilon)}{\Gamma(1-\epsilon)^3 \Gamma(1-6\epsilon)} \int_{-i\infty}^{+i\infty} \frac{dz_2 dz_3}{(2\pi i)^2} \Gamma(-z_2) \Gamma(-z_3) \Gamma(z_2 + 1) \Gamma(z_3 + 1) \\ &\quad \times \frac{\Gamma(-2\epsilon - z_2 - 1) \Gamma(-\epsilon + z_2 + 1) \Gamma(-2\epsilon - z_3 - 1) \Gamma(-\epsilon - z_2 - z_3 - 1) \Gamma(z_3 - \epsilon)}{\Gamma(-2\epsilon - z_2) \Gamma(-2\epsilon - z_3)}.\end{aligned}\quad (8.24)$$

After resolving the singularities in ϵ and expanding under the integration sign, all the twofold integrals can be reduced to onefold integrals using Barnes' lemmas and their corollaries. The remaining onefold integrals are trivial to compute by closing the contour and summing up residues. We find,

$$\begin{aligned}\mathcal{F}_{10}(\epsilon) &= -\frac{120}{\epsilon^4} + \frac{2004}{\epsilon^3} - \frac{14112}{\epsilon^2} + \frac{1}{\epsilon} \left(240\zeta_3 + 60696 \right) + 1980\zeta_4 - 4008\zeta_3 \\ &\quad - 201528 + \epsilon \left(6960\zeta_5 + 1680\zeta_2\zeta_3 - 33066\zeta_4 + 28224\zeta_3 + 612360 \right) \\ &\quad + \epsilon^2 \left(32700\zeta_6 + 6840\zeta_3^2 - 116232\zeta_5 - 28056\zeta_2\zeta_3 + 232848\zeta_4 - 121392\zeta_3 \right. \\ &\quad \left. - 1837080 \right) + \mathcal{O}(\epsilon^3).\end{aligned}\quad (8.25)$$

8.7 The master integral $\mathcal{F}_5$

We start by replacing the sums in the denominator of the integrand of $\mathcal{F}_5$ by three MB integrals,

$$\begin{aligned}\Phi_4^S(\epsilon) \mathcal{F}_5(\epsilon) &= \int \frac{d\Phi_4^S}{(s_{14} + s_{15})s_{23}s_{345}} \\ &= \int_{-i\infty}^{+i\infty} \frac{dz_1 dz_2 dz_3}{(2\pi i)^3} \Gamma(-z_1) \Gamma(-z_2) \Gamma(-z_3) \Gamma(1 + z_1 + z_2) \Gamma(1 + z_3) \\ &\quad \times \int \frac{d\Phi_4^S}{s_{14}^{1+z_1} s_{15}^{-z_1} s_{23}s_{34}^{1+z_1+z_2} s_{45}^{-z_1} s_{35}^{-z_2}}.\end{aligned}\quad (8.26)$$

We insert the energies and angles parametrization for the final-state particles and introduce MB integrations for the angular integrals. Note that the first angular integral necessarily involves three massless propagators and can thus not be done in closed form as a ${}_2F_1$, so

we insert the three-fold MB representation (5.17) for it. We then arrive at a representation for $\mathcal{F}_5$ as a six-fold MB integral convoluted with two angular integrations. It turns out that after performing the change of variables $z_3 \rightarrow z_3 + z_5$ and $z_6 \rightarrow z_6 + z_1 + z_2$, the integrals over z_1 , z_2 and z_5 can be done in closed form using Barnes' first lemma. The remaining two angular integrations can easily be performed in terms of hypergeometric functions, and all but one MB integration can be performed using Barnes' lemmas. We thus arrive at a one-fold MB representation for $\mathcal{F}_5$,

$$\mathcal{F}_5(\epsilon) = -\frac{\Gamma(6-6\epsilon)\Gamma(1-3\epsilon)}{\epsilon\Gamma(2-6\epsilon)\Gamma(2-2\epsilon)\Gamma(1-\epsilon)} \times \int_{-i\infty}^{+i\infty} \frac{dz_1}{2\pi i} \frac{\Gamma(-z_1)\Gamma(z_1+1)^2\Gamma(-z_1-2\epsilon)\Gamma(z_1-\epsilon+1)}{\Gamma(z_1-3\epsilon+2)}. \quad (8.27)$$

One might be tempted to sum up the residues at $z_1 = n$ and $z_1 = -2\epsilon + n$, $n \in \mathbb{N}^\times$, for finite values of ϵ to obtain an expression for $\mathcal{F}_5$ as a combination of two hypergeometric functions at 1 valid to all orders in ϵ . The two hypergeometric functions are however separately divergent (even for finite values of ϵ) and only their sum is finite. We therefore only compute a Laurent series for $\mathcal{F}_5$. Resolving singularities in ϵ and summing up residues in terms of harmonic sums we obtain,

$$\begin{aligned} \mathcal{F}_5(\epsilon) = & -\frac{120}{\epsilon} \zeta_2 - 960 \zeta_3 + 684 \zeta_2 + \epsilon \left(-4620 \zeta_4 + 5472 \zeta_3 - 1188 \zeta_2 \right) \\ & + \epsilon^2 \left(-17160 \zeta_5 - 720 \zeta_2 \zeta_3 + 26334 \zeta_4 - 9504 \zeta_3 + 648 \zeta_2 \right) \\ & + \epsilon^3 \left(-64110 \zeta_6 - 2880 \zeta_3^2 + 97812 \zeta_5 + 4104 \zeta_2 \zeta_3 - 45738 \zeta_4 + 5184 \zeta_3 \right) \\ & + \mathcal{O}(\epsilon^4). \end{aligned} \quad (8.28)$$

8.8 The master integral $\mathcal{F}_8$

The master integral $\mathcal{F}_8$ is defined by

$$\Phi_4^S(\epsilon) \mathcal{F}_8(\epsilon) = \int \frac{d\Phi_4^S}{(s_{13} + s_{15})(s_{23} + s_{24})s_{34}s_{35}}. \quad (8.29)$$

We start by introducing two MB integrations in order to remove the sums in the denominator of the integral $\mathcal{F}_8$. The energies are integrated out in terms of Γ functions, and the angular integrations over the particles four and five are performed using eq. (5.16). At this stage we have three integrations left to do: the two MB integrations and the integral over $\cos \theta_3 = 2y - 1$

$$\begin{aligned} \mathcal{F}_8(\epsilon) = & -\frac{6\Gamma(6-6\epsilon)}{\epsilon\Gamma(1-\epsilon)^4\Gamma(1-6\epsilon)} \int_{-i\infty}^{+i\infty} \frac{dz_1 dz_2}{(2\pi i)^2} \int_0^1 dy y^{z_2-\epsilon} (1-y)^{z_1-\epsilon} \Gamma(-z_1) \\ & \times \Gamma(z_1+1) \Gamma(-z_2) \Gamma(z_2+1) \Gamma(-\epsilon-z_1) \Gamma(-\epsilon-z_2) \Gamma(-2\epsilon+z_1+z_2) \\ & \times {}_2F_1(1, z_1+1; 1-\epsilon; y) {}_2F_1(1, z_2+1; 1-\epsilon; 1-y). \end{aligned} \quad (8.30)$$

In order to proceed, we first apply the identity

$${}_2F_1(a, b; c; x) = (1-x)^{-b} {}_2F_1\left(c-a, b; c; \frac{x}{x-1}\right), \quad (8.31)$$

and then insert an MB representation for each hypergeometric function in the integrand of eq. (8.30). The reason to apply eq. (8.31) before inserting the MB integrations comes from the fact that in this way one of the four MB integrations can be performed using Barnes' first lemma. We then arrive at the following three-fold MB representation for $\mathcal{F}_8$,

$$\begin{aligned} \mathcal{F}_8(\epsilon) = & -\frac{6\epsilon\Gamma(6-6\epsilon)}{\Gamma(1-\epsilon)^4\Gamma(1-6\epsilon)} \int_{-i\infty}^{+i\infty} \frac{dz_2 dz_3 dz_4}{(2\pi i)^3} \Gamma(-z_2) \Gamma(-z_3) \Gamma(-z_4) \\ & \times \Gamma(z_3+1) \Gamma(z_2-2\epsilon) \Gamma(-z_2-z_4) \Gamma(z_2+z_4+1) \Gamma(-\epsilon-z_3) \Gamma(z_3-\epsilon) \\ & \times \frac{\Gamma(-2\epsilon+z_2-z_3) \Gamma(-\epsilon-z_4) \Gamma(z_4-\epsilon)}{\Gamma(-2\epsilon+z_2+1) \Gamma(-2\epsilon-z_3-z_4)}. \end{aligned} \quad (8.32)$$

In the rest of this section we show how we can compute a Laurent expansion for this integral. We proceed in the standard way and resolve singularities in ϵ . At the end of this procedure, we have a collection of MB integrals of dimensionality at most three with integration contours that are straight vertical lines. These integrals can then be safely expanded in ϵ under the integration sign. In the following we discuss the computation of the two and three-fold integrals.

Three-fold MB integrals. There is one three-fold integral contributing to $\mathcal{F}_8$,

$$\begin{aligned} \mathcal{F}_{8,3} = & \int_{-i\infty}^{+i\infty} \frac{dz_2 dz_3 dz_4}{(2\pi i)^3} \Gamma(-z_2) \Gamma(z_2) \Gamma(-z_4)^2 \Gamma(z_4) \Gamma(-z_2-z_4) \Gamma(-z_3-z_4) \\ & \times \frac{\Gamma(1-z_3-z_4) \Gamma(z_2+z_4+1) \Gamma(z_3+z_4)^2 \Gamma(z_2+z_3+z_4)}{\Gamma(z_2+1) \Gamma(z_3)}, \end{aligned} \quad (8.33)$$

where we omit all Γ function prefactors and where the integration contours are straight vertical lines defined by

$$\text{Re}(z_3) = 0.28 \quad \text{and} \quad \text{Re}(z_4) = 0.97 \quad \text{and} \quad \text{Re}(z_5) = -0.36. \quad (8.34)$$

We start by closing the z_3 contour to the right and take residues,

$$\begin{aligned} & \int_{-i\infty}^{+i\infty} \frac{dz_2 dz_4}{(2\pi i)^2} \frac{\Gamma(-z_2) \Gamma(z_2) \Gamma(-z_2-z_4) \Gamma(-z_4)^2 \Gamma(z_4) \Gamma(z_2+z_4+1)}{\Gamma(z_2+1)} \\ & \times \sum_{n=1}^{\infty} \left\{ \frac{\Gamma(n+z_2)}{n \Gamma(n-z_4)} \psi(n+z_2) - \frac{\Gamma(n+z_2)}{n \Gamma(n-z_4)} \psi(n-z_4) - \frac{\Gamma(n+z_2)}{n^2 \Gamma(n-z_4)} \right\}. \end{aligned} \quad (8.35)$$

The first two sums can be performed in terms of Γ functions and their derivatives. We illustrate this on the first term (the second term is similar),

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\Gamma(n+z_2)}{n \Gamma(n-z_4)} \psi(n+z_2) &= \lim_{\eta \rightarrow 0} \frac{\partial}{\partial \eta} \sum_{n=1}^{\infty} \frac{\Gamma(n+z_2+\eta)}{n \Gamma(n-z_4)} \\ &= \lim_{\eta \rightarrow 0} \frac{\partial}{\partial \eta} \left\{ \frac{\Gamma(\eta+z_2)}{\Gamma(-z_4)} [\psi(-z_4) - \psi(-\eta-z_2-z_4)] \right\} \\ &= -\frac{\Gamma(z_2)}{\Gamma(-z_4)} \left[\psi(z_2+1) \psi(-z_2-z_4) - \frac{1}{z_2} \psi(-z_2-z_4) - z_2 \psi(z_2+1) \psi(-z_4) \right. \\ & \quad \left. + \frac{1}{z_2} \psi(-z_4) - \psi'(-z_2-z_4) \right]. \end{aligned} \quad (8.36)$$

In this way, the first two terms can effectively be reduced to the computation of two-fold integrals, and we will therefore not discuss them any further in this section.

The third term can also be summed up in closed form. However, unlike the first two terms, the sum cannot be expressed in terms of Γ functions and their derivatives alone, but the sum evaluates to a ${}_4F_3$ function. We therefore arrive at the following single three-fold MB integral,

$$\mathcal{F}_{8,3} = - \int_{-i\infty}^{+i\infty} \frac{dz_2 dz_4}{(2\pi i)^2} \frac{\Gamma(-z_2) \Gamma(z_2) \Gamma(-z_2 - z_4) \Gamma(-z_4)^2 \Gamma(z_4)}{\Gamma(1 - z_4)} \quad (8.37)$$

$$\times \Gamma(z_2 + z_4 + 1) {}_4F_3(1, 1, 1, z_2 + 1; 2, 2, 1 - z_4; 1) .$$

Although it looks like we have managed to reduce the three-fold integral to a two-fold integral, it is still secretly three-fold, except that we have ‘hidden’ one integration inside the ${}_4F_3$ function. The advantage of this representation is that we can change the representation for the ${}_4F_3$ function in the integrand. More precisely, we perform the change of variables $z_4 \rightarrow -z_4 - z_2$ and chose the contours to be straight vertical lines given by

$$\text{Re}(z_2) = \frac{1}{3} \quad \text{and} \quad \text{Re}(z_4) = \frac{1}{5} . \quad (8.38)$$

We then insert an Euler integral representation for the ${}_4F_3$ function as an integral over a ${}_3F_2$. The ${}_3F_2$ function turns out to be reducible,

$${}_3F_2(1, 1, 1; 2, 2; t) = \frac{\text{Li}_2(t)}{t} , \quad (8.39)$$

and so we finally arrive at

$$\mathcal{F}_{8,3} = - \int_{-i\infty}^{+i\infty} \frac{dz_2 dz_4}{(2\pi i)^2} \int_0^1 dt t^{z_2-1} (1-t)^{z_4-1} \text{Li}_2(t)$$

$$\times \frac{\Gamma(-z_2) \Gamma(z_2) \Gamma(1 - z_4) \Gamma(-z_2 - z_4) \Gamma(z_2 + z_4)^2}{\Gamma(z_2 + 1)} , \quad (8.40)$$

Next, we would like to exchange the MB and the Euler integration and sum up the residues of the poles of the Γ functions in the integrand. However, we are only allowed to do so if the Euler integration does not produce any new poles whose residues need to be taken into account. It is easy to see that in our case the Euler integral converges whenever $\text{Re}(z_2)$ and $\text{Re}(z_4)$ are positive. We can thus close both contours to the right and exchange the Euler and MB integrations and then sum up the residues coming from the Γ functions.

We start by taking residues in z_2 and then the residues in z_4 . There are several cases to be considered separately:

1. The poles at $z_2 = n_2 \in \mathbb{N}^\times$ give rise to poles in z_4 at $n_4 \in \mathbb{N}^\times$. We hence obtain double sums of the form

$$\sum_{n_2, n_4=1}^{\infty} \binom{n_2 + n_4}{n_2} \frac{S_{\tilde{i}_1}(n_2)}{n_2^{j_1}} \frac{S_{\tilde{i}_2}(n_4)}{n_4^{j_2}} \frac{S_{\tilde{i}_3}(n_2 + n_4)}{(n_2 + n_4)^{j_3}} t^{n_2} (1-t)^{n_4} . \quad (8.41)$$

Sums of this type can be performed using **XSummer** [98, 99], and give rise to complicated multiple polylogarithms whose arguments are rational functions of t and $(1-t)$. Using symbols (see appendix D), all these complicated multiple polylogarithms can be reduced to harmonic polylogarithms with indices 0 and 1 in t .

2. Taking the residues at the poles at $z_2 = -z_4 + n_2$, $n_2 \in \mathbb{N}^\times$ gives rise to the expression

$$-\int_0^1 dt \operatorname{Li}_2(t) \sum_{n_2=1}^{\infty} \int_{-i\infty}^{+i\infty} \frac{dz_4}{2\pi i} \frac{z_4^2 n_2! \Gamma(-z_4)^2 \Gamma(z_4)}{n_2^2 (n_2 - z_4)^2 \Gamma(n_2 - z_4)} t^{n_2 - z_4 - 1} (1-t)^{z_4 - 1}. \quad (8.42)$$

Next we want to close the z_4 contour and sum up the corresponding residues. From the previous discussion, we are forced to close the contour to the right, and the summand obviously only has poles at $z_4 = n_4 \in \mathbb{N}^\times$ inside the integration contour. There is however a subtlety, and we cannot just sum up the residues. Indeed, it is easy to see that

- for $n_4 < n_2$, there are double poles. Shifting the summation variable $n_2 \rightarrow n_2 + n_4$, these residues give rise to double sums similar to eq. (8.41) and can again be performed using **XSummer**. We obtain complicated multiple polylogarithms with rational functions of t as argument. Using symbols, they can again be simplified to harmonic polylogarithms with indices 0 and 1 in t .
- for $n_4 = n_2$, there is a simple pole. This gives rise to a simple sum which is trivial to perform in terms of harmonic polylogarithms in t .
- for $n_4 > n_2$, there is a simple pole. Shifting the summation variable $n_4 \rightarrow n_2 + n_4$ this gives rise to a single double sum $S(1-t, 1-1/t)$, with

$$S(x, y) = - \sum_{n_2, n_4=0}^{\infty} \frac{n_2! n_4!}{(n_2 + n_4 + 1)!} \frac{x^{n_2}}{n_2 + 1} \frac{y^{n_4}}{n_4 + 1}. \quad (8.43)$$

This sum is not of the type (8.41), and we therefore need a different way to sum up the series. This procedure will be discussed in the rest of this section.

One way to sum the series $S(x, y)$ is to recognize that it is related to an integral over an Appell F_3 function. More precisely we can write

$$\begin{aligned} S(x, y) &= -\frac{1}{xy} \sum_{n_2, n_4=0}^{\infty} \frac{n_2! n_4!}{(n_2 + n_4 + 1)!} \frac{x^{n_2+1}}{n_2 + 1} \frac{y^{n_4+1}}{n_4 + 1} \\ &= -\frac{1}{xy} \int_0^x d\xi \int_0^y d\chi \sum_{n_2, n_4=0}^{\infty} \frac{(n_2!)^2 (n_4!)^2}{(n_2 + n_4 + 1)!} \frac{\xi^{n_2}}{n_2!} \frac{\chi^{n_4}}{n_4!} \\ &= -\frac{1}{xy} \int_0^x d\xi \int_0^y d\chi F_3(1, 1, 1, 1; 2; \xi, \chi). \end{aligned} \quad (8.44)$$

The particular Appell F_3 function we obtain turns out to be reducible,

$$F_3(\alpha, \gamma - \alpha, \beta, \gamma - \beta; \gamma; \xi, \chi) = (1 - \chi)^{\alpha + \beta - \gamma} {}_2F_1(\alpha, \beta; \gamma; \xi + \chi - \xi\chi), \quad (8.45)$$

and so we obtain a simple two-fold integral representation for S ,

$$\begin{aligned} S(x, y) &= -\frac{1}{xy} \int_0^x d\xi \int_0^y d\chi {}_2F_1(1, 1; 2; \xi + \chi - \xi\chi) \\ &= \frac{1}{xy} \int_0^x d\xi \int_0^y d\chi \frac{\log(1 - \xi) + \log(1 - \chi)}{\xi + \chi - \xi\chi}. \end{aligned} \quad (8.46)$$

This integral is trivial to perform in terms of multiple polylogarithms with rational functions in x and y as arguments (see appendix D). It then follows that $S(1 - t, 1 - 1/t)$ can be expressed in terms of multiple polylogarithms with rational functions in t as arguments. Using symbols, we arrive at the following simple expression,

$$\begin{aligned} S\left(1 - t, 1 - \frac{1}{t}\right) &= \frac{t}{(1 - t)^2} \left[-4\text{Li}_3(t) + 2\text{Li}_2(t) \log t \right. \\ &\quad \left. + \frac{1}{6} \log^3 t + \frac{\pi^2}{3} \log t + 4\zeta_3 \right]. \end{aligned} \quad (8.47)$$

Putting everything together, we arrive at a representation for $\mathcal{F}_{8,3}$ as a one-fold integral over harmonic polylogarithms. This integral is trivial to perform, and we obtain

$$\mathcal{F}_{8,3} = \frac{23\pi^6}{22680} - 2\zeta_3^2. \quad (8.48)$$

Two-fold MB integrals. There is only one two-fold MB integral contributing to $\mathcal{F}_8$ that cannot be reduced to simpler integrals by Barnes' lemmas and their corollaries. This integral reads

$$\begin{aligned} \mathcal{F}_{8,2}(\epsilon) &= \int_{-i\infty}^{+i\infty} \frac{dz_3 dz_4}{(2\pi i)^2} \frac{\Gamma(-z_3)^3 \Gamma(z_3) \Gamma(z_3 + 1) \Gamma(-z_4)^3 \Gamma(z_4) \Gamma(z_4 + 1)}{2\epsilon \Gamma(-z_3 - z_4)} \\ &\quad \times \left[3\epsilon\psi(-z_3) + \epsilon\psi(z_3) - 2\epsilon\psi(-z_3 - z_4) + \epsilon\psi(-z_4) + \epsilon\psi(z_4) - 1 \right], \end{aligned} \quad (8.49)$$

where the integration contours are straight vertical lines defined by

$$\text{Re}(z_3) = -0.64 \quad \text{and} \quad \text{Re}(z_4) = -0.22. \quad (8.50)$$

The integral could in principle be done by closing contours to the left and summing up residues. This leads to double sums of the form (8.41) — thousands of them due to the presence of multiple poles — but without any parametric dependence. We therefore took a different route, which we present in the following.

We start by noting that the *integrand* of eq. (8.49) agrees, up to higher order terms of $\mathcal{O}(\epsilon)$, with the function

$$- \frac{\Gamma(z_3 + 1) \Gamma(-z_4)^2 \Gamma(z_4 + 1) \Gamma(-\epsilon - z_3)^3 \Gamma(z_3 - \epsilon) \Gamma(-\epsilon - z_4) \Gamma(z_4 - \epsilon)}{2\epsilon \Gamma(-2\epsilon - z_3 - z_4)}. \quad (8.51)$$

It would however be wrong to conclude that then necessarily the *integrals*, seen as a Laurent expansion in ϵ , are also equal to the same accuracy, because the Laurent expansion of eq. (8.51) around $\epsilon = 0$ might require to shift the integration contours to avoid pinch

singularities in the limit $\epsilon \rightarrow 0$. It is however easy to see that, for the contours given in eq. (8.50), no new pinch singularity is created for $\epsilon \rightarrow 0$ in eq. (8.51). We therefore consider from now on ϵ to be infinitesimal but finite: it is large enough to separate poles at, e.g., $z_2 = 0$ from $z_2 = \epsilon$, but small enough to ensure that no poles change their nature, i.e., poles that were left (right) of the contour (8.50) in eq. (8.49) remain left (right) of the contour in eq. (8.51). If we chose ϵ in this way, we conclude that

$$\mathcal{F}_{8,2}(\epsilon) = \tilde{\mathcal{F}}_{8,2}(\epsilon) + \mathcal{O}(\epsilon), \quad (8.52)$$

where $\tilde{\mathcal{F}}_{8,2}$ is given by eq. (8.51) integrated over the straight vertical lines defined by eq. (8.50). Our aim will be to find an ϵ expansion for $\tilde{\mathcal{F}}_{8,2}$.

We perform the change of variables $(z_3, z_4) \rightarrow (-1 - z_3, -z_4)$, and we close the z_3 contour to the right and take residues. There are two towers of poles we need to take into account:

$$z_3 = n_3 \in \mathbb{N} \quad \text{and} \quad z_3 = -1 - \epsilon + n_3, \quad n_3 \in \mathbb{N}^\times. \quad (8.53)$$

Two comments are in order:

1. At this stage our assumption that ϵ be infinitesimal but finite is vital, because otherwise the two towers of poles would glue together and thus give rise to double poles.
2. The second tower of poles runs over the set $\{-\epsilon, -\epsilon + 1, \dots\}$. For technical reasons that will become clear below, it is easier to explicitly take into account the residue at $z_3 = -1 - \epsilon$, and to compensate for this by adding it back,

$$\tilde{\mathcal{F}}_{8,2}(\epsilon) = \mathcal{R}_{8,2}(\epsilon) + F_{8,2}(\epsilon), \quad (8.54)$$

where $F_{8,2}$ is the integral obtained from $\tilde{\mathcal{F}}_{8,2}$ by deforming the z_3 contour such that the pole at $z_3 = -1 - \epsilon$ is now to the right of the contour. The residue at $z_3 = -1 - \epsilon$ can be computed in closed form and gives rise to

$$\begin{aligned} \mathcal{R}_{8,2}(\epsilon) = & -\frac{\Gamma(1-\epsilon)^2\Gamma(1+\epsilon)\Gamma(1-2\epsilon)^3}{16\epsilon^5(1+\epsilon)\Gamma(1-3\epsilon)} {}_3F_2(1, 1, 1-\epsilon; 1-3\epsilon, \epsilon+2; 1) \\ & + \frac{\Gamma(1-2\epsilon)^4\Gamma(1-\epsilon)^2\Gamma(1+\epsilon)^2}{8\epsilon^6\Gamma(1-4\epsilon)} - \frac{3\Gamma(1-2\epsilon)^3\Gamma(1-\epsilon)^3\Gamma(1+\epsilon)^2}{16\epsilon^6\Gamma(1-3\epsilon)}. \end{aligned} \quad (8.55)$$

Next we compute $F_{8,2}$ by closing the z_3 contour to the right and summing up residues. The resulting sums can easily be performed in terms of hypergeometric functions, and we get

$$\begin{aligned} F_{8,2}(\epsilon) = & -\int_{-i\infty}^{+i\infty} \frac{dz_4}{2\pi i} \frac{\Gamma(1-z_4)\Gamma(z_4)^2\Gamma(-\epsilon-z_4)\Gamma(z_4-\epsilon)}{2\epsilon\Gamma(z_4-3\epsilon)\Gamma(-2\epsilon+z_4+1)} \\ & \times \left[\Gamma(-\epsilon-1)\Gamma(1-\epsilon)^3\Gamma(z_4-3\epsilon) {}_3F_2(1-\epsilon, 1-\epsilon, 1-\epsilon; \epsilon+2, -2\epsilon+z_4+1; 1) \right. \\ & \left. + \Gamma(-2\epsilon)^3\Gamma(\epsilon+1)\Gamma(-2\epsilon+z_4+1) {}_3F_2(-2\epsilon, -2\epsilon, -2\epsilon; -\epsilon, z_4-3\epsilon; 1) \right]. \end{aligned} \quad (8.56)$$

In order to proceed, we insert an Euler integral representation for each of the ${}_3F_2$ functions. It is then easy to see that the Euler integrals are convergent for $\text{Re}(z_4) > 0$ and ϵ infinitesimal but finite, and so we can exchange the Euler and MB integrations provided that we close the integration contour to the right. The important point is that the ${}_2F_1$ functions appearing inside the Euler integrals are independent of z_4 , and so they can be pulled out of the MB integral. Summing up the residues in z_4 , we then arrive at the following integral representation for $F_{8,2}$.

$$\begin{aligned}
 F_{8,2}(\epsilon) = & \frac{\Gamma(1-\epsilon)}{2\epsilon^3} \int_0^1 dt t^{-\epsilon} (1-t)^{-\epsilon} \\
 & \times \left\{ (1-t)^{-\epsilon} \frac{\Gamma(1-\epsilon)^4 \Gamma(1+\epsilon)}{\epsilon^2 (1+\epsilon) (1-t)} {}_2F_1(1-\epsilon, 1-\epsilon; \epsilon+2; t) \right. \\
 & + t^{-\epsilon-1} (1-t)^{-1-\epsilon} \frac{\Gamma(1-2\epsilon)^2 \Gamma(1-\epsilon) \Gamma(1+\epsilon)^2}{4\epsilon^3} {}_2F_1(-2\epsilon, -2\epsilon; -\epsilon; t) \\
 & - (1-t)^{-1-\epsilon} \frac{\Gamma(1-2\epsilon)^2 \Gamma(1-\epsilon) \Gamma(1+\epsilon)^2}{4\epsilon^3 t} {}_2F_1(-2\epsilon, -2\epsilon; -\epsilon; t) \\
 & - \frac{\Gamma(1-2\epsilon)^2 \Gamma(1+\epsilon)}{4\epsilon(1+\epsilon)t} {}_2F_1(-2\epsilon, -2\epsilon; -\epsilon; t) {}_2F_1(\epsilon+1, \epsilon+1; \epsilon+2; 1-t) \\
 & - t^\epsilon (1-t)^{-\epsilon} \frac{\Gamma(1-\epsilon)^4 \Gamma(1+\epsilon)}{\epsilon^2 (1+\epsilon) (1-t)} {}_2F_1(1-\epsilon, 1-\epsilon; \epsilon+2; t) \\
 & \left. - t^\epsilon \frac{\Gamma(1-\epsilon)^3}{(1+\epsilon)^2} {}_2F_1(1-\epsilon, 1-\epsilon; \epsilon+2; t) {}_2F_1(\epsilon+1, \epsilon+1; \epsilon+2; 1-t) \right\}. \tag{8.57}
 \end{aligned}$$

The terms involving a single ${}_2F_1$ function in the integrand immediately evaluate to ${}_3F_2$ functions, which can be expanded in ϵ using **HypExp**. In addition, the fourth term can be done in closed form as follows: we insert an MB representation for each ${}_2F_1$ and perform the Euler integration as a Beta function. One of the two remaining MB integrals evaluates to a ${}_2F_1$ evaluated at 1, which reduces to Γ functions through Gauss' identity. The remaining one-fold MB integral then immediately evaluates to a ${}_4F_3$ function, which can be expanded in ϵ using **HypExp**. We were not able to find a closed form for the last remaining Euler integral. We therefore insert an Euler integration for each ${}_2F_1$, and we obtain the expression,

$$- \frac{\Gamma(1-\epsilon)^3 \Gamma(1+\epsilon)}{2\epsilon^3 \Gamma(1+2\epsilon)} \mathcal{I}(\epsilon), \tag{8.58}$$

where $\mathcal{I}$ denotes the integral

$$\mathcal{I}(\epsilon) = \int_0^1 dt du dv (1-t)^{-\epsilon} (1-u)^{2\epsilon} u^{-\epsilon} v^\epsilon (1-tu)^{\epsilon-1} (1-(1-t)v)^{-\epsilon-1}. \tag{8.59}$$

It is easy to see that $\mathcal{I}$ is finite as $\epsilon \rightarrow 0$, and so we can expand in ϵ under the integration sign and perform the integration over t , u and v recursively using the techniques of appendix D. This is a trivial exercise that leads to

$$\mathcal{I}(\epsilon) = 2\zeta_3 - \frac{7\pi^4}{180}\epsilon + \left(25\zeta_5 - \frac{\pi^2}{2}\zeta_3\right)\epsilon^2 + \left(-6\zeta_3^2 - \frac{809\pi^6}{22680}\right)\epsilon^3 + \mathcal{O}(\epsilon^4). \tag{8.60}$$

We have thus obtained the ϵ expansion of $F_{8,2}$, and thus of $\mathcal{F}_{8,2}$. We find,

$$\mathcal{F}_{8,2}(\epsilon) = \left(\frac{\pi^2}{6} \zeta_3 - \frac{9}{2} \zeta_5 \right) \frac{1}{\epsilon} - 18\gamma_E \zeta_5 + 4\zeta_3^2 + \frac{2}{3} \gamma_E \pi^2 \zeta_3 - \frac{817\pi^6}{45360} + \mathcal{O}(\epsilon), \quad (8.61)$$

where $\gamma_E = \Gamma'(1)$ denotes the Euler-Mascheroni constant.

The result for $\mathcal{F}_8$. We have now computed all the two and three-fold integrals contributing to $\mathcal{F}_8$. The remaining one-fold integrals are trivial to compute and we obtain

$$\begin{aligned} \mathcal{F}_8(\epsilon) = & -\frac{60}{\epsilon^5} + \frac{822}{\epsilon^4} + \frac{1}{\epsilon^3} (240 \zeta_2 - 4050) + \frac{1}{\epsilon^2} (2400 \zeta_3 - 3288 \zeta_2 + 9180) \\ & + \frac{1}{\epsilon} (13320 \zeta_4 - 32880 \zeta_3 + 16200 \zeta_2 - 9720) + 51840 \zeta_5 + 3360 \zeta_2 \zeta_3 \\ & - 182484 \zeta_4 + 162000 \zeta_3 - 36720 \zeta_2 + 3888 + \epsilon (207600 \zeta_6 + 11760 \zeta_3^2 \\ & - 710208 \zeta_5 - 46032 \zeta_2 \zeta_3 + 899100 \zeta_4 - 367200 \zeta_3 + 38880 \zeta_2) + \mathcal{O}(\epsilon^2). \end{aligned} \quad (8.62)$$

8.9 The master integral $\mathcal{F}_9$

In this section we describe the computation of the most complicated master integral, the integral

$$\Phi_4^S(\epsilon) \mathcal{F}_9(\epsilon) = \int \frac{d\Phi_4^S}{s_{15}(s_{14} + s_{15})s_{23}s_{34}s_{345}}. \quad (8.63)$$

We start in the usual way and derive an MB representation for $\mathcal{F}_9$ by inserting the energies and angles parametrisation and inserting MB integrations for the angular integrals. After applying Barnes' lemmas and their corollaries several times, we arrive at the following MB representation for $\mathcal{F}_9$,

$$\mathcal{F}_9(\epsilon) = \mathcal{F}_{9,1}(\epsilon) + \mathcal{F}_{9,2}(\epsilon), \quad (8.64)$$

where

$$\begin{aligned} \mathcal{F}_{9,1} = & \frac{3(1+6\epsilon)\Gamma(6-6\epsilon)\Gamma(1-4\epsilon)}{2\epsilon(1+4\epsilon)\Gamma(1-6\epsilon)\Gamma(1-\epsilon)^3} \int_{-i\infty}^{+i\infty} \frac{dz_1 dz_2 dz_3 dz_4}{(2\pi i)^4} \Gamma(-z_2) \Gamma(-z_3) \\ & \times \Gamma(1-z_1-z_2) \Gamma(1-z_1-z_3) \Gamma(z_1+z_2+z_3) \Gamma(-z_1-z_2-z_3-z_4) \\ & \times \Gamma(z_1+z_2+z_4+1) \Gamma(z_1+z_3+z_4+1) \Gamma(-z_1-z_2-2\epsilon) \\ & \times \frac{\Gamma(z_1-\epsilon-1) \Gamma(-z_1-z_4-\epsilon-1) \Gamma(z_4-\epsilon+1)}{\Gamma(1-z_2) \Gamma(1-z_3) \Gamma(-z_1-z_2-4\epsilon) \Gamma(z_4-2\epsilon+1)}, \end{aligned} \quad (8.65)$$

$$\begin{aligned} \mathcal{F}_{9,2} = & -\frac{3(1+6\epsilon)\Gamma(6-6\epsilon)\Gamma(1-4\epsilon)}{2\epsilon(1+4\epsilon)\Gamma(1-6\epsilon)\Gamma(1-\epsilon)^3} \int_{-i\infty}^{+i\infty} \frac{dz_1 dz_2 dz_3 dz_4}{(2\pi i)^4} \Gamma(1-z_1) \Gamma(-z_2) \\ & \times \Gamma(-z_3) \Gamma(-z_1-z_3+1) \Gamma(z_1+z_2+z_3) \Gamma(-z_1-z_2-z_3-z_4) \\ & \times \Gamma(z_1+z_2+z_4+1) \Gamma(z_1+z_3+z_4+1) \Gamma(-z_1-z_2-2\epsilon) \\ & \times \frac{\Gamma(z_1-\epsilon-1) \Gamma(-z_1-z_4-\epsilon-1) \Gamma(z_4-\epsilon+1)}{\Gamma(1-z_2) \Gamma(1-z_3) \Gamma(-z_1-4\epsilon) \Gamma(z_4-2\epsilon+1)}. \end{aligned} \quad (8.66)$$

We can resolve singularities for $\mathcal{F}_{9,1}$ and $\mathcal{F}_{9,2}$ and expand in ϵ up to and including $\mathcal{O}(\epsilon)$. The result is a collection of high dimensional MB integrals which, after closing the contour

and taking residues, result in multi-fold harmonic sums. While we were able to perform all the harmonic sums in terms of zeta values for all MB integrals up to $\mathcal{O}(\epsilon^0)$, at $\mathcal{O}(\epsilon)$ new polygamma functions appear in the integrand which make the combinatorics of the sums rather intricate. We therefore chose a different method to evaluate the integral $\mathcal{F}_9$, which we describe in the rest of this section.

We start by noting that the $\mathcal{F}_9$ is finite in $D = 6$ dimensions. This can easily be checked by replacing ϵ by $\epsilon - 1$ in the MB representations (8.65) and (8.66) and resolving singularities. Our goal is to find a parametric integral representation for $\mathcal{F}_9$ in $D = 6 - 2\epsilon$ dimensions and to expand under the integration and perform the parametric integrations recursively. The result in $D = 6 - 2\epsilon$ can then be related to the (divergent) result in $D = 4 - 2\epsilon$ using the dimensional recurrence relation for $\mathcal{F}_9$ of section 6.

It is easy to derive a parametric representation for $\mathcal{F}_9$ using the technique described in appendix C. We find

$$\mathcal{F}_9(D = 6 - 2\epsilon) = \frac{\Gamma(12 - 6\epsilon)\Gamma(3 - 3\epsilon)\Gamma(1 - \epsilon)}{\Gamma(5 - 6\epsilon)\Gamma(2 - \epsilon)^4} \left[\mathcal{I}_{9,1}(\epsilon) + \mathcal{I}_{9,2}(\epsilon) \right], \quad (8.67)$$

with

$$\begin{aligned} \mathcal{I}_{9,1}(\epsilon) &= - \int_0^\infty dt_1 dt_2 \int_0^1 dx_1 dx_2 dx_3 t_1^{2-4\epsilon} (1+t_1)^{\epsilon-1} t_2^{1-2\epsilon} \\ &\quad \times x_1^{-\epsilon} (1-x_1)^{2-4\epsilon} x_2^{1-3\epsilon} (1-x_2)^{-\epsilon} x_3^{-\epsilon} (1+t_2 x_3)^{1-3\epsilon} (1+t_2 x_2 x_3)^\epsilon \\ &\quad \times (t_1 t_2^2 x_1 x_2 x_3 + t_2^2 x_2 x_3 + t_1 t_2 x_1 x_2 + t_1 t_2 x_3 + t_2 x_2 x_3 + t_2 + t_1 + 1)^{3\epsilon-3}, \\ \mathcal{I}_{9,2}(\epsilon) &= \int_0^\infty dt_1 dt_2 \int_0^1 dx_1 dx_2 dx_3 t_1^{2-4\epsilon} (1+t_1)^{\epsilon-1} t_2^{1-2\epsilon} \\ &\quad \times x_1^{1-\epsilon} (1-x_1)^{2-4\epsilon} x_2^{1-3\epsilon} (1-x_2)^{-\epsilon} x_3^{-\epsilon} (1+t_2 x_3)^{1-3\epsilon} (1+t_2 x_2 x_3)^\epsilon \\ &\quad \times (t_1 t_2^2 x_1 x_2 x_3 + t_2^2 x_1 x_2 x_3 + t_2 x_1 + t_1 t_2 x_1 x_2 + t_1 t_2 x_3 + t_2 x_1 x_2 x_3 + t_1 + x_1)^{3\epsilon-3}, \end{aligned} \quad (8.68)$$

$$(8.69)$$

Several comments are in order about the parametric integrals we just defined. First, one can easily check that both $\mathcal{I}_{9,1}$ and $\mathcal{I}_{9,2}$ are individually finite as $\epsilon \rightarrow 0$. Second, at first glance our goal to integrate out the integration variables one-by-one seems rather hopeless due to the appearance of the huge polynomial factor. However, as we will see shortly, there is a sufficient condition that allows one to test whether a parametric integral can be performed in terms of multiple polylogarithms, and this criterion is fulfilled for the integrands of $\mathcal{I}_{9,1}$ and $\mathcal{I}_{9,2}$. We very briefly summarize this criterion in the following, and we refer to ref. [100] or to appendix D for more details. In order to understand the criterion, it is important to first understand multiple polylogarithms and their integration.

Multiple polylogarithms are generalizations of the ordinary logarithm and the *classical* polylogarithms,

$$\ln z = \int_1^z \frac{dt}{t} \quad \text{and} \quad \text{Li}_n(z) = \int_0^z \frac{dt}{t} \text{Li}_{n-1}(t), \quad (8.70)$$

with $\text{Li}_1(z) = -\ln(1-z)$. Multiple polylogarithms are defined recursively via the iterated integral [101, 102]

$$G(a_1, \dots, a_n; z) = \int_0^z \frac{dt}{t - a_1} G(a_2, \dots, a_n; t), \quad (8.71)$$

with $G(z) = 1$ and where $a_i, z \in \mathbb{C}$. In the special case where all the a_i 's are zero, we define, using the obvious vector notation $\vec{a}_n = \underbrace{(a, \dots, a)}_n$,

$$G(\vec{0}_n; z) = \frac{1}{n!} \ln^n z. \quad (8.72)$$

In the special case where $a_i \in \{-1, 0, 1\}$, multiple polylogarithms are related to harmonic polylogarithms,

$$H(a_1, \dots, a_n; z) = (-1)^p G(a_1, \dots, a_n; z), \quad (8.73)$$

where p denotes the number of elements in $(a_1, \dots, a_n)$ equal to $+1$.

Suppose now that we are given some integral over some rational function. We would like to be able to perform the integrations one-by-one in terms of multiple polylogarithms. Unfortunately, such a procedure is not possible for a generic integral. In ref. [100] a sufficient condition was derived that allows one to test whether such a recursive integration is possible. For a detailed description of the criterion we refer to ref. [100], or to appendix D, where (a version of) the criterion is reviewed. In a nutshell, as can easily be seen from the definition of multiple polylogarithms, a sufficient condition to perform the parametric integrations recursively using multiple polylogarithms is that at each integration step we can find an integration variable in which all the denominators appearing in the integrand are linear. More formally, let $S^{(i)}$ be the set of all polynomial factor in the integrand of $\mathcal{I}_{9,i}$. Next, we define the sets $S_{(x,y,\dots)}^{(i)}$, where $x, y, \dots$ are integration variables, as the sets of irreducible non-monomial polynomial factors in the integrand after integration over $(x, y, \dots)$ in this order. For more details how to construct these sets we refer to ref. [100] and to appendix D. Here it suffices to say that the sets $S_{(x,y,\dots)}^{(i)}$ can be constructed solely from the knowledge of the sets $S^{(i)}$. A sufficient criterion for a recursive integration in terms of multiple polylogarithms is that there is an ordering of the integration variables such that the sets $S_{(x,y,\dots)}^{(i)}$ corresponding to this ordering contain at least one integration variable in which all the polynomials of the set are linear. Miraculously, this condition is satisfied for both $\mathcal{I}_{9,1}$ and $\mathcal{I}_{9,2}$. We have

$$S^{(1)} = \{1 + t_1, 1 - x_1, 1 - x_2, 1 + t_2 x_3, 1 + t_2 x_2 x_3, \quad (8.74)$$

$$t_1 t_2^2 x_1 x_2 x_3 + t_2^2 x_2 x_3 + t_1 t_2 x_1 x_2 + t_1 t_2 x_3 + t_2 x_2 x_3 + t_2 + t_1 + 1\},$$

$$S^{(2)} = \{1 + t_1, 1 - x_1, 1 - x_2, 1 + t_2 x_3, 1 + t_2 x_2 x_3, \quad (8.75)$$

$$t_1 t_2^2 x_1 x_2 x_3 + t_2^2 x_1 x_2 x_3 + t_2 x_1 + t_1 t_2 x_1 x_2 + t_1 t_2 x_3 + t_2 x_1 x_2 x_3 + t_1 + x_1\}$$

$$S_{(t_1)}^{(1)} = \{1 - x_1, 1 - x_2, 1 + t_2 x_1 x_2, 1 + t_2 x_3, 1 + t_2 x_2 x_3, \quad (8.76)$$

$$- t_2 x_3 x_2 + t_2 x_1 x_3 x_2 + x_1 x_2 - x_3 x_2 + x_3 - 1\},$$

$$S_{(t_1)}^{(2)} = \{1 - x_1, 1 - x_2, 1 + t_2 x_1 x_2, 1 + t_2 x_3, 1 + t_2 x_2 x_3, \\ t_2 x_1 - t_2 x_2 x_1 + t_2 x_2 x_3 x_1 - t_2 x_3 + x_1 - 1\}, \quad (8.77)$$

$$S_{(t_1, x_1)}^{(1)} = \{1 - x_2, 1 + t_2 x_2, 1 - x_3, 1 + t_2 x_3, 1 + t_2 x_2 x_3, \\ t_2 x_2 x_3 + x_2 x_3 - x_3 + 1\}, \quad (8.78)$$

$$S_{(t_1, x_1)}^{(2)} = \{1 - x_2, 1 + t_2 x_2, 1 - x_3, 1 + t_2 x_3, 1 + t_2 x_2 x_3, \\ -t_2 x_2 + t_2 x_2 x_3 + t_2 + 1\}, \quad (8.79)$$

$$S_{(t_1, x_1, x_2)}^{(1)} = \{1 - x_3, 1 + t_2 x_3, 2t_2 x_3 - t_2 + x_3\}, \quad (8.80)$$

$$S_{(t_1, x_1, x_2)}^{(2)} = \{2 + t_2 - x_3, 1 - x_3, 1 + t_2 x_3\}, \quad (8.81)$$

$$S_{(t_1, x_1, x_2, x_3)}^{(1)} = \{1 + t_2, 1 + 2t_2\}, \quad (8.82)$$

$$S_{(t_1, x_1, x_2, x_3)}^{(2)} = \{1 + t_2, 2 + t_2\}. \quad (8.83)$$

We see that if we perform the integration in the order $(t_1, x_1, x_2, x_3, t_2)$ then at each step all the polynomials are linear in the next integration variable. The actual integration can be carried out in an algorithmic way. The procedure is however rather lengthy, so we do not discuss it here in detail, but we refer to appendix D for a detailed description of the integration algorithm. The result is

$$\begin{aligned} \mathcal{F}_9(D = 6 - 2\epsilon) = & 1663200\zeta_3 - 554400\pi^2 + 3326400 \\ & + 120\epsilon \left(+ 1309\pi^4 - 244203\zeta_3 + 2861\pi^2 + 135294 \right) \\ & - 2\epsilon^2 \left(-25779600\zeta_5 - 970200\pi^2\zeta_3 + 838657\pi^4 - 8149392\zeta_3 - 201756\pi^2 \right. \\ & \left. - 31378284 \right) + \frac{4}{15}\epsilon^3 \left(960575\pi^6 + 180873000\zeta_3^2 - 1978358850\zeta_5 \right. \\ & \left. - 33612075\pi^2\zeta_3 + 56663280\zeta_3 + 3240501\pi^4 + 6836130\pi^2 + 810381510 \right) \\ & + \mathcal{O}(\epsilon^4). \end{aligned} \quad (8.84)$$

Using the dimensional recurrence relations for $\mathcal{F}_9$ derived in section 6 we then finally find the value of $\mathcal{F}_9$ in $D = 4 - 2\epsilon$ dimensions,

$$\begin{aligned} \mathcal{F}_9(\epsilon) = & \frac{160}{\epsilon^5} - \frac{1712}{\epsilon^4} + \frac{1}{\epsilon^3} \left(-120\zeta_2 + 2784 \right) + \frac{1}{\epsilon^2} \left(-120\zeta_3 + 1284\zeta_2 + 31968 \right) \\ & + \frac{1}{\epsilon} \left(2520\zeta_4 + 1284\zeta_3 - 2088\zeta_2 - 216864 \right) + 15720\zeta_5 + 1920\zeta_2\zeta_3 \\ & - 26964\zeta_4 - 2088\zeta_3 - 23976\zeta_2 + 795744 + \epsilon \left(82520\zeta_6 + 9600\zeta_3^2 \right. \\ & \left. - 168204\zeta_5 - 20544\zeta_2\zeta_3 + 43848\zeta_4 - 23976\zeta_3 + 162648\zeta_2 - 2449440 \right) \\ & + \mathcal{O}(\epsilon^2). \end{aligned} \quad (8.85)$$

9 Conclusions and outlook

In this article, we presented a method for the expansion of phase-space integrals in kinematic parameters. We applied our method to perform a threshold expansion of triple-real radiation partonic cross-sections for Higgs production at hadron colliders. Our method reduces the phase-space integrals in the coefficients of the expansion to a small number of *soft* master integrals. These master integrals emerge not only in Higgs production processes but in all processes which are topologically equivalent, such as Drell-Yan production. We have computed the master integrals using a combination of techniques for Mellin-Barnes integration and the integration of real parametric integrals. We presented explicit expressions for the first two coefficients in the threshold expansion of all triple-real radiation partonic cross-sections which are necessary for the inclusive Higgs cross-section at N₃LO, both in terms of *soft* master integrals and as a Laurent series expansion in the dimensional regulator ϵ .

We can apply our method to compute further coefficients in the threshold expansion. This requires the reduction of higher rank cut loop integrals in their powers of propagators and irreducible numerators. It also requires the evaluation of some new master integrals originating from diagrams which did not contribute to the first two terms of the threshold expansion. We believe that it is feasible to compute in this way a sufficient number of terms in the threshold expansion for phenomenology purposes. A second possibility is to apply the reverse-unitarity method to compute the same cross-sections without resorting to a threshold expansion. The work that we have presented here will be particularly important for solving the unexpanded master integrals. With reverse unitarity, one can derive differential equations for the master integrals in arbitrary kinematics. For their solution, we can use the soft limits presented here as boundary conditions. Subleading terms in the expansion can serve as a check of the solutions or as an aid to guess them.

A complete computation of the N₃LO coefficient requires also virtual contributions integrated over the phase space of up to two real partons in the final state. The corresponding one-loop, two-loop and three-loop amplitudes are already known in the literature as a Laurent expansion in ϵ [103–113]. Integrating them over phase-space requires in addition their universal single-real and double-real infrared limits at higher orders in ϵ . A threshold expansion of mixed real and virtual corrections requires an extension of our method. While the threshold expansion for real emission matrix elements can be performed by uniformly rescaling all final-state parton momenta, $p_i \rightarrow \bar{z}p_i$, and expanding in the small parameter $\bar{z}$, for real-virtual matrix-elements the scaling of the loop momentum cannot be determined without further consideration. To determine the correct expansion at threshold we consider the strategy of regions [114, 115]. We applied this method to all 2-loop real-virtual master integrals contributing to Higgs production at NNLO and could reproduce the soft limits of the integrals. In the following we will illustrate this technique on the example of the integral

$$\mathcal{I} = \text{Diagram} = \int d\Phi_2 \frac{1}{(p_3 + p_2)^2} \int \frac{d^D k}{i\pi^{D/2}} \frac{1}{k^2 (k + p_1)^2 (k - p_3)^2 (k + p_{12})^2}. \quad (9.1)$$

We introduce the light-cone coordinates, e.g. for q^μ ,

$$q_\pm = \frac{1}{\sqrt{2}} (q^0 \pm q^1), \quad (9.2)$$

and $q_\perp^\mu$ denotes the remaining $(D-2)$ Euclidian transverse components. Furthermore we choose our reference frame such that

$$\begin{aligned} p_{1,+} &= \frac{1}{\sqrt{2}}, & p_{1,-} &= 0, & p_{1,\perp} &= 0, \\ p_{2,+} &= 0, & p_{2,-} &= \frac{1}{\sqrt{2}}, & p_{2,\perp} &= 0. \end{aligned} \quad (9.3)$$

We make the soft region manifest by redefining the loop momentum and the final state parton momentum in terms of the soft-expansion parameter $\bar{z}$

$$k_- \rightarrow \bar{z} k_-, \quad k_\perp \rightarrow \sqrt{\bar{z}} k_\perp, \quad p_3 \rightarrow \bar{z} p_3. \quad (9.4)$$

After expanding in $\bar{z}$ we find the leading soft integral to be

$$\mathcal{I}^S = \bar{z}^{-1-3\epsilon} \int d\Phi_2^S \frac{1}{\sqrt{2} p_{3+}} \int \frac{d^D k}{i\pi^{D/2}} \frac{1}{k^2 (k^2 + \sqrt{2} k_-) (k^2 - 2k_+ p_{3-}) (\sqrt{2} k_+ + 1)}.$$

All external kinematic scales are factored out of the integrand and the integral can be easily computed using Schwinger parameters. This yields

$$\mathcal{I}^S = -3 \, 2^{2\epsilon-3} \pi^{-1+\epsilon} \bar{z}^{-1-3\epsilon} \frac{\Gamma(1-\epsilon)^2 \Gamma(\epsilon+1)}{\epsilon^3 \Gamma(1-3\epsilon)}, \quad (9.5)$$

in agreement with the result in ref. [71]. In full kinematics other regions, which are suppressed in the soft limit, contribute to the integral.

Finally, the computation of the collinear counterterms for the partonic cross-section has been performed in refs. [116, 117]. Alternatively, this task can easily be achieved numerically following the procedure of ref. [40] and using the results for the partonic cross-sections at NNLO through $\mathcal{O}(\epsilon)$ of refs. [71, 72].

We believe that the prospects for a complete computation of the $N_3\text{LO}$ coefficient for Higgs production and other processes are excellent, and we are looking forward to completing this task in future publications.

Acknowledgments

The authors are grateful to F. Herzog for stimulating discussions. This work was supported by the ERC grant “IterQCD” and the Swiss National Science Foundation project 200021_143781.

A Parametrisation of the $2 \rightarrow H + (N - 2)$ phase space

In this appendix we derive the parametrisation of the $2 \rightarrow H + (N - 2)$ phase space in terms of Lorentz invariant quantities. The usual D -dimensional phase space measure is given in eq. (2.3) and we rescale the momenta as defined in eq. (2.6) and find for $s = 1$,

$$\begin{aligned} d\Phi_{N-1}(p_H, p_3, \dots, p_N; z; D) \\ = \bar{z}^{(D-2)(N-2)} \frac{d^D p_H}{(2\pi)^{D-1}} \delta_+(p_H^2 - M^2) \prod_{i=3}^N \frac{d^D p_i}{(2\pi)^{D-1}} \delta_+(p_i^2) (2\pi)^D \delta^{(D)}(p_{12} - p_H - p_{3\dots N}) , \end{aligned} \quad (\text{A.1})$$

for two incoming partons with momenta p_1 and p_2 producing a Higgs boson with momentum p_H and n additional partons with momenta p_i . We parametrise the final-state momenta as $p_i = (p_{i,0}, p_{i,z}, p_{i,\perp})$, so that $p_{i,\perp}$ is the $(D - 2)$ -dimensional transverse component of the momentum. Then we can rewrite the integration over the momenta of the final-state partons as,

$$d^D p_i \delta_+(p_i^2) = dp_{i,0} dp_{i,z} d^{D-2} p_{i,\perp} \Theta(p_{i,0}) \delta(p_{i,0}^2 - p_{i,z}^2 - p_{i,\perp}^2) , \quad (\text{A.2})$$

Next we choose a frame such that

$$p_1 = \frac{1}{2}(1, 1, 0, \dots, 0) \quad \text{and} \quad p_2 = \frac{1}{2}(1, -1, 0, \dots, 0) . \quad (\text{A.3})$$

In this frame we can write the t -channel invariants ($i \geq 3$) as

$$\begin{aligned} s_{1i} &= (p_1 - p_i)^2 = -2p_1 p_i = -(p_{i,0} - p_{i,z}) , \\ s_{2i} &= (p_2 - p_i)^2 = -2p_2 p_i = -(p_{i,0} + p_{i,z}) , \end{aligned} \quad (\text{A.4})$$

or equivalently

$$p_{i,0} = -\frac{1}{2}(s_{1i} + s_{2i}) \quad \text{and} \quad p_{i,z} = -\frac{1}{2}(s_{2i} - s_{1i}) , \quad (\text{A.5})$$

such that

$$p_{i,\perp}^2 = s_{1i} s_{2i} . \quad (\text{A.6})$$

The s -channel invariants ($i, j \geq 3$) can be written as

$$s_{i,j} = (s_{1i} s_{2j} + s_{1j} s_{2i}) - 2p_{i,\perp} \cdot p_{j,\perp} . \quad (\text{A.7})$$

This yields

$$d^D p_i \delta_+(p_i^2) = \frac{1}{2} d^{D-2} p_{i,\perp} ds_{1i} ds_{2i} \Theta(-s_{1i}) \Theta(-s_{2i}) \delta(s_{1i} s_{2i} - p_{i,\perp}^2) . \quad (\text{A.8})$$

After eliminating the integration over the Higgs momentum using momentum conservation the measure becomes

$$\begin{aligned} d\Phi_{N-1}(D) &= \bar{z}^{(D-2)(N-2)-1} \prod_{i=3}^N \frac{d^{D-2} p_{i,\perp}}{(2\pi)^{D-1}} ds_{1i} ds_{2i} \Theta(-s_{1i}) \Theta(-s_{2i}) \delta(s_{1i} s_{2i} - p_{i,\perp}^2) \\ &\quad \times \frac{\pi}{2^{N-3}} \delta_+ \left((p_{3\dots N} - p_{12})^2 - z p_{3\dots N}^2 \right) . \end{aligned} \quad (\text{A.9})$$

It is well known that rotationally invariant integrals of the form

$$I = \int \prod_i d^D p_i f(p_i \cdot p_j) \quad (\text{A.10})$$

can be rewritten by making the integration over the scalar products manifest:

$$I = \frac{1}{2^N} \int \prod_{i=1}^N \Omega_{D+1-i} \prod_{j=1}^i d(p_i \cdot p_j) [\mathbf{Gram}(p_1, \dots, p_N)]^{\frac{D-N-1}{2}} \Theta[\mathbf{Gram}(p_1, \dots, p_N)] f(p_i \cdot p_j). \quad (\text{A.11})$$

Here $\mathbf{Gram}(p_1, \dots, p_N)$ denotes the Gram determinant of the vectors p_i , which is defined as

$$\mathbf{Gram}(p_1, \dots, p_N) = \det(p_i \cdot p_j)_{1 \leq i, j \leq N}. \quad (\text{A.12})$$

We observe that the measure as well as the matrix element are invariant under rotations in the transverse space. Therefore, we can use this relation to rewrite the $D - 2$ dimensional transverse integration

$$\begin{aligned} \prod_{i=3}^N d^{D-2} p_{i,\perp} &= 2^{2-N} \prod_{i=3}^N \Omega_{D+1-i} \prod_{j=3}^i d(p_{i,\perp} \cdot p_{j,\perp}) \\ &\times [\mathbf{Gram}(p_{3,\perp}, \dots, p_{N,\perp})]^{\frac{D-N-1}{2}} \Theta[\mathbf{Gram}(p_{3,\perp}, \dots, p_{N,\perp})]. \end{aligned} \quad (\text{A.13})$$

Using the definition (A.7) of the s -channel invariants, we can rewrite the integration over the scalar products as

$$\prod_{i=3}^N \prod_{j=3}^i d(p_{i,\perp} \cdot p_{j,\perp}) = (-2)^{\frac{(N-2)(N-3)}{2}} \left(\prod_{i=3}^N dp_{i,\perp}^2 \right) \left(\prod_{3 \leq i < j \leq N} ds_{ij} \right). \quad (\text{A.14})$$

The integration over the norm of the transverse components can be performed using the on-shell δ -functions, so that we find

$$\begin{aligned} \prod_{i=3}^N d^D p_i \delta_+(p_i^2) &= (-1)^{\frac{(N-2)(N-3)}{2}} 2^{6-3N-\frac{(N-2)(N-3)}{2}} \\ &\times \left(\prod_{i=3}^N \Omega_{D+1-i} \right) \left(\prod_{\substack{1 \leq i, j \leq N \\ i \neq j, (i,j) \neq (1,2)}} ds_{ij} \right) G_N(\{s_{ij}\})^{\frac{D-N-1}{2}} \Theta[G_N(\{s_{ij}\})], \end{aligned} \quad (\text{A.15})$$

where we have written the Gram determinant as

$$G_N(\{s_{ij}\}) = \det(s_{1i}s_{2j} + s_{1j}s_{2i} - s_{ij})_{3 \leq i, j \leq N}. \quad (\text{A.16})$$

Thus we finally arrive at the phase space measure

$$d\Phi_{N-1}(D) = \mathcal{N}_{N-2}(D) \bar{z}^{(N-2)(D-2)-1} \left(\prod_{\substack{1 \leq i, j \leq N \\ i \neq j, (i,j) \neq (1,2)}} ds_{ij} \right) \\ \times \delta \left(1 - \sum_{i=3}^N (s_{1i} + s_{2i}) + (1-z) \sum_{i=3}^N \sum_{j=3}^{i-1} s_{ij} \right) G_N(\{s_{ij}\})^{\frac{D-N-1}{2}} \Theta[G_N(\{s_{ij}\})], \quad (\text{A.17})$$

with

$$\mathcal{N}_{N-2}(D) = (-1)^{\frac{(N-2)(N-3)}{2}} 2^{-(N-2)\frac{D}{2}} (2\pi)^{(N-1)-(N-2)D} \prod_{i=3}^N \Omega_{D-i+1}, \quad (\text{A.18})$$

as in eq. (4.4) and

$$\Omega_D = \frac{2\pi^{\frac{D}{2}}}{\Gamma\left(\frac{D}{2}\right)}. \quad (\text{A.19})$$

B Derivation of the phase space volume

In this appendix we derive an expression for the $2 \rightarrow H + (N-2)$ phase space volume that is valid for all orders in z and ϵ . We start by recalling the general phase space factorisation:

$$d\Phi_{k+1}(m^2, s) = \int_{m^2}^s \frac{d\mu^2}{2\pi} d\Phi_{l+1}(\mu^2, s) d\Phi_{k-l+1}(m^2, \mu^2) \quad (\text{B.1})$$

If we assume that we know the phase space volume at $N^k\text{LO}$, i.e. for $2 \rightarrow H + k$,

$$\Phi_{k+1}(m^2, s) = \int d\Phi_{k+1}(m^2, s), \quad (\text{B.2})$$

this relation allows us to rewrite the phase space volume at $N^{k+l}\text{LO}$ as

$$\Phi_{k+l+1}(m^2, s) = \int \int_{m^2}^s \frac{d\mu^2}{2\pi} d\Phi_{l+1}(\mu^2, s) d\Phi_{k+1}(m^2, \mu^2). \quad (\text{B.3})$$

We are specifically interested in rewriting the phase space volume at $N^{k+1}\text{LO}$ as a convolution of the $N^k\text{LO}$ phase space with a two particle phase space. Specialising to $l = 1$, we can use this formula to inductively derive the phase space volume for arbitrary orders. We have,

$$\begin{aligned} \Phi_{k+2}(m^2, s) &= \int \int_{m^2}^s \frac{d\mu^2}{2\pi} d\Phi_2(\mu^2, s) d\Phi_{k+1}(m^2, \mu^2) \\ &= \int_{m^2}^s \frac{d\mu^2}{2\pi} d\Phi_2(\mu^2, s) \Phi_{k+1}(m^2, \mu^2). \end{aligned} \quad (\text{B.4})$$

From eq. (A.17) we obtain an explicit parametrisation of the two particle phase space and we find

$$\begin{aligned}\Phi_{k+2}(m^2, s) &= \frac{1}{4}(2\pi)^{-2+2\epsilon} s^{-\epsilon} \Omega_{2-2\epsilon} \int_{m^2}^s \frac{d\mu^2}{2\pi} \int_0^\infty dx_{1,k+3} dx_{2,k+3} (x_{1,k+3} x_{2,k+3})^{-\epsilon} \\ &\times \delta(1 - x_{1,k+3} - x_{2,k+3}) \left(1 - \frac{\mu^2}{s}\right)^{1-2\epsilon} \Phi_{k+1}(m^2, \mu^2).\end{aligned}\quad (\text{B.5})$$

We can perform the integral over $x_{1,k+3}$ and $x_{2,k+3}$ in terms of beta functions and make the transformation $\mu^2 = sx$, $m^2 = sz$ to obtain

$$\Phi_{k+2}(z, s) = (4\pi)^{-2+\epsilon} s^{1-\epsilon} \frac{\Gamma(1-\epsilon)}{\Gamma(2-2\epsilon)} \int_z^1 dx (1-x)^{1-2\epsilon} \Phi_{k+1}(zs, xs). \quad (\text{B.6})$$

In the following we use eq. (B.6) to prove inductively the following result:

$$\begin{aligned}\Phi_{n+1}(m^2, s) &= \frac{1}{2}(4\pi)^{1-2n+n\epsilon} \frac{\Gamma(1-\epsilon)^n}{\Gamma(2n(1-\epsilon))} s^{n-1-n\epsilon} (1-z)^{2n-1-2n\epsilon} \\ &\times {}_2F_1((n-1)(1-\epsilon), n(1-\epsilon), 2n(1-\epsilon); 1-z),\end{aligned}\quad (\text{B.7})$$

First, eq. (B.7) correctly describes the phase space volume for $n = 1$ and $n = 2$. In order to derive Φ_{n+2} iteratively from Φ_{n+1} we use eq. (B.7) and find

$$\begin{aligned}\Phi_{n+2}(z, s) &= \frac{1}{2}(4\pi)^{1-2(n+1)+(n+1)\epsilon} \frac{\Gamma(1-\epsilon)^{n+1}}{\Gamma(2n(1-\epsilon))\Gamma(2-2\epsilon)} s^{n-(n+1)\epsilon} \\ &\times \int_z^1 dx (1-x)^{1-2\epsilon} x^{n-1-n\epsilon} \left(1 - \frac{z}{x}\right)^{2n-1-2n\epsilon} \\ &\times {}_2F_1\left((n-1)(1-\epsilon), n(1-\epsilon), 2n(1-\epsilon); 1 - \frac{z}{x}\right).\end{aligned}\quad (\text{B.8})$$

To solve the integral, we make the transformation $x = 1 - (1-z)y$ and find

$$\begin{aligned}\Phi_{n+2}(z, s) &= \mathcal{C} \int_0^1 dy y^{1-2\epsilon} (1-y)^{2n-1-2n\epsilon} (1 - (1-z)y)^{-n+n\epsilon} \\ &\times {}_2F_1\left((n-1)(1-\epsilon), n(1-\epsilon), 2n(1-\epsilon); (1-z) \frac{1-y}{1-(1-z)y}\right),\end{aligned}\quad (\text{B.9})$$

where we have factored out

$$\mathcal{C} = \frac{1}{2}(4\pi)^{1-2(n+1)+(n+1)\epsilon} \frac{\Gamma(1-\epsilon)^{n+1}}{\Gamma(2n(1-\epsilon))\Gamma(2-2\epsilon)} s^{n-(n+1)\epsilon} (1-z)^{2(n+1)-1-2(n+1)\epsilon}. \quad (\text{B.10})$$

Next, we introduce a Mellin-Barnes representation for ${}_2F_1$ and exchange the MB integration with the parametric integration. This allows us to perform the integration over y in terms of another ${}_2F_1$. Then we find

$$\begin{aligned}\Phi_{n+2}(z, s) &= \mathcal{C} \frac{\Gamma(2n(1-\epsilon))\Gamma(2-2\epsilon)}{\Gamma((n-1)(1-\epsilon))\Gamma(n(1-\epsilon))} \int_{-i\infty}^{+i\infty} \frac{d\xi}{2\pi i} (z-1)^\xi \Gamma(-\xi) \frac{\Gamma((n-1)(1-\epsilon) + \xi)}{\Gamma(2(n+1)(1-\epsilon) + \xi)} \\ &\times \Gamma(n(1-\epsilon) + \xi) {}_2F_1(n(1-\epsilon) + \xi, 2-2\epsilon, 2(n+1)(1-\epsilon) + \xi; 1-z).\end{aligned}\quad (\text{B.11})$$

Introduce another MB integral for the ${}_2F_1$, we arrive at

$$\begin{aligned} \Phi_{n+2}(z, s) = \mathcal{C} & \frac{\Gamma(2n(1-\epsilon))}{\Gamma((n-1)(1-\epsilon))\Gamma(n(1-\epsilon))} \int_{-i\infty}^{+i\infty} \frac{d\xi d\chi}{(2\pi i)^2} (z-1)^{\xi+\chi} \Gamma(-\xi)\Gamma(-\chi) \\ & \times \frac{\Gamma((n-1)(1-\epsilon)+\xi)\Gamma(n(1-\epsilon)+\xi+\chi)\Gamma(2-2\epsilon+\chi)}{\Gamma(2(n+1)(1-\epsilon)+\xi+\chi)}. \end{aligned} \quad (\text{B.12})$$

In the next step we perform the change of variables $\xi \rightarrow \xi + \chi$, such that

$$\begin{aligned} \Phi_{n+2}(z, s) = \mathcal{C} & \frac{\Gamma(2n(1-\epsilon))}{\Gamma((n-1)(1-\epsilon))\Gamma(n(1-\epsilon))} \int_{-i\infty}^{+i\infty} \frac{d\xi d\chi}{(2\pi i)^2} (z-1)^\xi \Gamma(\chi-\xi)\Gamma(-\xi) \\ & \times \frac{\Gamma((n-1)(1-\epsilon)+\xi-\chi)\Gamma(n(1-\epsilon)+\xi)\Gamma(2-2\epsilon+\xi)}{\Gamma(2(n+1)(1-\epsilon)+\xi)}. \end{aligned} \quad (\text{B.13})$$

The integral over χ can now be performed using Barnes' first lemma, and we find

$$\begin{aligned} \Phi_{n+2}(z, s) = \mathcal{C} & \frac{\Gamma(2n(1-\epsilon))}{\Gamma((n-1)(1-\epsilon))\Gamma(n(1-\epsilon))} \int_{-i\infty}^{+i\infty} \frac{d\xi}{2\pi i} \Gamma(-\xi) (z-1)^\xi \\ & \times \frac{\Gamma(n(1-\epsilon)+\xi)\Gamma((n+1)(1-\epsilon)+\xi)}{\Gamma(2(n+1)(1-\epsilon)+\xi)}. \end{aligned} \quad (\text{B.14})$$

The integral over ξ is just the MB representation of a ${}_2F_1$ so that we finally find

$$\begin{aligned} \Phi_{n+2}(z, s) = \frac{1}{2} (4\pi)^{1-2(n+1)+(n+1)\epsilon} s^{n-(n+1)\epsilon} (1-z)^{2(n+1)-1-2(n+1)\epsilon} & \frac{\Gamma(1-\epsilon)^{n+1}}{\Gamma(2(n+1)(1-\epsilon))} \\ & \times {}_2F_1(n(1-\epsilon), (n+1)(1-\epsilon), 2(n+1)(1-\epsilon); 1-z). \end{aligned} \quad (\text{B.15})$$

We have therefore inductively shown the validity of (B.7) for all n .

C From Mellin-Barnes integrals to parametric integrals

In this appendix we describe how one can derive an Euler-type integral from a Mellin-Barnes integral with a balanced integrand. Roughly speaking, a MB integral is said to be *balanced* if for each integration variable z_i the number of Γ functions of the form $\Gamma(\dots - z_i)$ is equal to the number of Γ functions of the form $\Gamma(\dots + z_i)$. More precisely, the integral

$$\int_{-i\infty}^{+i\infty} \frac{dz_i}{2\pi i} \prod_{k_1=1}^{n_+} \Gamma(a_{k_1} + z_i)^{\alpha_{k_1}} \prod_{k_2=1}^{n_-} \Gamma(b_{k_2} - z_i)^{\beta_{k_2}}, \quad \alpha_{k_i}, \beta_{k_i} \in \mathbb{Z}, \quad (\text{C.1})$$

is said to be *balanced* if $\sum_{k_1}^{n_+} \alpha_{k_1} = \sum_{k_2}^{n_-} \beta_{k_2}$. We assume in the following that the contours are straight vertical lines such that the real parts of the arguments of all the Γ functions are positive.⁶ In that case we can always derive an Euler-type integral representation for

⁶Note that in dimensional regularisation we might need to require ϵ to be finite for such a contour to exist.

the MB integral. We start by noting that if an integral is balanced, then we can always express its integrand as a product of Beta functions,

$$B(x, y) = \int_0^\infty dt t^{x-1} (1+t)^{-x-y}. \quad (\text{C.2})$$

The integral (C.2) is convergent whenever $\text{Re}(x), \text{Re}(y) > 0$. It is easy to convince oneself that this condition is satisfied whenever the real parts of all arguments of the Γ functions are positive in the original MB integral. We can therefore replace each Beta function by its integral representation (C.2) in the *integrand* of the MB integral and, because all the integral are convergent, we can exchange the MB integrations and the integrations coming from the Beta functions. This leaves us with an integral of the form

$$\int_0^\infty \left(\prod_{n=1}^N dt_i \right) R_0(\mathbf{t}) R_\epsilon(\mathbf{t})^\epsilon \int_{-i\infty}^{+i\infty} \prod_{m=1}^M \frac{dz_i}{2\pi i} R_i(\mathbf{t})^{z_i}, \quad (\text{C.3})$$

where $\mathbf{t} = (t_1, \dots, t_N)$ and the R_k are ratios of products of the t_i and $1+t_i$. Next, we would like to perform the MB integrations. This can be done using the formula

$$\int_{z_0-i\infty}^{z_0+i\infty} \frac{dz}{2\pi i} a^z = \delta(1-a), \quad a > 0. \quad (\text{C.4})$$

Indeed, parametrizing the contour as $z = z_0 + it$, we obtain

$$\int_{z_0-i\infty}^{z_0+i\infty} \frac{dz}{2\pi i} a^z = a^{z_0} \int_{-\infty}^{+\infty} \frac{dt}{2\pi} e^{it \ln a} = a^{z_0} \delta(\ln a) = \delta(1-a). \quad (\text{C.5})$$

Equation (C.3) can thus be written in the form

$$\int_0^\infty \left(\prod_{n=1}^N dt_i \right) R_0(\mathbf{t}) R_\epsilon(\mathbf{t})^\epsilon \prod_{m=1}^M \delta(1 - R_i(\mathbf{t})). \quad (\text{C.6})$$

We can solve the δ constraints, and the result is the desired parametric integral.

In the following we discuss two very simple examples that illustrate the above procedure. We start by discussing Barnes' first lemma, i.e., we consider the integral

$$\mathcal{I} = \int_{-i\infty}^{+i\infty} \frac{dz}{2\pi i} \Gamma(a+z) \Gamma(b+z) \Gamma(c-z) \Gamma(d-z). \quad (\text{C.7})$$

We assume that the integration contour and a, b, c and d are such that the real parts of all Γ functions are positive. We rewrite the integrand in terms of Beta functions,

$$\begin{aligned} \mathcal{I} &= \Gamma(a+c) \Gamma(b+d) \int_{-i\infty}^{+i\infty} \frac{dz}{2\pi i} B(a+z, c-z) B(b+z, d-z) \\ &= \Gamma(a+c) \Gamma(b+d) \int_{-i\infty}^{+i\infty} \frac{dz}{2\pi i} \int_0^\infty dt_1 dt_2 t_1^{a+z-1} (1+t_1)^{-a-c} t_2^{b+z-1} (1+t_2)^{-b-d} \\ &= \Gamma(a+c) \Gamma(b+d) \int_0^\infty dt_1 dt_2 t_1^{a-1} (1+t_1)^{-a-c} t_2^{b-1} (1+t_2)^{-b-d} \delta(1-t_1 t_2), \end{aligned} \quad (\text{C.8})$$

where the last step follows from eq. (C.4). Solving the δ -function constraint leads to a one-fold integral that can immediately be recognized as a Beta function, and we recover the usual form of Barnes' first lemma,

$$\begin{aligned}\mathcal{I} &= \Gamma(a+c)\Gamma(b+d) \int_0^\infty dt_1 t_1^{a+d-1} (1+t_1)^{-a-b-c-d} \\ &= \frac{\Gamma(a+c)\Gamma(a+d)\Gamma(b+c)\Gamma(b+d)}{\Gamma(a+b+c+d)}.\end{aligned}\tag{C.9}$$

The second example we are going to discuss is Gauss' hypergeometric function. We consider the integral

$$\mathcal{J} = \int_{-i\infty}^{+i\infty} \frac{dz}{2\pi i} \Gamma(-z) \frac{\Gamma(a+z)\Gamma(b+z)}{\Gamma(c+z)} x^z.\tag{C.10}$$

We again assume that all conditions for convergence are satisfied. Rewriting the integrand in terms of Beta functions, we obtain

$$\begin{aligned}\mathcal{J} &= \frac{\Gamma(b)}{\Gamma(c-a)} \int_{-i\infty}^{+i\infty} \frac{dz}{2\pi i} B(b+z, -z) B(a+z, c-a) x^z \\ &= \frac{\Gamma(b)}{\Gamma(c-a)} \int_0^\infty dt_1 dt_2 t_1^{a-1} (1+t_1)^{-c} t_2^{b-1} (1+t_2)^{-b} \delta\left(1 - \frac{x t_1 t_2}{1+t_1}\right).\end{aligned}\tag{C.11}$$

Solving the δ -function constraint with respect to t_2 and performing the change of variables $t_1 \rightarrow \xi/(1-\xi)$, we immediately arrive at the usual integral representation for the ${}_2F_1$ function,

$$\begin{aligned}\mathcal{J} &= \frac{\Gamma(b)}{\Gamma(c-a)} \int_0^\infty dt_1 t_1^{a-1} (1+t_1)^{b-c} (1+t_1+x t_1)^{-b} \\ &= \frac{\Gamma(b)}{\Gamma(c-a)} \int_0^1 d\xi \xi^{a-1} (1-\xi)^{c-a-1} (1+x\xi)^{-b} \\ &= \frac{\Gamma(a)\Gamma(b)}{\Gamma(c)} {}_2F_1(a, b; c; -x).\end{aligned}\tag{C.12}$$

D Symbolic integration of parametric integrals

In this section we describe an algorithmic approach to compute certain classes of parametric integrals. We stress that this approach is not genuinely new, but has already been successfully applied, in some variant or another, to the computation of Feynman integrals, e.g., [100, 118–122]. More precisely, consider an integral of the form

$$\mathcal{I}(\{y_j\}; \epsilon) = \int_0^\infty \left(\prod_{i=1}^n dx_i \right) \prod_{k=1}^m P_k(\{x_i\}; \{y_j\})^{a_k + \epsilon b_k},\tag{D.1}$$

where a_k and b_k are integers and the $P_k(\{x_i\}; \{y_j\})$ are polynomials with integer coefficients, which we assume irreducible over $\mathbb{Z}$, i.e., they cannot be factorized into a product of non-constant polynomials of lower degree. We assume that the integration range is $[0, \infty]$ for

each integration variable x_i . While many statements remain true for generic integration boundaries, in certain cases the algorithm we are going to describe breaks down, due to the appearance of boundary contributions. We can however always map a generic integration region $x_i \in [a, b]$ to $y_i \in [0, \infty]$ by the change of variable

$$y_i = \frac{x_i - b}{x_i - a}. \quad (\text{D.2})$$

We furthermore assume that the integral is convergent for $\epsilon = 0$, and so we can expand in ϵ under the integration sign. Note that in the simplest divergent cases where the singularities in the integrand factorize we can reduce the problem to a convergent integral by subtracting the divergencies.

Our goal is to compute the coefficients of the Taylor expansion of $\mathcal{I}(\{y_j\}; \epsilon)$ by integrating out the integration variables recursively one-by-one in terms of multiple polylogarithms, eq. (8.71). Obviously, not every integral can be performed in this way. In the rest of this appendix we describe a sufficient condition, first obtained in ref. [100], to be satisfied by the integrand of $\mathcal{I}(\{y_j\}; \epsilon)$ so that it can be integrated in terms of multiple polylogarithms. We then review an algorithm for integrating these classes of integrals, and illustrate the procedure explicitly on a simple example.

D.1 Denominator reduction

In this section we review (a variant of) the sufficient condition of ref. [100] to determine whether an integral can be performed recursively in terms of multiple polylogarithms. The main idea is that we need to determine an ordering of the integration variables such that at each step during the integration all the denominators are linear in the next integration variable. We start by defining the set S of all the polynomials that are not monomials and that appear inside the integrand of eq. (D.1),

$$S = \left\{ P_k(\{x_i\}; \{y_j\}) \right\}. \quad (\text{D.3})$$

To start the integration, we have to assume that there is one integration variable, say x_a such that all the element of S are linear in x_a . In that case we may write

$$P_k(\{x_i\}; \{y_j\}) = Q_k^a(\{x_i\}; \{y_j\}) x_a + R_k^a(\{x_i\}; \{y_j\}), \quad (\text{D.4})$$

where $Q_k^a(\{x_i\}; \{y_j\})$ and $R_k^a(\{x_i\}; \{y_j\})$ are polynomials that are independent of x_a . Note that, after expansion in ϵ , the integrand may also contain logarithms of the $P_k(\{x_i\}; \{y_j\})$, which can be rewritten in terms of multiple polylogarithms,

$$\begin{aligned} \log P_k &= \log \left(Q_k^a x_a + R_k^a \right) = \log R_k^a + \log \left(1 + \frac{Q_k^a}{R_k^a} x_a \right) \\ &= \log R_k^a + G \left(-\frac{R_k^a}{Q_k^a}; x_a \right), \end{aligned} \quad (\text{D.5})$$

where for clarity we suppressed the arguments of the polynomials. Furthermore, we can use the shuffle algebra of multiple polylogarithms to replace every product of multiple

polylogarithms by a sum,

$$G(a_1, \dots, a_{n_1}; z) G(a_{n_1+1}, \dots, a_{n_1+n_2}; z) = \sum_{\sigma \in \Sigma(n_1, n_2)} G(a_{\sigma(1)}, \dots, a_{\sigma(n_1+n_2)}; z), \quad (\text{D.6})$$

where $\Sigma(n_1, n_2)$ denotes the set of all shuffles of $n_1 + n_2$ elements, i.e., the subset of the symmetric group $S_{n_1+n_2}$ defined by

$$\begin{aligned} & \Sigma(n_1, n_2) \\ &= \{ \sigma \in S_{n_1+n_2} \mid \sigma^{-1}(1) < \dots < \sigma^{-1}(n_1) \text{ and } \sigma^{-1}(n_1+1) < \dots < \sigma^{-1}(n_1+n_2) \}. \end{aligned} \quad (\text{D.7})$$

Thus we can assume without loss generality that the integration over x_a takes the form

$$\int_0^\infty \frac{dx_a}{(Q_1^a x_a + R_1^a)^{-a_1} \dots (Q_m^a x_a + R_m^a)^{-a_m}} G(\vec{a}; x_a). \quad (\text{D.8})$$

Partial fractioning the factors in the denominator, e.g.,

$$\frac{1}{(Q_k^a x_a + R_k^a)(Q_l^a x_a + R_l^a)} = \frac{1}{Q_k^a R_l^a - Q_l^a R_k^a} \left(\frac{1}{x_a + R_k^a/Q_k^a} - \frac{1}{x_a + R_l^a/Q_l^a} \right), \quad (\text{D.9})$$

we obtain a sum of integrals that can be reduced to the recursive definition of multiple polylogarithms, eq. (8.71). We can thus easily compute a primitive with respect to x_a , and then take the limits $x_a \rightarrow 0$ and $x_a \rightarrow \infty$. We will address the issue of limits in the next subsection.

We would like to iterate this procedure and integrate over the next variable. This is however only possible if inside the new integrand we can still find an integration variable in which all polynomials are linear. The polynomials appearing inside the integrand are Q_k^a and R_k^a , which have been introduced through eq. (D.5) and (D.9), as well as the combinations $Q_k^a R_l^a - Q_l^a R_k^a$ introduced by partial fractioning. This last polynomial however need not necessarily be linear, even if the Q_k^a and R_k^a are. In order to proceed, it is therefore mandatory that all the $Q_k^a R_l^a - Q_l^a R_k^a$ factor into polynomials that are linear in a certain variable.

In ref. [100] a criterion was given that allows one to determine a priori whether the above procedure terminates, i.e., whether there is an ordering of the integration variables such that all the denominators stay linear at each integration step. We start by defining the set $S_{(x_a)}$ as the set of irreducible factors that appear inside the polynomials Q_k^a , R_k^a and $Q_k^a R_l^a - Q_l^a R_k^a$. Then, if we can find an integration variable x_b such that all the elements of $S_{(x_a)}$ are linear in x_b , we can restart the above procedure and integrate over x_b . If we iterate this procedure and are able to construct a sequence of sets of polynomials

$$S_{(x_a)}, S_{(x_a, x_b)}, S_{(x_a, x_b, x_c)}, \dots \quad (\text{D.10})$$

such that in each set all the polynomials are linear in at least one integration variable, then we have found an ordering of the integration variables such that we can recursively integrate out all the integration variables in terms of multiple polylogarithms. We stress that this condition is sufficient, but not necessary: even if we fail to find a suitable sequence (D.10),

the integral might still be expressible in terms of multiple polylogarithms (e.g., after a suitable change of variables). In addition, note that for the last integration step it is not necessary for the polynomials to be linear: we can then factor the polynomials into linear factors, whose roots involve algebraic expressions of the parameters $\{y_j\}$.

D.2 Symbolic integration

We have now a criterion to determine if a given integral of the type (D.1) can be integrated recursively in terms of multiple polylogarithms, and we have already explained how to perform the first integration step. We assume from now on that we have found a sequence (D.10) and the associated ordering of the integration variables, and that we have performed the integration over x_a as described in the previous subsection.

In order to continue and perform the next integration, we need to address two issues

1. We need to take the limits $x_a \rightarrow 0$ and $x_a \rightarrow \infty$ of the primitive with respect to x_a .
2. If we want to integrate next over x_b , we have to face the problem that x_b might be ‘hidden’ inside the arguments of multiple polylogarithms of the form

$$G\left(\dots, -\frac{R_k^a}{Q_k^a}, \dots\right), \quad (\text{D.11})$$

i.e., x_b appears inside the polynomials Q_k^a and R_k^a . In order to compute a primitive with respect to x_b , we need to rewrite all multiple polylogarithms of the form (D.11) as $G(\vec{a}; x_b)$, where $\vec{a}$ is independent of x_b .

The limit $x_a \rightarrow 0$ can easily be taken by using the fact that

$$\lim_{x_a \rightarrow 0} G(\vec{a}_1; x_a) = 0, \quad \vec{a}_1 \neq \vec{0}. \quad (\text{D.12})$$

While eq. (D.12) is sufficient to compute the value at $x_a = 0$ in many circumstances, it can happen that the primitive has spurious poles and / or logarithmic singularities at $x_a = 0$. In the presence of poles we need to expand the multiple polylogarithms around $x_a = 0$. This task can easily be achieved using the series representation of multiple polylogarithms. Indeed, multiple polylogarithms can equally well be represented as multiple nested sums,

$$\text{Li}_{m_k, \dots, m_1}(z_k, \dots, z_1) = \sum_{n_1 > \dots > n_k > 0} \frac{z_1^{n_1}}{n_1^{m_1}} \cdots \frac{z_k^{n_k}}{n_k^{m_k}} = \sum_{n=1}^{\infty} \frac{z_1^n}{n^{m_1}} Z_{m_2, \dots, m_k}(n-1; z_2, \dots, z_k), \quad (\text{D.13})$$

where $Z_{m_2, \dots, m_k}(n-1; z_2, \dots, z_k)$ denote the Z -sums defined in ref. [98]. The integral representation and the series representation are related by

$$G(\underbrace{0, \dots, 0}_{m_1-1}, a_1, \dots, \underbrace{0, \dots, 0}_{m_k-1}, a_k; x) = (-1)^k \text{Li}_{m_k, \dots, m_1}\left(\frac{a_{k-1}}{a_k}, \dots, \frac{x}{a_1}\right), \quad a_i \neq 0. \quad (\text{D.14})$$

If $a_k = 0$, we can use the shuffle algebra to extract pure logarithms in x_a , e.g., if $b \neq 0$,

$$G(b, 0; x_a) = G(0; x_a)G(b; x_a) - G(0, b; x_a). \quad (\text{D.15})$$

We can thus easily obtain the expansion around $x_a = 0$ in terms of Z -sums to arbitrary order.

Next, we discuss how to take the limit $x_a \rightarrow \infty$. If we define $\bar{x}_a = 1/x_a$, then the problem can be reduced to taking the limit $\bar{x}_a \rightarrow 0$,

$$\lim_{x_a \rightarrow \infty} G(\vec{a}; x_a) = \lim_{\bar{x}_a \rightarrow 0} G\left(\vec{a}; \frac{1}{\bar{x}_a}\right). \quad (\text{D.16})$$

We would like to use the inversion relations for the multiple polylogarithms in the right-hand side to express them as linear combinations of multiple polylogarithms of the form $G(\dots; \bar{x}_a)$, for which the limit $\bar{x}_a \rightarrow 0$ is trivial. Unfortunately, the corresponding functional equations are often unknown. Note at this point that the problem of finding the inversion relations is formally equivalent to the second problem we have not yet addressed, namely how to bring the primitive with respect to x_a into a form where x_b only enters through multiple polylogarithms of the form $G(\vec{a}; x_b)$: in both cases we are looking for functional equations that bring a certain multiple polylogarithm into a ‘canonical form’ where a certain variable only appears as the explicit upper integration boundary of the multiple polylogarithms. In the rest of this section we argue that, under certain conditions which we will later show to be equivalent to the criterion derived in the previous subsection, plus the hypothesis that the integration ranges are $[0, \infty]$, we can always bring multiple polylogarithms into the ‘canonical form’

$$\sum_i c_i G(\vec{a}_i; x), \quad (\text{D.17})$$

for some variable x such that a_i is independent of x and the coefficients c_i involve only multiple polylogarithms that are independent of x . Note that this result is similar to the result obtained in ref. [100]. In the following we give a constructive algorithm that allows us to derive the canonical form (D.17).

In order to achieve a rewriting of our multiple polylogarithms in canonical form, we need to derive the corresponding functional equations. The natural language to discuss functional equations among multiple polylogarithms are symbols [123–127] and the Hopf algebra of multiple polylogarithms [102]. We start by giving a concise review of symbols.

One possible way to define the symbol of a multiple polylogarithm is to consider its total differential [102],

$$dG(a_{n-1}, \dots, a_1; a_n) = \sum_{i=1}^{n-1} G(a_{n-1}, \dots, a_{i-1}, a_{i+1}, \dots, a_1; a_n) d \ln \left(\frac{a_i - a_{i+1}}{a_i - a_{i-1}} \right), \quad (\text{D.18})$$

and to define the symbol recursively by [126]

$$\mathcal{S}(G(a_{n-1}, \dots, a_1; a_n)) = \sum_{i=1}^{n-1} \mathcal{S}(G(a_{n-1}, \dots, a_{i-1}, a_{i+1}, \dots, a_1; a_n)) \otimes \left(\frac{a_i - a_{i+1}}{a_i - a_{i-1}} \right). \quad (\text{D.19})$$

As an example, the symbols of the classical polylogarithms and the ordinary logarithms are given by

$$\mathcal{S}(\text{Li}_n(z)) = -(1-z) \otimes \underbrace{z \otimes \dots \otimes z}_{(n-1) \text{ times}} \text{ and } \mathcal{S}\left(\frac{1}{n!} \ln^n z\right) = \underbrace{z \otimes \dots \otimes z}_{n \text{ times}}. \quad (\text{D.20})$$

In addition the symbol satisfies the following identities,

$$\begin{aligned} \dots \otimes (a \cdot b) \otimes \dots &= \dots \otimes a \otimes \dots + \dots \otimes b \otimes \dots, \\ \dots \otimes (\pm 1) \otimes \dots &= 0, \\ \mathcal{S}\left(G(\vec{a}; x) G(\vec{b}; y)\right) &= \mathcal{S}(G(\vec{a}; x)) \text{ III } \mathcal{S}(G(\vec{b}; y)), \end{aligned} \quad (\text{D.21})$$

where III denotes the shuffle product on tensors.

We can make the following observation about the symbol of a multiple polylogarithm: if the a_i are independent of x , then the symbol of $G(a_1, \dots, a_n; x)$ contains exactly one term that contains x in all its entries, and this term can be chosen to be of the form

$$\mathcal{S}(G(a_1, \dots, a_n; x)) = (a_n - x) \otimes \dots \otimes (a_1 - x) + \dots \quad (\text{D.22})$$

In order to proof this statement, we focus on the term in the total differential of $G(a_1, \dots, a_n; x)$ proportional to dx ,

$$\begin{aligned} dG(a_1, \dots, a_n; x) &= G(a_2, \dots, a_n; x) \frac{dx}{x - a_1} + \dots \\ &= G(a_2, \dots, a_n; x) d \log(a_1 - x) + \dots, \end{aligned} \quad (\text{D.23})$$

where the last step follows from

$$d \log(a_1 - x) = \frac{dx - da_1}{x - a_1}, \quad (\text{D.24})$$

and where the dots indicate terms in the total differential that are independent of dx . The statement then follows recursively. Note that this statement is independent of whether the a_i are zero or not.

Assume now that we are given an integrable symbol T (which will correspond later to the symbol of the multiple polylogarithm we want to bring into canonical form) which is of uniform weight w and has rational coefficients. If T does not satisfy this last condition, we deal separately with the contributions of different weight and / or different rational function prefactor. Let us suppose that the entries are drawn from a set S . Without loss of generality, we may assume S to consist of irreducible polynomials in some variables x_i , $1 \leq i \leq n$, and for simplicity we assume for now that the polynomials are linear in *all* the x_i (we will see in the next subsection that the correct restriction is that S satisfies the reduction criterion of the previous subsection). Furthermore, we assume that we have fixed an ordering on the variables, which we will take in the following as $(x_1, \dots, x_n)$. For each variable x_i , we define a linear map ϕ_{x_i} which acts on elementary tensors s by

$$\phi_{x_i}(s) = \begin{cases} G\left(-\frac{b_1}{a_1}, \dots, -\frac{b_w}{a_w}; x_i\right), & \text{if } s = (a_w x_i + b_w) \otimes \dots \otimes (a_1 x_i + b_1), \\ 0, & \text{otherwise.} \end{cases} \quad (\text{D.25})$$

Morally speaking, the map ϕ_{x_i} assigns to T the combination of multiple polylogarithms which will give the same terms that have x_i in all entries through eq. (D.22).

Using the maps ϕ_{x_i} , we can now formulate an algorithm that assigns to T a multiple polylogarithm in the canonical form associated to the ordering of the variables $(x_1, \dots, x_n)$. We start by defining a new symbol by subtracting off the contribution from $\phi_{x_1}(T)$,

$$T_1 = T - \mathcal{S}[\phi_{x_1}(T)]. \quad (\text{D.26})$$

By construction, each term in the symbol T_1 has at least one entry that is independent of x_1 . Next, concentrate on the terms of the form

$$\sum_{(i_1, \dots, i_w)} c_{i_1, \dots, i_w} b_{i_1} \otimes (a_{i_2} x_1 + b_{i_2}) \otimes \dots \otimes (a_{i_w} x_1 + b_{i_w}), \quad (\text{D.27})$$

where by hypothesis the b_{i_k} are independent of x_1 . As we have subtracted of the contribution from $\phi_{x_1}(T)$, these terms cannot come from a multiple polylogarithm of the form $G(\dots; x_1)$ of weight w , but it can only arise from the product

$$\log b_{i_1} G\left(-\frac{b_{i_w}}{a_{i_w}}, \dots, -\frac{b_{i_2}}{a_{i_2}}; x_1\right) \rightarrow \phi_{x_2}(b_{i_1}) \phi_{x_1}\left((a_{i_2} x_1 + b_{i_2}) \otimes \dots \otimes (a_{i_w} x_1 + b_{i_w})\right). \quad (\text{D.28})$$

It is easy to convince oneself that the difference

$$T_2 = T_1 - \sum_{(i_1, \dots, i_w)} c_{i_1, \dots, i_w} \mathcal{S}\left[\phi_{x_2}(b_{i_1}) \phi_{x_1}\left((a_{i_2} x_1 + b_{i_2}) \otimes \dots \otimes (a_{i_w} x_1 + b_{i_w})\right)\right] \quad (\text{D.29})$$

contains only terms for which at most $(w-2)$ entries depend on x_1 . We can now go on and recursively subtract contributions with different multiplicities of x_1 . Assume for example that we have subtracted all contributions where x_1 appears in more than $(w-r)$ entries, and that the resulting symbol is T_r . We can then concentrate on the terms

$$\sum_{(i_1, \dots, i_w)} c_{i_1, \dots, i_w} b_{i_1} \otimes \dots \otimes b_{i_r} \otimes (a_{i_{r+1}} x_1 + b_{i_{r+1}}) \otimes \dots \otimes (a_{i_w} x_1 + b_{i_w}). \quad (\text{D.30})$$

It is easy to convince oneself that in the difference

$$T_{r+1} = T_r - \sum_{(i_1, \dots, i_w)} c_{i_1, \dots, i_w} \mathcal{S}\left[\phi_{x_2}(b_{i_1} \otimes \dots \otimes b_{i_r}) \phi_{x_1}\left((a_{i_{r+1}} x_1 + b_{i_{r+1}}) \otimes \dots \otimes (a_{i_w} x_1 + b_{i_w})\right)\right] \quad (\text{D.31})$$

x_1 appear in at most $(w-r-1)$ entries. We continue this procedure until we reach T_w , which is independent of x_1 , and we restart the algorithm with T_w and ϕ_{x_2} . We then repeat this procedure until we have exhausted all the integration variables, and the algorithm stops. The result of this algorithm is, by construction, a function of the form

$$\sum_{(i_1, \dots, i_n)} c_{i_1, \dots, i_n} G(\vec{a}_{i_n}; x_n) \dots G(\vec{a}_{i_1}; x_1), \quad (\text{D.32})$$

whose symbol is T and such that the $\vec{a}_{i_k}$ are sequences of rational functions that are in the variables x_k , $k > i_k$, i.e., the sought-for canonical form for T .

Using this algorithm we can easily take the limit $x_a \rightarrow \infty$, by applying it to the right-hand side of eq. (D.16) with the ordering $(\bar{x}_a, x_b, \dots)$. We can also use it to rewrite the integral over x_a such that we can easily integrate over the next integration variable x_b . However, the result of the algorithm will at this stage still deliver the wrong answer, as we have so far only constructed a function in canonical form whose *symbol* matches the symbol of a given function. To illustrate this, assume we run the algorithm on the function in the right-hand side of eq. (D.16) with the ordering $(\bar{x}_a, x_b, \dots)$. The result is a function $\mathcal{G}(\bar{x}_a, x_b, \dots)$ in canonical form such that

$$\mathcal{S} \left[G \left(\vec{a}; \frac{1}{\bar{x}_a} \right) - \mathcal{G}(\bar{x}_a, x_b, \dots) \right] = 0. \quad (\text{D.33})$$

It would however be wrong to conclude that $G \left(\vec{a}; \frac{1}{\bar{x}_a} \right)$ and $\mathcal{G}(\bar{x}_a, x_b, \dots)$ are equal, because the symbol maps to 0 all terms proportional to multiple zeta values. In ref. [128] an algorithm was described that allows to reconstruct these zeta-valued terms using the full Hopf algebra structure of multiple polylogarithms, augmented by some ideas by Brown [129]. In the following we only give a very brief account on how to reconstruct the zeta-valued terms, and we refer to ref. [128] for a detailed description of the algorithm.

In the following we assume for simplicity that the function $G \left(\vec{a}; \frac{1}{\bar{x}_a} \right) - \mathcal{G}(\bar{x}_a, x_b, \dots)$ is real,⁷ so we do not need to worry about imaginary parts proportional to $i\pi$. Next we act with $\Delta_{2,1,\dots,1}$ on the difference, where $\Delta_{2,1,\dots,1}$ is the component of the iterated coproduct where the first component has weight two and all other components have weight one. Without loss of generality we can write

$$\begin{aligned} & \Delta_{2,1,\dots,1} \left[G \left(\vec{a}; \frac{1}{\bar{x}_a} \right) - \mathcal{G}(\bar{x}_a, x_b, \dots) \right] \\ &= \sum_{(i_1, \dots, i_{w-1})} c_{i_1, \dots, i_{w-1}} A_{i_1} \otimes \log a_{i_2} \otimes \dots \otimes \log a_{i_{w-1}}, \end{aligned} \quad (\text{D.34})$$

where the a_{i_k} are irreducible polynomials and A_{i_1} is a combination of multiple polylogarithms of weight two. Without loss of generality we can assume that we have collected all term that have the same ‘tail’ $\log a_{i_2} \otimes \dots \otimes \log a_{i_{w-1}}$, i.e., we can assume

$$\log a_{i_2} \otimes \dots \otimes \log a_{i_{w-1}} \neq \log a_{j_2} \otimes \dots \otimes \log a_{j_{w-1}} \text{ if } (i_2, \dots, i_{w-1}) \neq (j_2, \dots, j_{w-1}). \quad (\text{D.35})$$

As we know that the symbol of the function vanishes, eq. (D.33), we necessarily conclude that A_{i_1} is proportional to ζ_2 , i.e., $A_{i_1} = k_{i_1} \zeta_2$, for some rational number k_{i_1} . This rational number can easily be determined by evaluating A_{i_1} numerically at a single point using any of the standard libraries to evaluate multiple polylogarithms [88, 130–135], and running for

⁷There is no obstacle to consider complex-valued functions. Imaginary parts can be extracted in exactly the same way using $\Delta_{1,\dots,1}$.

example the PSLQ algorithm [136]. Equation (D.34) then takes the form

$$\begin{aligned} \Delta_{2,1,\dots,1} \left[G \left(\vec{a}; \frac{1}{\bar{x}_a} \right) - \mathcal{G}(\bar{x}_a, x_b, \dots) \right] \\ = \sum_{(i_1, \dots, i_{w-1})} c_{i_1, \dots, i_{w-1}} k_{i_1} \zeta_2 \otimes \log a_{i_2} \otimes \dots \otimes \log a_{i_{w-1}}, \end{aligned} \quad (\text{D.36})$$

Next we drop ζ_2 , i.e., we only keep the tail of each elementary tensor. If we also drop the log signs, we obtain a symbol associated with the terms proportional to ζ_2 ,

$$\Delta_{2,1,\dots,1} \left[G \left(\vec{a}; \frac{1}{\bar{x}_a} \right) - \mathcal{G}(\bar{x}_a, x_b, \dots) \right] \rightarrow \sum_{(i_1, \dots, i_{w-1})} c_{i_1, \dots, i_{w-1}} k_{i_1} a_{i_2} \otimes \dots \otimes a_{i_{w-1}}. \quad (\text{D.37})$$

Running the algorithm described at the beginning of this section on the symbol in eq. (D.37) we obtain a function $\mathcal{G}_2(\bar{x}_a, x_b, \dots)$ of weight $w - 2$ in canonical form such that

$$\Delta_{2,1,\dots,1} \left[G \left(\vec{a}; \frac{1}{\bar{x}_a} \right) - \mathcal{G}(\bar{x}_a, x_b, \dots) - \zeta_2 \mathcal{G}_2(\bar{x}_a, x_b, \dots) \right] = 0. \quad (\text{D.38})$$

We have in this way determined all the contributions proportional to ζ_2 , and the result is by construction in canonical form. We then repeat exactly the same exercise by acting with $\Delta_{3,1,\dots,1}$ to determine the terms proportional to ζ_3 , and we continue in this way until we have exhausted all possibilities and the algorithm stops. As a result we obtain the expression of $G \left(\vec{a}; \frac{1}{\bar{x}_a} \right)$ in canonical form at *function level*.

This terminates the algorithm to bring multiple polylogarithms into the canonical form corresponding to a certain ordering of the variables. This operation is sufficient to take the limit $x_a \rightarrow \infty$, and to bring the integral into a form where the next integration can easily be performed. We have however not yet shown that this algorithm will always terminate on the class of integrals we consider. In particular, when formulating the algorithm, we assumed that all entries that appear in the symbol are linear in all variables. In the next subsection we show that this condition can be relaxed, and that the algorithm always terminates if the integrand satisfies the reduction criterion of the previous subsection and if the integration range is $[0, \infty]$.

D.3 Symbolic integration and denominator reduction

In this subsection we show that the algorithm we just described always terminates for integrals of the type (D.1) such that the set S of polynomials satisfies the reduction criterion. From now on we assume that we have found an ordering of the integration variables, which we take as $(x_1, \dots, x_n)$ and we can find a sequence

$$S, S_{(x_1)}, S_{(x_1, x_2)}, S_{(x_1, x_2, x_3)}, \dots \quad (\text{D.39})$$

such that all the elements of $S_{(x_1, \dots, x_k)}$ are linear in x_{k+1} . We start by showing that under these hypotheses and after having integrated out $(x_1, \dots, x_{k-1})$,

1. the symbol of the primitive with respect to x_k has all its entries drawn from the set $\bar{S}_{(x_1, \dots, x_{k-1})} = \{x_k, \dots, x_n\} \cup S_{(x_1, \dots, x_{k-1})}$.

2. the symbol of the function after integration over x_k , i.e., after taking the limits $x_k \rightarrow 0, \infty$ of the primitive, has all its entries drawn from $\tilde{S}_{(x_1, \dots, x_k)} = \{x_{k+1}, \dots, x_n\} \cup S_{(x_1, \dots, x_k)}$.

We start by proving the first statement, the second then immediately follows by taking the appropriate limits. As in the following we constantly switch between polynomials of the sets we just defined, and polynomials as entries of symbols, we introduce the notation $[P]$ to refer to P as an entry of a symbol. Note that we then have the identities

$$\begin{aligned} [PQ] &= [P] + [Q], \\ [P/Q] &= [P] - [Q], \\ [\pm 1] &= 0, \\ \text{etc.} \end{aligned} \tag{D.40}$$

We proceed by iteration in the number k of variables we have already integrated out. We start by analyzing what happens after the first integration. It is obvious from eqs. (D.5) and (D.9) that after the first integration the primitive only involves multiple polylogarithms $G(a_1, \dots, a_w; x_1)$ with

$$a_i \in \{0, -R_k^1/Q_k^1\}. \tag{D.41}$$

It is then easy to see (e.g., from the polygon approach to the symbol of ref. [127]) that the symbol can only have the following entries:

$$\begin{aligned} &[x_k], [R_k^1], \\ &[-R_k^1/Q_k^1] = [R_k^1] - [Q_k^1], \\ &\left[1 + \frac{x_1 Q_k^1}{R_k^1}\right] = [R_k^1 + Q_k^1 x_1] - [R_k^1] = [P_k] - [R_k^1], \\ &\left[1 + \frac{Q_k^1 R_l^1}{R_k^1 Q_l^1}\right] = [Q_k^1 R_l^1 - R_k^1 Q_l^1] - [R_k^1] - [Q_l^1]. \end{aligned} \tag{D.42}$$

The polynomials that appear inside the symbol are precisely those that appear in S and $S_{(x_1)}$, which finishes the proof of the first statement for the first integration.

Next consider taking the limits $x_1 \rightarrow 0$ and $x_1 \rightarrow \infty$. By definition, R_k^1 and Q_k^1 are independent of the limit, so we only need to consider the limits of $[x_1]$ and $[P_k]$. $[x_1]$ will give rise to logarithmic singularities in the limit, and these terms must cancel if the integral is convergent. For $[P_k]$ we have,

$$\begin{aligned} \lim_{x_1 \rightarrow 0} [P_k] &= [R_k^1], \\ \lim_{x_1 \rightarrow \infty} [P_k] &= \lim_{\bar{x}_1 \rightarrow 0} \{[\bar{x}_1 R_k^1 + Q_k^1] - [\bar{x}_1]\} = [Q_k^1] + \lim_{\bar{x}_1 \rightarrow 0} [x_1]. \end{aligned} \tag{D.43}$$

The logarithmic singularity in the last line must again cancel for convergent integrals, and so we see that the only polynomials that appear in the limit are those in $S_{(x_1)}$. This finishes the proof of the second statement for the first integration. We stress that it is important that the integration region is $[0, \infty]$, because otherwise we have to take into account effects coming from the integration boundaries.

Let us now suppose that the two statements are true for the first $r - 1$ integrations. The set $S_{(x_1, \dots, x_{r-1})}$ then consists of polynomials of the form $\tilde{P}_k = \tilde{Q}_k^r x_r + \tilde{R}_k^r$. Let us now compute the primitive with respect to x_r . We start by running the algorithm to bring the multiple polylogarithms in the integrand into canonical form. As this involves the application of the map ϕ_{x_r} , we obtain multiple polylogarithms of the form $G(a_1, \dots, a_w; x_r)$ with

$$a_i \in \left\{ 0, -\tilde{R}_k^r / \tilde{Q}_k^r \right\}. \quad (\text{D.44})$$

If we compute the primitive, we integrate over linear functions in x_r from the set $S_{(x_1, \dots, x_{r-1})}$, and it is easy to see that this does not change eq. (D.44). Using exactly the same argument as for the first integration, we see that these multiple polylogarithms only contribute terms to the symbols whose entries are drawn from $\bar{S}_{(x_1, \dots, x_{r-1})}$. In addition we have multiple polylogarithms that are independent of x_r , whose symbols involve those elements of $\tilde{S}_{(x_1, \dots, x_{r-1})}$ that are independent of x_r . As these functions do not change if we take the primitive with respect to x_r , we see that they do not alter the conclusion. So the symbol of the primitive with respect to x_r must have all its entries drawn from $\bar{S}_{(x_1, \dots, x_{r-1})}$. Taking the limits $x_r \rightarrow 0$ and $x_r \rightarrow \infty$ just like for the first integration then finishes the proof.

Having proved the two statements, we can show that our algorithm always terminates for the class of integral we consider. More precisely, we have to show that our algorithm can always produce the canonical form for the next integration step. This is done by applying the map ϕ_{x_r} , which requires all the entries in the symbol to be either independent of x_r or linear in x_r . By construction, this condition is always fulfilled for the elements of $\tilde{S}_{(x_1, \dots, x_{r-1})}$. It is easy to check that the same argument shows that the map $\phi_{x_{r+1}}$, which is called recursively by ϕ_{x_r} is well-defined, and so we can always find a canonical form for the integrand.

D.4 A toy example

In this section we discuss an example that illustrates how the algorithm described in the previous subsection can be used to compute parametric integrals. In particular, we used this algorithm to compute the parametric integrals (8.68) and (8.69). While these integrals are a straightforward application of the algorithm, intermediate expression are rather long, so we prefer to use a simpler integral where we can explicitly show all the steps. The integral we are going to consider is

$$\begin{aligned} \mathcal{I}(\epsilon) = \int_0^\infty dx_1 dx_2 dx_3 x_1^\epsilon (1+x_1)^{3\epsilon-2} x_2^{-\epsilon} (1+x_2)^{-4\epsilon-2} x_3^{2\epsilon} (1+x_3)^{-\epsilon-1} \\ \times (1+x_2+x_3+x_1x_3)^{-2\epsilon-1}. \end{aligned} \quad (\text{D.45})$$

The integral is finite as $\epsilon \rightarrow 0$, and we want to compute the first few terms in the Taylor expansion

$$\mathcal{I}(\epsilon) = \mathcal{I}_0 + \mathcal{I}_1 \epsilon + \mathcal{I}_2 \epsilon^2 + \mathcal{I}_3 \epsilon^3 + \mathcal{O}(\epsilon^4). \quad (\text{D.46})$$

The first coefficient is trivial to compute,

$$\mathcal{I}_0 = \frac{\pi^2}{9} - \frac{2}{3}. \quad (\text{D.47})$$

We will now illustrate our algorithm in detail on the coefficient $\mathcal{I}_1$. If we integrate out the integration variables in the order (x_1, x_2, x_3) , we obtain

$$\begin{aligned} S &= \{1 + x_1, 1 + x_2, 1 + x_3, 1 + x_2 + x_3 + x_1x_3\}, \\ S_{(x_1)} &= \{1 + x_2, 1 + x_3, 1 + x_2 + x_3\}, \\ S_{(x_1, x_2)} &= \{1 + x_3\}. \end{aligned} \quad (\text{D.48})$$

We see that all the sets are linear in all integration variables, so we perform the integrations one after the other.

The coefficient $\mathcal{I}_1$ is given by the integral

$$\begin{aligned} \mathcal{I}_1 &= \int_0^\infty \frac{dx_1 dx_2 dx_3}{(1+x_1)^2 (1+x_2)^2 (1+x_3) (1+x_2+x_3+x_1x_3)} \\ &\times \left[3G(-1; x_1) - 4G(-1; x_2) - 3G(-1; x_3) + G(0; x_1) - G(0; x_2) \right. \\ &\quad \left. + 2G(0; x_3) - 2G(-x_3-1; x_2) - 2G\left(\frac{-x_2-x_3-1}{x_3}; x_1\right) \right], \end{aligned} \quad (\text{D.49})$$

where we have already written all logarithms in terms of multiple polylogarithms, e.g.,

$$\begin{aligned} \log(1+x_2+x_3+x_1x_3) &= \log(1+x_3) + \log\left(1+\frac{x_2}{1+x_3}\right) + \log\left(1+\frac{x_1x_3}{1+x_2+x_3}\right) \\ &= G(-1; x_3) + G(-1-x_3; x_2) + G\left(\frac{-x_2-x_3-1}{x_3}; x_1\right). \end{aligned} \quad (\text{D.50})$$

It is easy to compute a primitive with respect to x_1 for the integrand of $\mathcal{I}_1$, e.g.,

$$\int \frac{dx_1}{1+x_2+x_3+x_1x_3} G(-1; x_1) = \frac{1}{x_3} G\left(\frac{-x_2-x_3-1}{x_3}, -1; x_1\right). \quad (\text{D.51})$$

In the following we only concentrate on this single term (which is in fact the most complicated one) to illustrate the procedure. All other terms can be dealt with in a similar way. Before we compute the limit of eq. (D.51) as $x_1 \rightarrow 0, \infty$, let us comment about the symbol of the primitive. We have

$$\begin{aligned} \mathcal{S}\left[G\left(\frac{-x_2-x_3-1}{x_3}, -1; x_1\right)\right] &= -(1+x_1) \otimes (1+x_2) \\ &\quad + (1+x_1) \otimes (1+x_2+x_3+x_1x_3) - (1+x_2+x_3) \otimes (1+x_2) \\ &\quad + (1+x_2+x_3) \otimes x_3 + (1+x_2+x_3+x_1x_3) \otimes (1+x_2) \\ &\quad - (1+x_2+x_3+x_1x_3) \otimes x_3. \end{aligned} \quad (\text{D.52})$$

We see that the entries in the symbol are drawn from the set $S \cup \{x_1, x_2, x_3\}$, as expected. The same is true for all other terms in the primitive.

The limit $x_1 \rightarrow 0$ is trivial,

$$\lim_{x_1 \rightarrow 0} G\left(\frac{-x_2-x_3-1}{x_3}, -1; x_1\right) = 0. \quad (\text{D.53})$$

The limit $x_1 \rightarrow \infty$ is obtained by letting $x_1 = 1/\bar{x}_1$ and deriving the inversion relation for this multiple polylogarithm. This is equivalent to bringing $G(-1, (-x_2 - x_3 - 1)/x_3; 1/\bar{x}_1)$ into canonical form with respect to the ordering of variables $(\bar{x}_1, x_2, x_3)$. We start by using our algorithm to construct a function $\mathcal{G}(\bar{x}_1, x_2, x_3)$ which is in canonical form and has the symbol given in eq. (D.52). We find

$$\begin{aligned} \mathcal{G}(\bar{x}_1, x_2, x_3) = & G\left(-\frac{x_3}{x_2 + x_3 + 1}, -1; \bar{x}_1\right) - G\left(-\frac{x_3}{x_2 + x_3 + 1}, 0; \bar{x}_1\right) \\ & - G(0, -1; \bar{x}_1) + G(0, 0; \bar{x}_1) - G(-1, -x_3 - 1; x_2) \\ & + G(-1; x_2) [G(0; x_3) - G(-1; x_3)] + G(0, -1; x_3) - G(0, 0; x_3). \end{aligned} \quad (\text{D.54})$$

It is easy to check that the symbol of $\mathcal{G}(\bar{x}_1, x_2, x_3)$ agrees with eq. (D.52). It would however be wrong to conclude that the *functions* are equal — they might differ by a rational multiple of ζ_2 . Evaluating the difference numerically at a single point, we obtain

$$G\left(\frac{-x_2 - x_3 - 1}{x_3}, -1; \frac{1}{\bar{x}_1}\right) - \mathcal{G}(\bar{x}_1, x_2, x_3) = -1.64493406684822643\dots = -\zeta_2. \quad (\text{D.55})$$

Taking the limit $x_1 \rightarrow \infty$ is now trivial, and we obtain

$$\begin{aligned} G\left(-1, \frac{-x_2 - x_3 - 1}{x_3}; x_1\right) = & G(0, 0; x_1) - G(-1; x_2) G(-1; x_3) \\ & + G(-1; x_2) G(0; x_3) - G(-1, -x_3 - 1; x_2) + G(0, -1; x_3) - G(0, 0; x_3) \\ & - \zeta_2 + \mathcal{O}(1/x_1). \end{aligned} \quad (\text{D.56})$$

Note that the function has a logarithmic singularity for $x_1 \rightarrow \infty$, which will cancel against similar contributions from other terms. Furthermore, note that the symbols of all the (finite) terms in eq. (D.56) have entries drawn from the set $S_{(x_1)} \cup \{x_2, x_3\}$, as expected. The previous steps can easily be implemented into a computer code and repeated for all the terms appearing in the primitive with respect to x_1 .

The result of the integration over x_1 is, by construction, already in canonical form with respect to (x_2, x_3) , and we can immediately compute the primitive with respect to x_2 , e.g.,

$$\int \frac{dx_2}{1 + x_2 + x_3} G(-1; x_2) = G(-1 - x_3, -1; x_2). \quad (\text{D.57})$$

The limit $x_2 \rightarrow 0$ is again trivial, while the limit $x_2 \rightarrow \infty$ can again be computed by letting $x_2 = 1/\bar{x}_2$ and constructing a function $\mathcal{G}(\bar{x}_2, x_3)$ in canonical form with the same symbol. We find

$$\begin{aligned} \mathcal{G}(\bar{x}_2, x_3) = & G\left(-\frac{1}{x_3 + 1}, -1; \bar{x}_2\right) - G\left(-\frac{1}{x_3 + 1}, 0; \bar{x}_2\right) - G(0, -1; \bar{x}_2) \\ & + G(0, 0; \bar{x}_2) - G(0, -1; x_3). \end{aligned} \quad (\text{D.58})$$

Numerical evaluation at a single point immediately shows that we do not need to add any term proportional to ζ_2 . Taking the limit is now trivial, and we get

$$G(-1 - x_3, -1; x_2) = G(0, 0; x_2) - G(0, -1; x_3) + \mathcal{O}(1/x_2). \quad (\text{D.59})$$

We are finally only left with the integral over x_3 . The primitive involves integrals like

$$\int \frac{dx_3}{1+x_3} G(-1, 0; x_3) = G(-1, -1, 0; x_3). \quad (\text{D.60})$$

The limit $x_3 \rightarrow 0$ is trivial and for the limit $x_3 \rightarrow \infty$ we construct a function $\mathcal{G}(\bar{x}_3)$ with the same symbol as $G(-1, -1, 0; 1/\bar{x}_3)$. We find

$$\mathcal{G}(\bar{x}_3) = G(-1, 0, 0; \bar{x}_3) - G(-1, -1, 0; \bar{x}_3) + G(0, -1, 0; \bar{x}_3) - G(0, 0, 0; \bar{x}_3). \quad (\text{D.61})$$

Next we have to determine terms proportional to ζ_2 . This is achieved by acting with $\Delta_{2,1}$ on the difference,

$$\begin{aligned} \Delta_{2,1} [G(-1, -1, 0; 1/\bar{x}_3) - \mathcal{G}(\bar{x}_3)] \\ = [G(-1, 0; \bar{x}_3) + G(-1, 0; \bar{x}_3) - G(0, 0; \bar{x}_3)] \otimes G(-1; \bar{x}_3) \\ + \left[G(0, 0; \bar{x}_3) - G\left(-1, 0; \frac{1}{\bar{x}_3}\right) - G(-1, 0; \bar{x}_3) \right] \otimes G(0; \bar{x}_3). \end{aligned} \quad (\text{D.62})$$

Evaluating the first entries numerically at a single point reveals

$$\begin{aligned} \Delta_{2,1} [G(-1, -1, 0; 1/\bar{x}_3) - \mathcal{G}(\bar{x}_3)] &= -\zeta_2 \otimes G(-1; \bar{x}_3) + \zeta_2 \otimes G(0; \bar{x}_3) \\ &= \Delta_{2,1} [-\zeta_2 G(-1; \bar{x}_3) + \zeta_2 G(0; \bar{x}_3)]. \end{aligned} \quad (\text{D.63})$$

Finally, evaluating the full difference at a single point, we obtain

$$\begin{aligned} G(-1, -1, 0; 1/\bar{x}_3) - [\mathcal{G}(\bar{x}_3) - \zeta_2 G(-1; \bar{x}_3) + \zeta_2 G(0; \bar{x}_3)] &= 1.20205690315959 \dots \\ &= \zeta_3. \end{aligned} \quad (\text{D.64})$$

Taking the limit $\bar{x}_3$ is now trivial, and we finally get

$$\mathcal{I}_1 = -5\zeta_3 + \frac{2\pi^2}{9} + \frac{5}{3}. \quad (\text{D.65})$$

The higher terms in the ϵ expansion can be obtained in exactly the same way. For this particular integral we find for example

$$\begin{aligned} \mathcal{I}_2 &= \frac{149\pi^4}{216} - 10\zeta_3 - \frac{16\pi^2}{9} - \frac{157}{6}, \\ \mathcal{I}_3 &= -\frac{910}{3}\zeta_5 + \frac{149\pi^4}{108} + \frac{607}{6}\zeta_3 - \frac{277\pi^2}{18}\zeta_3 + \frac{29\pi^2}{3} + \frac{1175}{12}. \end{aligned} \quad (\text{D.66})$$

References

- [1] Z. Bern et al., *Four-Jet Production at the Large Hadron Collider at Next-to-Leading Order in QCD*, *Phys. Rev. Lett.* **109** (2012) 042001 [[arXiv:1112.3940](#)] [[INSPIRE](#)].
- [2] H. Ita et al., *Precise Predictions for $Z + 4$ Jets at Hadron Colliders*, *Phys. Rev. D* **85** (2012) 031501 [[arXiv:1108.2229](#)] [[INSPIRE](#)].
- [3] C. Berger et al., *Precise Predictions for $W + 4$ Jet Production at the Large Hadron Collider*, *Phys. Rev. Lett.* **106** (2011) 092001 [[arXiv:1009.2338](#)] [[INSPIRE](#)].

- [4] C. Berger et al., *Next-to-Leading Order QCD Predictions for $W+3$ -Jet Distributions at Hadron Colliders*, *Phys. Rev. D* **80** (2009) 074036 [[arXiv:0907.1984](#)] [[INSPIRE](#)].
- [5] C. Berger et al., *Precise Predictions for $W + 3$ Jet Production at Hadron Colliders*, *Phys. Rev. Lett.* **102** (2009) 222001 [[arXiv:0902.2760](#)] [[INSPIRE](#)].
- [6] R.K. Ellis, K. Melnikov and G. Zanderighi, *$W+3$ jet production at the Tevatron*, *Phys. Rev. D* **80** (2009) 094002 [[arXiv:0906.1445](#)] [[INSPIRE](#)].
- [7] T. Melia, K. Melnikov, R. Rontsch and G. Zanderighi, *NLO QCD corrections for W^+W^- pair production in association with two jets at hadron colliders*, *Phys. Rev. D* **83** (2011) 114043 [[arXiv:1104.2327](#)] [[INSPIRE](#)].
- [8] K. Melnikov and G. Zanderighi, *$W+3$ jet production at the LHC as a signal or background*, *Phys. Rev. D* **81** (2010) 074025 [[arXiv:0910.3671](#)] [[INSPIRE](#)].
- [9] V. Hirschi, R. Frederix, S. Frixione, M.V. Garzelli, F. Maltoni and R. Pittau, *Automation of one-loop QCD corrections*, *JHEP* **05** (2011) 044 [[arXiv:1103.0621](#)] [[INSPIRE](#)].
- [10] G. Bevilacqua, M. Czakon, C. Papadopoulos and M. Worek, *Hadronic top-quark pair production in association with two jets at Next-to-Leading Order QCD*, *Phys. Rev. D* **84** (2011) 114017 [[arXiv:1108.2851](#)] [[INSPIRE](#)].
- [11] G. Bevilacqua et al., *HELAC-NLO*, *Comput. Phys. Commun.* **184** (2013) 986 [[arXiv:1110.1499](#)] [[INSPIRE](#)].
- [12] G. Bevilacqua, M. Czakon, C. Papadopoulos, R. Pittau and M. Worek, *Assault on the NLO Wishlist: $pp \rightarrow t\bar{t}b\bar{b}$* , *JHEP* **09** (2009) 109 [[arXiv:0907.4723](#)] [[INSPIRE](#)].
- [13] G. Bevilacqua, M. Czakon, C. Papadopoulos and M. Worek, *Dominant QCD Backgrounds in Higgs Boson Analyses at the LHC: A Study of $pp \rightarrow t\bar{t} + 2$ jets at Next-To-Leading Order*, *Phys. Rev. Lett.* **104** (2010) 162002 [[arXiv:1002.4009](#)] [[INSPIRE](#)].
- [14] A. Bredenstein, A. Denner, S. Dittmaier and S. Pozzorini, *NLO QCD corrections to $pp \rightarrow t\bar{t}b\bar{b} + X$ at the LHC*, *Phys. Rev. Lett.* **103** (2009) 012002 [[arXiv:0905.0110](#)] [[INSPIRE](#)].
- [15] A. Denner, S. Dittmaier, S. Kallweit and S. Pozzorini, *NLO QCD corrections to $WWb\bar{b}$ production at hadron colliders*, *Phys. Rev. Lett.* **106** (2011) 052001 [[arXiv:1012.3975](#)] [[INSPIRE](#)].
- [16] F. Cascioli, P. Maierhofer and S. Pozzorini, *Scattering Amplitudes with Open Loops*, *Phys. Rev. Lett.* **108** (2012) 111601 [[arXiv:1111.5206](#)] [[INSPIRE](#)].
- [17] R.K. Ellis, Z. Kunszt, K. Melnikov and G. Zanderighi, *One-loop calculations in quantum field theory: from Feynman diagrams to unitarity cuts*, *Phys. Rept.* **518** (2012) 141 [[arXiv:1105.4319](#)] [[INSPIRE](#)].
- [18] R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, R. Pittau and P. Torrielli, *Four-lepton production at hadron colliders: aMC@NLO predictions with theoretical uncertainties*, *JHEP* **02** (2012) 099 [[arXiv:1110.4738](#)] [[INSPIRE](#)].
- [19] S. Catani, F. Krauss, R. Kuhn and B. Webber, *QCD matrix elements + parton showers*, *JHEP* **11** (2001) 063 [[hep-ph/0109231](#)] [[INSPIRE](#)].
- [20] L. Lönnblad, *Correcting the color dipole cascade model with fixed order matrix elements*, *JHEP* **05** (2002) 046 [[hep-ph/0112284](#)] [[INSPIRE](#)].

- [21] F. Krauss, *Matrix elements and parton showers in hadronic interactions*, *JHEP* **08** (2002) 015 [[hep-ph/0205283](#)] [[INSPIRE](#)].
- [22] S. Frixione and B.R. Webber, *Matching NLO QCD computations and parton shower simulations*, *JHEP* **06** (2002) 029 [[hep-ph/0204244](#)] [[INSPIRE](#)].
- [23] P. Nason, *A New method for combining NLO QCD with shower Monte Carlo algorithms*, *JHEP* **11** (2004) 040 [[hep-ph/0409146](#)] [[INSPIRE](#)].
- [24] N. Lavesson and L. Lönnblad, *W+jets matrix elements and the dipole cascade*, *JHEP* **07** (2005) 054 [[hep-ph/0503293](#)] [[INSPIRE](#)].
- [25] M.L. Mangano, M. Moretti, F. Piccinini and M. Treccani, *Matching matrix elements and shower evolution for top-quark production in hadronic collisions*, *JHEP* **01** (2007) 013 [[hep-ph/0611129](#)] [[INSPIRE](#)].
- [26] J. Alwall et al., *Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions*, *Eur. Phys. J. C* **53** (2008) 473 [[arXiv:0706.2569](#)] [[INSPIRE](#)].
- [27] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with Parton Shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070 [[arXiv:0709.2092](#)] [[INSPIRE](#)].
- [28] J. Alwall, S. de Visscher and F. Maltoni, *QCD radiation in the production of heavy colored particles at the LHC*, *JHEP* **02** (2009) 017 [[arXiv:0810.5350](#)] [[INSPIRE](#)].
- [29] S. Hoeche, F. Krauss, S. Schumann and F. Siegert, *QCD matrix elements and truncated showers*, *JHEP* **05** (2009) 053 [[arXiv:0903.1219](#)] [[INSPIRE](#)].
- [30] K. Hamilton, P. Richardson and J. Tully, *A Modified CKKW matrix element merging approach to angular-ordered parton showers*, *JHEP* **11** (2009) 038 [[arXiv:0905.3072](#)] [[INSPIRE](#)].
- [31] K. Hamilton and P. Nason, *Improving NLO-parton shower matched simulations with higher order matrix elements*, *JHEP* **06** (2010) 039 [[arXiv:1004.1764](#)] [[INSPIRE](#)].
- [32] S. Hoeche, F. Krauss, M. Schonherr and F. Siegert, *Automating the POWHEG method in Sherpa*, *JHEP* **04** (2011) 024 [[arXiv:1008.5399](#)] [[INSPIRE](#)].
- [33] S. Hoeche, F. Krauss, M. Schonherr and F. Siegert, *NLO matrix elements and truncated showers*, *JHEP* **08** (2011) 123 [[arXiv:1009.1127](#)] [[INSPIRE](#)].
- [34] L. Lönnblad and S. Prestel, *Matching Tree-Level Matrix Elements with Interleaved Showers*, *JHEP* **03** (2012) 019 [[arXiv:1109.4829](#)] [[INSPIRE](#)].
- [35] S. Hoeche, F. Krauss, M. Schonherr and F. Siegert, *A critical appraisal of NLO+PS matching methods*, *JHEP* **09** (2012) 049 [[arXiv:1111.1220](#)] [[INSPIRE](#)].
- [36] K. Hamilton, P. Nason and G. Zanderighi, *MINLO: Multi-Scale Improved NLO*, *JHEP* **10** (2012) 155 [[arXiv:1206.3572](#)] [[INSPIRE](#)].
- [37] S. Hoeche, F. Krauss, M. Schonherr and F. Siegert, *QCD matrix elements + parton showers: The NLO case*, *JHEP* **04** (2013) 027 [[arXiv:1207.5030](#)] [[INSPIRE](#)].
- [38] T. Gehrmann, S. Hoeche, F. Krauss, M. Schonherr and F. Siegert, *NLO QCD matrix elements + parton showers in $e^+e^- \rightarrow \text{hadrons}$* , *JHEP* **01** (2013) 144 [[arXiv:1207.5031](#)] [[INSPIRE](#)].
- [39] R. Frederix and S. Frixione, *Merging meets matching in MC@NLO*, *JHEP* **12** (2012) 061 [[arXiv:1209.6215](#)] [[INSPIRE](#)].

- [40] S. Buehler, F. Herzog, A. Lazopoulos and R. Mueller, *The Fully differential hadronic production of a Higgs boson via bottom quark fusion at NNLO*, *JHEP* **07** (2012) 115 [[arXiv:1204.4415](#)] [[INSPIRE](#)].
- [41] A. G.-D. Ridder, T. Gehrmann, E. Glover and J. Pires, *Second order QCD corrections to jet production at hadron colliders: the all-gluon contribution*, [arXiv:1301.7310](#) [[INSPIRE](#)].
- [42] M. Czakon and A. Mitov, *NNLO corrections to top pair production at hadron colliders: the quark-gluon reaction*, *JHEP* **01** (2013) 080 [[arXiv:1210.6832](#)] [[INSPIRE](#)].
- [43] M. Czakon and A. Mitov, *NNLO corrections to top-pair production at hadron colliders: the all-fermionic scattering channels*, *JHEP* **12** (2012) 054 [[arXiv:1207.0236](#)] [[INSPIRE](#)].
- [44] P. Baernreuther, M. Czakon and A. Mitov, *Percent Level Precision Physics at the Tevatron: First Genuine NNLO QCD Corrections to $q\bar{q} \rightarrow t\bar{t} + X$* , *Phys. Rev. Lett.* **109** (2012) 132001 [[arXiv:1204.5201](#)] [[INSPIRE](#)].
- [45] S. Catani, L. Cieri, D. de Florian, G. Ferrera and M. Grazzini, *Diphoton production at hadron colliders: a fully-differential QCD calculation at NNLO*, *Phys. Rev. Lett.* **108** (2012) 072001 [[arXiv:1110.2375](#)] [[INSPIRE](#)].
- [46] S. Catani, L. Cieri, G. Ferrera, D. de Florian and M. Grazzini, *Vector boson production at hadron colliders: A Fully exclusive QCD calculation at NNLO*, *Phys. Rev. Lett.* **103** (2009) 082001 [[arXiv:0903.2120](#)] [[INSPIRE](#)].
- [47] G. Ferrera, M. Grazzini and F. Tramontano, *Associated WH production at hadron colliders: a fully exclusive QCD calculation at NNLO*, *Phys. Rev. Lett.* **107** (2011) 152003 [[arXiv:1107.1164](#)] [[INSPIRE](#)].
- [48] M. Grazzini, *NNLO predictions for the Higgs boson signal in the $H \rightarrow WW \rightarrow l\nu l\nu$ and $H \rightarrow ZZ \rightarrow 4l$ decay channels*, *JHEP* **02** (2008) 043 [[arXiv:0801.3232](#)] [[INSPIRE](#)].
- [49] K. Melnikov and F. Petriello, *Electroweak gauge boson production at hadron colliders through $O(\alpha_s^2)$* , *Phys. Rev. D* **74** (2006) 114017 [[hep-ph/0609070](#)] [[INSPIRE](#)].
- [50] C. Anastasiou, G. Dissertori and F. Stockli, *NNLO QCD predictions for the $H \rightarrow WW \rightarrow l\nu l\nu$ signal at the LHC*, *JHEP* **09** (2007) 018 [[arXiv:0707.2373](#)] [[INSPIRE](#)].
- [51] C. Anastasiou, K. Melnikov and F. Petriello, *Fully differential Higgs boson production and the di-photon signal through next-to-next-to-leading order*, *Nucl. Phys. B* **724** (2005) 197 [[hep-ph/0501130](#)] [[INSPIRE](#)].
- [52] C. Anastasiou, K. Melnikov and F. Petriello, *Higgs boson production at hadron colliders: Differential cross sections through next-to-next-to-leading order*, *Phys. Rev. Lett.* **93** (2004) 262002 [[hep-ph/0409088](#)] [[INSPIRE](#)].
- [53] D. Graudenz, M. Spira and P. Zerwas, *QCD corrections to Higgs boson production at proton proton colliders*, *Phys. Rev. Lett.* **70** (1993) 1372 [[INSPIRE](#)].
- [54] S. Dawson, *Radiative corrections to Higgs boson production*, *Nucl. Phys. B* **359** (1991) 283 [[INSPIRE](#)].
- [55] A. Djouadi, M. Spira and P. Zerwas, *Production of Higgs bosons in proton colliders: QCD corrections*, *Phys. Lett. B* **264** (1991) 440 [[INSPIRE](#)].
- [56] M. Spira, A. Djouadi, D. Graudenz and P. Zerwas, *Higgs boson production at the LHC*, *Nucl. Phys. B* **453** (1995) 17 [[hep-ph/9504378](#)] [[INSPIRE](#)].

- [57] V. Ravindran, J. Smith and W.L. van Neerven, *NNLO corrections to the total cross-section for Higgs boson production in hadron hadron collisions*, *Nucl. Phys. B* **665** (2003) 325 [[hep-ph/0302135](#)] [[INSPIRE](#)].
- [58] R.V. Harlander and W.B. Kilgore, *Next-to-next-to-leading order Higgs production at hadron colliders*, *Phys. Rev. Lett.* **88** (2002) 201801 [[hep-ph/0201206](#)] [[INSPIRE](#)].
- [59] C. Anastasiou and K. Melnikov, *Higgs boson production at hadron colliders in NNLO QCD*, *Nucl. Phys. B* **646** (2002) 220 [[hep-ph/0207004](#)] [[INSPIRE](#)].
- [60] C. Anastasiou, S. Buehler, F. Herzog and A. Lazopoulos, *Total cross-section for Higgs boson hadroproduction with anomalous Standard Model interactions*, *JHEP* **12** (2011) 058 [[arXiv:1107.0683](#)] [[INSPIRE](#)].
- [61] R. Hamberg, W. van Neerven and T. Matsuura, *A Complete calculation of the order α_s^2 correction to the Drell-Yan K factor*, *Nucl. Phys. B* **359** (1991) 343 [Erratum *ibid.* **B 644** (2002) 403] [[INSPIRE](#)].
- [62] C. Anastasiou, L.J. Dixon, K. Melnikov and F. Petriello, *Dilepton rapidity distribution in the Drell-Yan process at NNLO in QCD*, *Phys. Rev. Lett.* **91** (2003) 182002 [[hep-ph/0306192](#)] [[INSPIRE](#)].
- [63] S. Moch and A. Vogt, *Higher-order soft corrections to lepton pair and Higgs boson production*, *Phys. Lett. B* **631** (2005) 48 [[hep-ph/0508265](#)] [[INSPIRE](#)].
- [64] C. Anastasiou and K. Melnikov, *Pseudoscalar Higgs boson production at hadron colliders in NNLO QCD*, *Phys. Rev. D* **67** (2003) 037501 [[hep-ph/0208115](#)] [[INSPIRE](#)].
- [65] C. Anastasiou, L.J. Dixon and K. Melnikov, *NLO Higgs boson rapidity distributions at hadron colliders*, *Nucl. Phys. Proc. Suppl.* **116** (2003) 193 [[hep-ph/0211141](#)] [[INSPIRE](#)].
- [66] C. Anastasiou, L.J. Dixon, K. Melnikov and F. Petriello, *High precision QCD at hadron colliders: Electroweak gauge boson rapidity distributions at NNLO*, *Phys. Rev. D* **69** (2004) 094008 [[hep-ph/0312266](#)] [[INSPIRE](#)].
- [67] S. Laporta, *High precision calculation of multiloop Feynman integrals by difference equations*, *Int. J. Mod. Phys. A* **15** (2000) 5087 [[hep-ph/0102033](#)] [[INSPIRE](#)].
- [68] T. Gehrmann and E. Remiddi, *Differential equations for two loop four point functions*, *Nucl. Phys. B* **580** (2000) 485 [[hep-ph/9912329](#)] [[INSPIRE](#)].
- [69] F. Tkachov, *A Theorem on Analytical Calculability of Four Loop Renormalization Group Functions*, *Phys. Lett. B* **100** (1981) 65 [[INSPIRE](#)].
- [70] K. Chetyrkin and F. Tkachov, *Integration by Parts: The Algorithm to Calculate β -functions in 4 Loops*, *Nucl. Phys. B* **192** (1981) 159 [[INSPIRE](#)].
- [71] C. Anastasiou, S. Buehler, C. Duhr and F. Herzog, *NNLO phase space master integrals for two-to-one inclusive cross sections in dimensional regularization*, *JHEP* **11** (2012) 062 [[arXiv:1208.3130](#)] [[INSPIRE](#)].
- [72] A. Pak, M. Rogal and M. Steinhauser, *Production of scalar and pseudo-scalar Higgs bosons to next-to-next-to-leading order at hadron colliders*, *JHEP* **09** (2011) 088 [[arXiv:1107.3391](#)] [[INSPIRE](#)].
- [73] E. Remiddi and J. Vermaseren, *Harmonic polylogarithms*, *Int. J. Mod. Phys. A* **15** (2000) 725 [[hep-ph/9905237](#)] [[INSPIRE](#)].

- [74] O. Tarasov, *Connection between Feynman integrals having different values of the space-time dimension*, *Phys. Rev. D* **54** (1996) 6479 [[hep-th/9606018](#)] [[INSPIRE](#)].
- [75] V.A. Smirnov and O. Veretin, *Analytical results for dimensionally regularized massless on-shell double boxes with arbitrary indices and numerators*, *Nucl. Phys. B* **566** (2000) 469 [[hep-ph/9907385](#)] [[INSPIRE](#)].
- [76] C. Anastasiou, E.N. Glover and C. Oleari, *The two loop scalar and tensor pentabox graph with lightlike legs*, *Nucl. Phys. B* **575** (2000) 416 [Erratum *ibid.* **B 585** (2000) 763] [[hep-ph/9912251](#)] [[INSPIRE](#)].
- [77] C. Anastasiou, T. Gehrmann, C. Oleari, E. Remiddi and J. Tausk, *The Tensor reduction and master integrals of the two loop massless crossed box with lightlike legs*, *Nucl. Phys. B* **580** (2000) 577 [[hep-ph/0003261](#)] [[INSPIRE](#)].
- [78] C. Anastasiou, J. Tausk and M. Tejeda-Yeomans, *The On-shell massless planar double box diagram with an irreducible numerator*, *Nucl. Phys. Proc. Suppl.* **89** (2000) 262 [[hep-ph/0005328](#)] [[INSPIRE](#)].
- [79] R. Lee, *Space-time dimensionality D as complex variable: Calculating loop integrals using dimensional recurrence relation and analytical properties with respect to D* , *Nucl. Phys. B* **830** (2010) 474 [[arXiv:0911.0252](#)] [[INSPIRE](#)].
- [80] R.N. Lee and V.A. Smirnov, *The Dimensional Recurrence and Analyticity Method for Multicomponent Master Integrals: Using Unitarity Cuts to Construct Homogeneous Solutions*, *JHEP* **12** (2012) 104 [[arXiv:1209.0339](#)] [[INSPIRE](#)].
- [81] G. Somogyi, *Angular integrals in d dimensions*, *J. Math. Phys.* **52** (2011) 083501 [[arXiv:1101.3557](#)] [[INSPIRE](#)].
- [82] W. van Neerven, *Dimensional regularization of mass and infrared singularities in two loop on-shell vertex functions*, *Nucl. Phys. B* **268** (1986) 453 [[INSPIRE](#)].
- [83] Y. Schröder and M. Steinhauser, *Four-loop decoupling relations for the strong coupling*, *JHEP* **01** (2006) 051 [[hep-ph/0512058](#)] [[INSPIRE](#)].
- [84] K. Chetyrkin, J.H. Kuhn and C. Sturm, *QCD decoupling at four loops*, *Nucl. Phys. B* **744** (2006) 121 [[hep-ph/0512060](#)] [[INSPIRE](#)].
- [85] E. Furlan, *Gluon-fusion Higgs production at NNLO for a non-standard Higgs sector*, *JHEP* **10** (2011) 115 [[arXiv:1106.4024](#)] [[INSPIRE](#)].
- [86] S. Catani and M. Grazzini, *Infrared factorization of tree level QCD amplitudes at the next-to-next-to-leading order and beyond*, *Nucl. Phys. B* **570** (2000) 287 [[hep-ph/9908523](#)] [[INSPIRE](#)].
- [87] P. Nogueira, *Automatic Feynman graph generation*, *J. Comput. Phys.* **105** (1993) 279 [[INSPIRE](#)].
- [88] C.W. Bauer, A. Frink and R. Kreckel, *Introduction to the GiNaC framework for symbolic computation within the C++ programming language*, [cs/0004015](#).
- [89] J. Vermaseren, *New features of FORM*, [math-ph/0010025](#) [[INSPIRE](#)].
- [90] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer and T. Stelzer, *MadGraph 5 : Going Beyond*, *JHEP* **06** (2011) 128 [[arXiv:1106.0522](#)] [[INSPIRE](#)].
- [91] C. Anastasiou and A. Lazopoulos, *Automatic integral reduction for higher order perturbative calculations*, *JHEP* **07** (2004) 046 [[hep-ph/0404258](#)] [[INSPIRE](#)].

- [92] M.L. Mangano and S.J. Parke, *Multiparton amplitudes in gauge theories*, *Phys. Rept.* **200** (1991) 301 [[hep-th/0509223](#)] [[INSPIRE](#)].
- [93] T. Huber and D. Maître, *HypExp: A Mathematica package for expanding hypergeometric functions around integer-valued parameters*, *Comput. Phys. Commun.* **175** (2006) 122 [[hep-ph/0507094](#)] [[INSPIRE](#)].
- [94] M. Czakon, *Automatized analytic continuation of Mellin-Barnes integrals*, *Comput. Phys. Commun.* **175** (2006) 559 [[hep-ph/0511200](#)] [[INSPIRE](#)].
- [95] A. Smirnov and V. Smirnov, *On the Resolution of Singularities of Multiple Mellin-Barnes Integrals*, *Eur. Phys. J. C* **62** (2009) 445 [[arXiv:0901.0386](#)] [[INSPIRE](#)].
- [96] D.A. Kosower, *barnesroutines*, <http://projects.hepforge.org/mbtools/>.
- [97] J. Vermaseren, *Harmonic sums, Mellin transforms and integrals*, *Int. J. Mod. Phys. A* **14** (1999) 2037 [[hep-ph/9806280](#)] [[INSPIRE](#)].
- [98] S. Moch, P. Uwer and S. Weinzierl, *Nested sums, expansion of transcendental functions and multiscale multiloop integrals*, *J. Math. Phys.* **43** (2002) 3363 [[hep-ph/0110083](#)] [[INSPIRE](#)].
- [99] S. Moch and P. Uwer, *XSummer: Transcendental functions and symbolic summation in form*, *Comput. Phys. Commun.* **174** (2006) 759 [[math-ph/0508008](#)] [[INSPIRE](#)].
- [100] F. Brown, *The Massless higher-loop two-point function*, *Commun. Math. Phys.* **287** (2009) 925 [[arXiv:0804.1660](#)] [[INSPIRE](#)].
- [101] A.B. Goncharov, *Multiple polylogarithms, cyclotomy and modular complexes*, *Math. Res. Lett.* **5** (1998) 497 [[arXiv:1105.2076](#)] [[INSPIRE](#)].
- [102] A.B. Goncharov, *Multiple polylogarithms and mixed Tate motives*, [math/0103059](#).
- [103] C.F. Berger, V. Del Duca and L.J. Dixon, *Recursive Construction of Higgs-Plus-Multiparton Loop Amplitudes: The Last of the Phi-nite Loop Amplitudes*, *Phys. Rev. D* **74** (2006) 094021 [Erratum *ibid.* **D 76** (2007) 099901] [[hep-ph/0608180](#)] [[INSPIRE](#)].
- [104] S. Badger, E.N. Glover and K. Risager, *One-loop phi-MHV amplitudes using the unitarity bootstrap*, *JHEP* **07** (2007) 066 [[arXiv:0704.3914](#)] [[INSPIRE](#)].
- [105] E.N. Glover, P. Mastrolia and C. Williams, *One-loop phi-MHV amplitudes using the unitarity bootstrap: The General helicity case*, *JHEP* **08** (2008) 017 [[arXiv:0804.4149](#)] [[INSPIRE](#)].
- [106] S. Badger and E.N. Glover, *One-loop helicity amplitudes for $H \rightarrow j$ gluons: The All-minus configuration*, *Nucl. Phys. Proc. Suppl.* **160** (2006) 71 [[hep-ph/0607139](#)] [[INSPIRE](#)].
- [107] L.J. Dixon and Y. Sofianatos, *Analytic one-loop amplitudes for a Higgs boson plus four partons*, *JHEP* **08** (2009) 058 [[arXiv:0906.0008](#)] [[INSPIRE](#)].
- [108] S. Badger, J.M. Campbell, R.K. Ellis and C. Williams, *Analytic results for the one-loop NMHV Hqgg amplitude*, *JHEP* **12** (2009) 035 [[arXiv:0910.4481](#)] [[INSPIRE](#)].
- [109] S. Badger, E. Nigel Glover, P. Mastrolia and C. Williams, *One-loop Higgs plus four gluon amplitudes: Full analytic results*, *JHEP* **01** (2010) 036 [[arXiv:0909.4475](#)] [[INSPIRE](#)].
- [110] T. Gehrmann, M. Jaquier, E. Glover and A. Koukoutsakis, *Two-Loop QCD Corrections to the Helicity Amplitudes for $H \rightarrow 3$ partons*, *JHEP* **02** (2012) 056 [[arXiv:1112.3554](#)] [[INSPIRE](#)].

- [111] P. Baikov, K. Chetyrkin, A. Smirnov, V. Smirnov and M. Steinhauser, *Quark and gluon form factors to three loops*, *Phys. Rev. Lett.* **102** (2009) 212002 [[arXiv:0902.3519](#)] [[INSPIRE](#)].
- [112] R. Lee, A. Smirnov and V. Smirnov, *Analytic Results for Massless Three-Loop Form Factors*, *JHEP* **04** (2010) 020 [[arXiv:1001.2887](#)] [[INSPIRE](#)].
- [113] T. Gehrmann, E. Glover, T. Huber, N. Ikizlerli and C. Studerus, *Calculation of the quark and gluon form factors to three loops in QCD*, *JHEP* **06** (2010) 094 [[arXiv:1004.3653](#)] [[INSPIRE](#)].
- [114] M. Beneke and V.A. Smirnov, *Asymptotic expansion of Feynman integrals near threshold*, *Nucl. Phys. B* **522** (1998) 321 [[hep-ph/9711391](#)] [[INSPIRE](#)].
- [115] V.A. Smirnov, *Applied asymptotic expansions in momenta and masses*, *Springer Tracts Mod. Phys.* **177** (2002) 1.
- [116] M. Hschele, J. Hoff, A. Pak, M. Steinhauser and T. Ueda, *Higgs boson production at the LHC: NNLO partonic cross sections through order ϵ and convolutions with splitting functions to N^3LO* , *Phys. Lett. B* **721** (2013) 244 [[arXiv:1211.6559](#)] [[INSPIRE](#)].
- [117] S. Buehler and A. Lazopoulos, *Scale dependence and collinear subtraction terms for Higgs production in gluon fusion at N^3LO* , [arXiv:1306.2223](#) [[INSPIRE](#)].
- [118] V. Del Duca, C. Duhr, E. Nigel Glover and V.A. Smirnov, *The One-loop pentagon to higher orders in epsilon*, *JHEP* **01** (2010) 042 [[arXiv:0905.0097](#)] [[INSPIRE](#)].
- [119] V. Del Duca, C. Duhr and V.A. Smirnov, *The Two-Loop Hexagon Wilson Loop in $N = 4$ SYM*, *JHEP* **05** (2010) 084 [[arXiv:1003.1702](#)] [[INSPIRE](#)].
- [120] J. Ablinger, J. Blumlein, A. Hasselhuhn, S. Klein, C. Schneider and F. Wissbrock, *Massive 3-loop Ladder Diagrams for Quarkonic Local Operator Matrix Elements*, *Nucl. Phys. B* **864** (2012) 52 [[arXiv:1206.2252](#)] [[INSPIRE](#)].
- [121] F. Chavez and C. Duhr, *Three-mass triangle integrals and single-valued polylogarithms*, *JHEP* **11** (2012) 114 [[arXiv:1209.2722](#)] [[INSPIRE](#)].
- [122] C. Bogner and F. Brown, *Symbolic integration and multiple polylogarithms*, *PoS LL2012* (2012) 053 [[arXiv:1209.6524](#)] [[INSPIRE](#)].
- [123] G. A.B., *A simple construction of Grassmannian polylogarithms*, [arXiv:0908.2238](#) [[INSPIRE](#)].
- [124] K.T. Chen, *Iterated path integrals*, *Bull. Amer. Math. Soc.* **83** (1977) 831.
- [125] F.C. Brown, *Multiple zeta values and periods of moduli spaces $M_{0,n}(R)$* , *Annales Sci. Ecole Norm. Sup.* **42** (2009) 371 [[math/0606419](#)] [[INSPIRE](#)].
- [126] A.B. Goncharov, M. Spradlin, C. Vergu and A. Volovich, *Classical Polylogarithms for Amplitudes and Wilson Loops*, *Phys. Rev. Lett.* **105** (2010) 151605 [[arXiv:1006.5703](#)] [[INSPIRE](#)].
- [127] C. Duhr, H. Gangl and J.R. Rhodes, *From polygons and symbols to polylogarithmic functions*, *JHEP* **10** (2012) 075 [[arXiv:1110.0458](#)] [[INSPIRE](#)].
- [128] C. Duhr, *Hopf algebras, coproducts and symbols: an application to Higgs boson amplitudes*, *JHEP* **08** (2012) 043 [[arXiv:1203.0454](#)] [[INSPIRE](#)].
- [129] F. Brown, *On the decomposition of motivic multiple zeta values*, [arXiv:1102.1310](#) [[INSPIRE](#)].

- [130] T. Gehrmann and E. Remiddi, *Numerical evaluation of harmonic polylogarithms*, *Comput. Phys. Commun.* **141** (2001) 296 [[hep-ph/0107173](#)] [[INSPIRE](#)].
- [131] T. Gehrmann and E. Remiddi, *Numerical evaluation of two-dimensional harmonic polylogarithms*, *Comput. Phys. Commun.* **144** (2002) 200 [[hep-ph/0111255](#)] [[INSPIRE](#)].
- [132] D. Maître, *HPL, a mathematica implementation of the harmonic polylogarithms*, *Comput. Phys. Commun.* **174** (2006) 222 [[hep-ph/0507152](#)] [[INSPIRE](#)].
- [133] D. Maître, *Extension of HPL to complex arguments*, *Comput. Phys. Commun.* **183** (2012) 846 [[hep-ph/0703052](#)] [[INSPIRE](#)].
- [134] J. Vollinga and S. Weinzierl, *Numerical evaluation of multiple polylogarithms*, *Comput. Phys. Commun.* **167** (2005) 177 [[hep-ph/0410259](#)] [[INSPIRE](#)].
- [135] S. Buehler and C. Duhr, *CHAPLIN - Complex Harmonic Polylogarithms in Fortran*, [arXiv:1106.5739](#) [[INSPIRE](#)].
- [136] H.R.P. Ferguson, D.H. Bailey and S. Arno, *Analysis of PSLQ, an integer relation finding algorithm*, *Math. Comput.* **68** (1999) 351 [*NASA Ames Technical Report*, NAS 96 005].