

# The full greenhouse gases budget of Africa: Synthesis, uncertainties and vulnerabilities

## Synthesis, uncertainties and vulnerabilities

### Working Paper

**Author(s):**

Valentini, Riccardo; Arneth, Almuth; Bombelli, Antonio; Castaldi, Simona; Cazzolla Gatti, Roberto; Chevallier, Frédéric; Ciais, Philippe; Grieco, Elisa; Hartmann, Jens; Henry, Matieu; Houghton, R.A.; Jung, Martin; Kutsch, Werner L.; Malhi, Yadvinder Singh; Mayorga, Emilio; Merbold, Lutz; Papale, Dario; Peylin, Philippe; Raymond, Peter A.; Santini, Monia; Sitch, Stephen A.; Vaglio Laurin, Gaia; van der Werf, Guido R.; Williams, Christopher R.; Scholes, Robert J.

**Publication date:**

2013-05-17

**Permanent link:**

<https://doi.org/10.3929/ethz-b-000077875>

**Rights / license:**

[Creative Commons Attribution 3.0 Unported](#)

**Originally published in:**

Biogeosciences Discussions 10, <https://doi.org/10.5194/bgd-10-8343-2013>



This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# The full greenhouse gases budget of Africa: synthesis, uncertainties and vulnerabilities

R. Valentini<sup>1,2</sup>, A. Arneth<sup>3</sup>, A. Bombelli<sup>2</sup>, S. Castaldi<sup>2,4</sup>, R. Cazzolla Gatti<sup>1</sup>,  
F. Chevallier<sup>5</sup>, P. Ciais<sup>5</sup>, E. Grieco<sup>2</sup>, J. Hartmann<sup>6</sup>, M. Henry<sup>7</sup>, R. A. Houghton<sup>8</sup>,  
M. Jung<sup>9</sup>, W. L. Kutsch<sup>10</sup>, Y. Malhi<sup>11</sup>, E. Mayorga<sup>12</sup>, L. Merbold<sup>13</sup>,  
G. Murray-Tortarolo<sup>15</sup>, D. Papale<sup>1</sup>, P. Peylin<sup>5</sup>, B. Poulter<sup>5</sup>, P. A. Raymond<sup>14</sup>,  
M. Santini<sup>2</sup>, S. Sitch<sup>15</sup>, G. Vaglio Laurin<sup>2,16</sup>, G. R. van der Werf<sup>17</sup>,  
C. A. Williams<sup>18</sup>, and R. J. Scholes<sup>19</sup>

<sup>1</sup>Department for Innovation in Biological, Agro-food and Forest systems (DIBAF), University of Tuscia, via S. Camillo de Lellis, 01100 Viterbo, Italy

<sup>2</sup>Euro-Mediterranean Center on Climate Change (CMCC), Via Augusto Imperatore 16, 73100 Lecce, Italy

<sup>3</sup>Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research, Atmospheric Environmental Research, Kreuzeckbahn Str. 19, 82467 Garmisch-Partenkirchen, Germany

<sup>4</sup>Dipartimento di Scienze Ambientali, Biologiche e Farmaceutiche (DISTABIF), Seconda Università di Napoli, via Vivaldi 43, 81100 Caserta, Italy

<sup>5</sup>LSCE, CEA-CNRS-UVSQ, L'Orme des Merisiers, Bat 701, 91191 Gif-sur-Yvette, France

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



<sup>6</sup>Institute for Biogeochemistry and Marine Chemistry, 20146, Hamburg, Germany

<sup>7</sup>FAO, Forestry Department, UN-REDD Programme, Viale delle terme di Caracalla 1, 00153 Rome, Italy

<sup>8</sup>Woods Hole Research Center, 149 Woods Hole Road, Falmouth, MA 02540, USA

<sup>9</sup>Biogeochemical Model Data Integration Group, Max Planck Institute for Biogeochemistry, Hans-Knöll Str. 10, 07745 Jena, Germany

<sup>10</sup>Thuenen Institute for Climate-smart Agriculture, Bundesallee 50, 38116 Braunschweig, Germany

<sup>11</sup>Environmental Change Institute, School of Geography and the Environment, University of Oxford, UK

<sup>12</sup>Applied Physics Laboratory, University of Washington, Seattle, USA

<sup>13</sup>Department of Environmental Systems Science, Institute of Agricultural Sciences IAS, ETH Zurich, 8092 Zurich, Switzerland

<sup>14</sup>Yale School of Forestry and Environmental Studies, 195 Prospect St, new Haven CT, 06511, USA

<sup>15</sup>University of Exeter, Rennes Drive, Exeter EX4 4RJ, UK

<sup>16</sup>Department of Civil Engineering and Computer Science Engineering, University of Tor Vergata, Rome, Italy

<sup>17</sup>Faculty of Earth and Life Sciences, VU University Amsterdam, De Boelelaan 1085, 1081HV Amsterdam, the Netherlands

<sup>18</sup>Graduate School of Geography, Clark University, Worcester, Massachusetts, USA

<sup>19</sup>Council for Scientific and Industrial Research, P.O. Box 395, Pretoria 0001 South Africa

Received: 23 January 2013 – Accepted: 26 March 2013 – Published: 17 May 2013

Correspondence to: R. Valentini (rik@unitus.it)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Abstract

This paper, developed under the framework of the RECCAP initiative, aims at providing improved estimates of the carbon and GHG (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) balance of continental Africa. The various components and processes of the African carbon and GHG budget were considered, and new and available data derived by different methodologies (based on inventories, ecosystem fluxes, models, and atmospheric inversions) were integrated. The related uncertainties were quantified and current gaps and weakness in knowledge and in the monitoring systems were also considered in order to provide indications on the future requirements. The vast majority of the results seem to agree that Africa is probably a small sink of carbon on an annual scale, with an average value of  $-0.61 \pm 0.58 \text{ PgCyr}^{-1}$ . Nevertheless the emissions of CH<sub>4</sub> and N<sub>2</sub>O may turn Africa into a source in terms of CO<sub>2</sub> equivalents. At sub-regional level there is a significant spatial variability in both sources and sinks, mainly due to the biome's differences and the different anthropic impacts, with southern Africa as the main source and central Africa, with its evergreen tropical forests, as the main sink. Emissions from land use change in Africa are significant (around  $0.32 \pm 0.05 \text{ PgCyr}^{-1}$ ) and even higher than the fossil fuel ones; this is a unique feature among all the continents. In addition there can be significant carbon losses from land even without changes in the land use (forest), as results from the impact of selective logging. Fires also play a significant role, with  $1.03 \pm 0.22 \text{ PgCyr}^{-1}$  of carbon emissions, mainly (90 %) originated by savanna and woodland burning. But whether fire carbon emissions are compensated by CO<sub>2</sub> uptake during the growing season, or are a non-reversible loss of CO<sub>2</sub>, remains unclear. Most of these figures are subjected to a significant interannual variability, on the order of  $\pm 0.5 \text{ PgCyr}^{-1}$  in standard deviation, accounting for around 25 % of the year-to-year variation in the global carbon budget.

These results, even if still highly uncertain, show the important role that Africa plays in the carbon cycle at global level, both in terms of absolute values and variability.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# 1 Introduction

Africa has the highest population annual growth rate in the world, at 2.3 % in 2008 (UN, 2010). The population of Africa exceeded one billion inhabitants in 2009 (UN, 2011). At the same time Africa has the lowest GDP of any continent and its economies rely mostly on natural resources, especially small scale agriculture: 64 % of people in Sub-Saharan Africa (SSA) are employed in the rural sector and an estimated 90 % of the continent's population depends on rain-fed crop production and pastoralism to meet its basic food supplies (IFAD, 2001; Patt and Winkler, 2007; UNSTAT, 2010). Population growth scenarios and increasing energy demand, the intensive exploitation of natural resources for food and wood products, and the associated land uses emissions make the African regional carbon budget and the processes driving GHG emission and up-take a particularly urgent topic for investigation.

Independent estimates of the African contribution to the global carbon cycle have recently become available (Williams et al., 2007; Bombelli et al., 2009; Ciais et al., 2011). These studies combined various sources of data and various methodologies, from bottom-up ecosystem inventories to land fluxes, biogeochemical models and atmospheric inversions. The results indicate a net biospheric sink of carbon, however there is still quite a significant range across the different estimates, from a near zero value to a net sink of  $1.0 \text{ PgCyr}^{-1}$ , with high uncertainties associated<sup>1</sup>.

This paper, developed under the framework of the REgional Carbon Cycle Assessment and Processes (RECCAP, <http://www.globalcarbonproject.org/reccap>) initiative of the Global Carbon Project (GCP) (Canadell et al., 2011), aims at providing an improved estimate of the carbon balance of continental Africa, along with balances for non-CO<sub>2</sub> greenhouse gases, through the integration of data from different sources and methodologies with the related quantification of the uncertainties. In particular, a comprehensive review of available information together with newly produced data, from in

<sup>1</sup>“C” indicates carbon from CO<sub>2</sub> only; when CH<sub>4</sub> and/or other non CO<sub>2</sub> GHG are included CO<sub>2</sub>-eq is used.

BGD

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



situ, space based and data assimilation systems, was conducted. Current gaps and weakness in knowledge and in the monitoring systems are also considered in order to provide indications on the future requirements. In particular this paper will try (1) to disentangle the various components and processes of the carbon budget and the potentials of today methodologies to address them and (2) to extend the scope of the previous studies by including a more comprehensive analysis of other greenhouse gases, specifically methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O).

## 2 Africa's GHG emissions and removals as reported to UNFCCC

We have estimated the Greenhouse Gases (GHG) emissions on the basis of the national communication to the United Nations Framework Convention on Climate Change (UNFCCC) updated to December 2011<sup>2</sup>. The results reveal a profile of emissions from the African continent that is remarkably different to that of other regions in the world. Firstly, the contribution of fossil fuel emissions is low (less than 4 % of total global CO<sub>2</sub> emissions; Canadell et al., 2009a) but increasing rapidly. In terms of per capita emissions from fossil fuel based energy consumption, Southern Africa, led by South Africa, has the highest per capita emission rate, at 8.5 MgCO<sub>2</sub>-eq person<sup>-1</sup> yr<sup>-1</sup>, i.e. 4 to 20 times the per capita emissions of other African regions.

<sup>2</sup>Data were processed from the latest (updated to December 2011) African countries national communication to the UNFCCC. Emission sources analysis has been developed considering the following sectors: anthropogenic emissions (regrouping emissions from the energy, industry and waste sectors); agriculture emissions; Land Use Change and Forestry (LUCF) emissions; and removals (mainly by forests). The emissions rates include, when available, CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O; all the values were converted to CO<sub>2</sub> equivalent (CO<sub>2</sub>-eq) on the basis of Global Warming Potential reported in IPCC (2007). The countries Libya, Mayotte, Reunion, Seychelles, Somalia, Zambia, Guinea Bissau, Liberia, Central African Republic and Equatorial Guinea were not considered in this analysis due to the lack or the inconsistency of the data reported in their National Communication.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏮

⏭

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The inventory data suggest that fossil fuel based emissions from Africa are almost equal to the emissions from Land Use Change and Forestry (LUCF), the latter including deforestation. Agriculture also plays an important role, amounting to 26 % of total emissions (including  $\text{N}_2\text{O}$  and  $\text{CH}_4$ ). It is important to underscore that LUCF emissions are more than one third (35 %) of total emissions in Africa, a share significantly higher than the 9 % at the global level (GCP, 2011). These emissions are partially counterbalanced by removals from the atmosphere of about  $2.20 \text{ PgCO}_2\text{-eq}$ , mainly due to the forest sector of tropical Africa, resulting in a net source of  $0.95 \text{ PgCO}_2\text{-eq}$ . Comparing these data for the year 2011 with the study of Bombelli et al. (2009) there is clear evidence of an increasing trend in both fossil fuel and agricultural emissions, while LUCF sector shows a decrease of about  $0.38 \text{ PgCO}_2\text{-eq}$  compared with the 1994 reference year. This, combined with the simultaneous reduction in removals caused by a decreasing sink, results in increased emissions from  $0.36$  to  $0.95 \text{ PgCO}_2\text{-eq}$  (Table 1) if compared to the 1994 reference year.

A sub-regional analysis shows that Central Africa plays a key role as carbon sink, but at the same time contributes the greatest emissions from the LUCF sector. This could be a sign of an increasing threat to local forest resources. Southern Africa (mainly due to South Africa), followed by North Africa contribute more than other regions to the African fossil fuel emissions – with Southern and North Africa being responsible for almost the whole component of the total emissions. In Western and Eastern Africa agricultural and LUCF emissions dominate.

Therefore, according to national reports to the UNFCCC and considering  $\text{CO}_2$  equivalents, Africa is a net source of  $0.95 \text{ PgCO}_2\text{-eq}$  per year, while focusing only on the carbon contribution of LUCF and removals by forests, the net carbon budget of the African continent is a sink of  $-0.31 \text{ PgCyr}^{-1}$ , that is a small carbon sink on a whole continent on a year basis. However it must be noted that this estimate is the result of the difference between two large and uncertain numbers.

### 3 The African carbon budget and its components

Despite the availability of carbon budget estimates overall, there are many individual processes that are relevant for understanding the global carbon cycle. Indeed, 17 % of global carbon emissions from land-use change, over half of the global gross fire emissions and about half of the interannual variability of the global carbon balance have been attributed to Africa (Williams et al., 2008; Canadell et al., 2009b; van der Werf et al., 2010). About 1/3 of the tropical biomass carbon sink (16 % of the total terrestrial carbon sink) over the period 2000–2005 is thought to be due to the African tropical forests (Lewis et al., 2009; Malhi, 2010). However, these estimates carry large uncertainties, particularly regarding the role of Africa as carbon source or sink; the role of forest degradation and deforestation; and the inter-annual variability of the carbon fluxes. A complete and accurate greenhouse gas account for Africa is not yet available. In the following sections we disentangle the various processes and their contribution to the African GHG budget.

#### 3.1 Africa land cover: estimates and differences between various studies

Estimation of land cover area and its change is critical for deriving bottom-up estimates from models, inventories and field data. This is particularly true when the land cover area is used to scale biomass estimates or fluxes of greenhouse gases from local to continental level. For this reason we try to quantify here this source of uncertainty by comparing different land cover maps for Africa.

Several attempts have been made during the past few years to standardize land cover maps globally using a common reference classification system (the FAO/UNEP Land Cover Classification System – LCCS, [http://www.glcn.org/ont.2\\_en.jsp](http://www.glcn.org/ont.2_en.jsp)). Despite the common classification approach it remains difficult to compare different land cover products. We considered three widely used global land cover products (GCL2000, GlobCover2006 and GlobCover2009) and tried to regroup the large number of classes into a smaller number: croplands, forests, shrublands/grasslands (including savannas),

BGD

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



flooded vegetation, bare soil, water bodies and other. The aggregated accuracy of the land cover products is 66 %. However there are still remarkable differences in the areas reported, in particular for forests (Table 2). Forested area is reported for Africa to be between 638.15 and 836.81 Mha – the range of variation is about 27 % of the median value. According to FAO (FRA, 2010) deforestation in the 2000–2010 period is estimated to be about 3.4 Mha yr<sup>-1</sup>. Thus deforestation cannot by itself explain such big differences. As a consequence of the unexplained differences in the forest areas, these land cover products cannot be used for deriving deforestation rates by simple differencing. A possible source of disagreement between the different land cover products is the fact that they were not produced for the same reference year. Other possible reasons are the differences of the definitions within a land cover class (e.g. forests can include open forest, closed forest, sparse forest, woodlands, savannas, etc.), the different land cover classes used by the different products, and the presence of mixed classes (i.e. mosaics of forests, savannas, croplands, etc.) without the proportions of the different components.

Such uncertainty in the area estimation of land cover classes can create large biases in estimating net fluxes and/or biomass changes, particularly in forests. In the other categories the variation in estimated areas is more limited. For croplands the range is 6.4 % of the median, and for shrublands/grasslands it is 14.8 %.

### 3.2 Biomass estimation

Considering that around 16–17 % of the world tropical forested area occurs in Africa (FAO, 2006), it is surprising how few carbon stocks estimations at continental level have been published. Brown and Gaston (1995) produced one of the first estimates using forest inventory data combined with land-use modeling. Others based their analysis on remote sensing data and field derived ground truth (Baccini et al., 2008, 2012; Saatchi et al., 2011). Lewis et al. (2009) analyzed a network of research plots across Africa to estimate long term changes in carbon stocks. At local level aboveground biomass has been assessed in a number of specific forest sites (i.e. Henry, 2010; Glenday, 2008;



For these aggregations, basic biomass statistics (Table 4) and histograms (Fig. 1) were computed to evaluate distribution patterns.

The Croplands aggregation is characterized by medium-low biomass values, with areas having higher biomass occupying about 20 % of the extent. The higher biomass cells are found in compact clusters inside a highly human-influenced mosaic: they mostly belong to class 30 (Table 3) which includes also forest patches. These areas have a role in C stocks changes and can be prioritized for forest recovery activities. The Forests aggregation is characterized by broad geographical distribution. The histogram (Fig. 1) has two peaks in a nearly bimodal distribution corresponding to two distinct areas of low and high biomass ranges. In this aggregation, cells belonging to class 40 (lowland tropical rainforest, Table 3) account for most of the areas of higher biomass. Their spatial distribution can be resumed in two large intact blocks in eastern and western Central Africa; a smaller area in West Africa at the Liberia-Sierra Leone border; a smaller fraction with lower biomass around the above blocks and close to the coast, in highly-anthropogenic areas. The shrublands/grasslands aggregation has lowest biomass values and corresponds mainly to drier environments. The Flooded vegetation aggregation includes areas of very high biomass, but its extent is relatively small. It occurs in clusters in the western Democratic Republic of Congo and Congo Republic. This block deserves to be prioritized for conservation activities.

We recognize that the GlobCover 2009 map has known issues in identifying flooded areas, while the WHRC biomass map has been built with limited ground truth. Furthermore, they do not cover non-tropical Africa. Nonetheless, despite these limitations, they are the freely accessible most updated products available so far.

According to this analysis (Table 4) the total tropical African aboveground ground biomass is 77.9 PgC, with different coefficients of variation according to the different land cover classes (Table 3). Considering that each dataset is possibly affected by a 10–15 % representation error, the combined error associated with the overall estimate of biomass in tropical Africa is in the order of 19–27 %.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.3 Atmospheric inversions

Inversions used in this study are Bayesian synthesis inversions where a prior set of surface fluxes is modified to match atmospheric CO<sub>2</sub> observations, given a numerical model of atmospheric transport and a statistical description of the uncertainties of both the prior fluxes and the observations. Over Africa, inversions are poorly constrained by measurements from the surface CO<sub>2</sub> network. The main stations of the surface network are located in South Africa (station CPT), Namibia (station NMB), Algeria (station ASK), Ivory Coast (station LTO) and in two islands downwind of the continent (stations IZO, in the Canary Islands, and ASC, in Ascension Island). Furthermore, the importance of vertical mixing by convection during the wet season associated with the seasonal displacement of the ITCZ over Africa makes the simulation of CO<sub>2</sub> transport particularly challenging, and reduces the influence of surface fluxes on surface station CO<sub>2</sub> variations in atmospheric transport model. As a consequence, inverted fluxes over Africa are subjected to large uncertainties and often reflect the prior fluxes used in the Bayesian inversion.

Early inversion estimates (from 5 inversions corresponding to the late-1990's period) compiled by Williams et al. (2007) suggested that Africa is close to CO<sub>2</sub> neutral. This would imply that land use CO<sub>2</sub> emissions (0.3 PgCyr<sup>-1</sup> in Friedlingstein et al., 2010) and fossil fuel CO<sub>2</sub> emissions (0.26 PgCyr<sup>-1</sup> in Canadell et al., 2009a) are approximately in balance with a residual CO<sub>2</sub> uptake by ecosystems processes not counted in the land use CO<sub>2</sub> emissions. The Net CO<sub>2</sub> flux estimates over Africa given by inversions published in the late 1990's and early 2000's are: 0.1 PgCyr<sup>-1</sup> during 1985–1995 (Bousquet et al., 1999), –0.1 PgCyr<sup>-1</sup> during 1996–1999 (Roedenbeck et al., 2003), –0.4 PgCyr<sup>-1</sup> and 0.25 PgCyr<sup>-1</sup> from the average of 13 inversions of the TRANSCOM project during 1992–1996, in Gurney et al. (2002) (based on annual mean CO<sub>2</sub> gradients) and in Gurney et al. (2003) (based on monthly mean CO<sub>2</sub> gradients), respectively. Positive numbers denote net sources of CO<sub>2</sub> to the atmosphere.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The results from five more recent inversions with fluxes “solved” at the transport model spatial resolution for four of them have been compiled (see display in <http://www.carboscope.eu/>, access 2 May 2011). Note that two of them correspond to a variant of the same system using different observation networks. Over Africa, inversion estimates of the CO<sub>2</sub> natural flux are 0.1 PgCyr<sup>-1</sup> during 1996–2008 using the inversion system of Roedenbeck et al. (2003), –0.26 PgCyr<sup>-1</sup> using the same system but over a shorter period (1999–2008) with 61 atmospheric stations versus 52 in the previous set up, 0.21 PgCyr<sup>-1</sup> during 2000–2007 using the CarbonTracker-Europe inversion of Peters et al. (2009), 0.32 PgCyr<sup>-1</sup> during 1996–2004 using the results described in Piao et al. (2009) (following the analytical formulation of Peylin et al., 2005), and –0.08 PgCyr<sup>-1</sup> during 1988–2008, according to the inversion of Chevallier et al. (2010). The inversion fluxes (including fossil fuel emissions) aggregated over Africa are showed in Fig. 2. The longest inversion (LSCE\_var.v1.0) gives over the period 1988–2008 a mean CO<sub>2</sub> flux of  $0.24 \pm 0.8$  PgCyr<sup>-1</sup> over northern Africa and a sink of  $-0.15 \pm 0.9$  PgCyr<sup>-1</sup> over southern Africa (the uncertainty representing 1-sigma for a single year).

Despite the large uncertainty associated with the mean continental scale natural CO<sub>2</sub> flux, the results from the five inversions displayed in Fig. 2 show coherent interannual variations of CO<sub>2</sub> fluxes, though with different amplitudes. A strong source anomaly (positive land-to-atmosphere CO<sub>2</sub> flux) during 1997–1998 associated with the positive-phase ENSO (El Niño Southern Oscillation) condition is obtained for two systems, while during 2001–2002 most systems provide also a slightly positive flux without any significant phase of ENSO.

Figure 3a shows the geographic distribution of the mean land-to-atmosphere CO<sub>2</sub> flux from these inversions, while Fig. 3b shows the spread of the five inversions, defined from the estimator of the standard deviation. Note that the estimated spatial distribution is highly controlled by the a priori flux distribution, the location of imposed biomass burning (except for the JENA systems), and the flux error correlation length used by the LSCE and JENA systems. Inversions appear to be consistent in placing a large uptake

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



over central African forests ( $-60 \text{ gC m}^{-2} \text{ yr}^{-1}$ ), although with an associated uncertainty of nearly similar magnitude. This feature is not surprising because all inversions allocate large prior errors to tropical forest fluxes, and since the region is not constrained by atmospheric measurements, the models tend to place large sink increments in this biome. On the other hand, the spread of the five inversions is much larger over savannas (Fig. 3b), and larger than the mean flux (indicating positive or negative fluxes). This disagreement may originate from different choices for the prior biomass burning fluxes or the prior natural fluxes. It is seen in Fig. 3a that some inversions use non-zero a priori fluxes, in particular for fire emissions in savannas.

In conclusion, although inversions are associated with significant uncertainties over the African continent, both earlier inversions and recent ones robustly converge toward a neutral to slightly positive  $\text{CO}_2$  balance for Africa, but this results has a standard deviation larger than  $1 \text{ PgCyr}^{-1}$ . New estimates of the African  $\text{CO}_2$  flux will soon be obtained from the observations from the Greenhouse gases Observing SATellite (GOSAT, Kuze et al., 2010), launched in January 2009 by the JAXA (Japan Aerospace Exploration Agency). The GOSAT satellite observations of column  $\text{CO}_2$  data cover the atmosphere over tropical continents much better than in situ stations, albeit with larger errors on individual soundings. The error on the  $\text{CO}_2$  balance of Africa is expected to be strongly reduced when GOSAT data have been assimilated (Chevallier et al., 2009), provided that these data will have no systematic bias. The errors on annual  $\text{CO}_2$  fluxes are expected to decrease from  $\pm 1.3 \text{ PgCyr}^{-1}$  (6 % error reduction from the assumed prior error) when using in situ data from the current surface network, down to  $\pm 0.4 \text{ PgCyr}^{-1}$  (72 % error reduction from the prior error) expected from the assimilation of GOSAT data, if those data have no bias.

### 3.4 Terrestrial fluxes up-scaled from in situ observations

In recent years there has been an expansion of the flux tower network in Africa (Bombelli et al., 2009). The total number of flux sites (see FLUXNET database)

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



includes more than 15 sites with continuous carbon dioxide, water vapour and energy fluxes, covering different plant functional types (from tropical forest, to transition forest and savannas) in different climatic regions. Some flux towers have been used in sporadic or seasonal campaign mode. The latest overview of the flux towers being active during CarboAfrica has been given in Merbold et al. (2009) and Sjöström et al. (2013).

The net annual carbon budget (NEE) of savanna sites ranges from a slight source of 0.25 to a sink of  $-4.53 \text{ tCha}^{-1} \text{ yr}^{-1}$  (Bombelli et al., 2009) with a median value being a sink of  $-0.14 \text{ tCha}^{-1} \text{ yr}^{-1}$ . Recent data from East African tropical wetlands (Saunders et al., 2012) show a quite significant sink ( $-10 \text{ tCha}^{-1} \text{ yr}^{-1}$  and  $-7 \text{ tCha}^{-1} \text{ yr}^{-1}$ ) for papyrus and cocoyam plantations, respectively.

There are still large gaps in our knowledge of NEE derived from in situ measurements for many ecosystems and climate regions of Africa. A first attempt to scale up the limited flux data was provided by Jung et al. (2011). Data-driven estimates of mean Gross Primary production (GPP), Total Ecosystem Respiration (TER), and NEE for the period 1982–2008 based on the integration of FLUXNET, satellite and meteorological observations using the Model Tree Ensemble (MTE) approach (Jung et al., 2009) are  $24.3 \pm 2.9 \text{ PgCyr}^{-1}$ ,  $19.0 \pm 1.4 \text{ PgCyr}^{-1}$ , and  $-3.7 \pm 1.6 \text{ PgCyr}^{-1}$  respectively (Jung et al., 2011). Uncertainties refer to one standard deviation of the spread of the 25 members of the model tree ensemble, and reflect the uncertainty in the structure of the model trees. Uncertainties related to FLUXNET data and global forcing data sets are not included in this measure. The NEE estimates are not consistent with the difference of TER and GPP because NEE was up-scaled independently. The “La Thuille” FLUXNET data set (<http://fluxnet.ornl.gov>) was used to train the model trees and thus they include insufficient data representation from Africa. Hence, uncertainties due to extrapolation to African ecosystem are likely to be substantial. Jung et al. (2011) inferred a systematic underestimation of 5–10 % of their global TER value, likely due to a biased sampling with enhanced representation of young and regrowing forests in the La Thuille data base and an even larger bias when considering African ecosystems only. The uncertainties of the NEE value are very large, as inferred by cross-validation



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Jung et al., 2011), because determinants of mean NEE, such as site-history, management, and disturbances, could not be accounted for explicitly in the MTE approach. The apparent strong sink of  $-3.7 \text{ Pg C yr}^{-1}$  is most likely a large overestimation, given that this number does not capture disturbed ecosystems being sources of  $\text{CO}_2$ . A second data-driven approach based on Artificial Neural Networks (ANN, Papale and Valentini, 2003) has been also used and gave as result a median African NEE of  $-2.22 \text{ Pg C yr}^{-1}$  (5 % and 95 % percentile  $-3.35$  and  $-1.08$ , respectively). Median values reported for North Africa are  $0.31 \text{ Pg C yr}^{-1}$  (percentile range  $-1.02$ ,  $0.42$ ) and  $-1.92 \text{ Pg C yr}^{-1}$  for Sub-Saharan Africa (5 % and 95 % 10 percentile  $-2.43$ ,  $-1.38$ ). The ANN approach is affected by the same uncertainty sources reported for the MTE results.

It is striking that there are no published field studies of NPP in African tropical forests where the major components of NPP have been directly measured. Malhi et al. (2011) conducted a review of NPP and its allocation in global tropical forests. For sites reporting woody production and canopy litterfall (i.e. neglecting root productivity), they found 33 sites in the Americas, 21 in Asia, 12 in Hawaii, and none in Africa, despite Africa being the second largest rainforest continent. For sites that also reported fine root productivity (and thus gave an estimate of total productivity) they found 35 sites worldwide, and again none in Africa. For lowland sites across the humid tropics, NPP ranged between 9 and  $15 \text{ tCha}^{-1} \text{ yr}^{-1}$ , and it is likely that NPP in African humid tropical forests is also within this range. In Amazonia, more seasonal forests did not appear to have much lower NPP than dry forests, with low GPP compensated by a higher carbon use efficiency (NPP/GPP).

NPP values from savannas were compiled by Bombelli et al. (2009) and they range from  $3.3$  to  $16.2 \text{ tCha}^{-1} \text{ yr}^{-1}$ , with a mean value of  $6.9 \text{ tCha}^{-1} \text{ yr}^{-1}$ .

To date no estimates of Gross Primary Productivity (GPP) have been reported from African tropical forests sites (although estimates should be forthcoming from flux towers and intensive carbon cycling sites in the next 1–2 yr). Humid tropical forests in the Americas and Asia report GPP in the range  $30$ – $40 \text{ tCha}^{-1} \text{ yr}^{-1}$  (Malhi, 2012), with the mean towards the lower end of this range. Taking a lower end value of  $30 \text{ tCha}^{-1} \text{ yr}^{-1}$



and multiplying by the area of African humid forests (about  $2.33 \times 10^6 \text{ km}^2$  in the GLC dataset) suggests a GPP of  $\sim 7.0 \text{ PgCyr}^{-1}$  in the humid forests. The African dry forests and woodlands are likely to have lower GPP (on the order of  $20 \text{ tCha}^{-1} \text{ yr}^{-1}$ , but are twice as extensive ( $4.28 \times 10^6 \text{ km}^2$ ) suggesting they contribute a further  $8.6 \text{ PgCyr}^{-1}$ . Thus the total GPP of African forests and woodlands could be around  $15.6 \text{ PgCyr}^{-1}$ , about 13 % of global terrestrial GPP.

Field studies in Amazonia suggest that the carbon use efficiency (NPP/GPP) is around 0.35 in humid forests, and rises to 0.45 in dry forests (Malhi, unpublished analysis). Applying these values to the humid and dry forest areas as defined in this paper (see Sect. 3.2) it results a total NPP of  $2.4 \text{ PgCyr}^{-1}$  in the humid forest zone and  $3.9 \text{ PgCyr}^{-1}$  in the dry forest/woodland zone, i.e.  $6.3 \text{ PgCyr}^{-1}$ , about 10 % of global terrestrial NPP, for the total forested zone.

The net change in biomass of tropical forests has been better reported, thanks to the pioneering work of the AfriTRON forest plot network (<http://www.geog.leeds.ac.uk/projects/afritron>). Lewis et al. (2009) collected data from repeated surveys of 79 forest plots in Central and West Africa, and reported a net biomass carbon sink of 0.63 (range 0.22–0.94)  $\text{tCha}^{-1} \text{ yr}^{-1}$ . This is very similar to the value of 0.45 (range 0.33–0.56)  $\text{tCha}^{-1} \text{ yr}^{-1}$  reported for 123 plots in Amazonia (Phillips et al., 2009), suggesting that a common driver, such as atmospheric  $\text{CO}_2$  fertilization, may be stimulating this biomass storage across the intact tropical forest biome. Scaling the data of Lewis et al. (2009) across the humid and dry forest biome, Malhi (2010) estimated a net biomass carbon sink of  $0.31 \pm 0.15 \text{ PgCyr}^{-1}$  in the African forest biome, i.e. 33 % of the global tropical forest carbon sink and 16 % of the global terrestrial carbon sink. This estimate does not incorporate any source or sink in the soils, and does not include the savanna biomes.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.5 DGVM based estimates of gross fluxes and net ecosystem productivity

Dynamic Global Vegetation Models (DGVMs) simulate vegetation distribution, seasonally changing phenology, and vegetation and soil carbon and water pools and fluxes. Vegetation is classified into plant functional types, based on today's observed bioclimatic limits for establishment and survival, and characteristic plant growth strategies. Leaf carbon assimilation is mostly modelled following Farquhar et al. (1980) and Collatz et al. (1991, 1992), or some modification thereof, and coupled to the plant and soil water balance via stomatal conductance. Vegetation and ecosystem net productivity are determined by carbon losses through autotrophic and heterotrophic respiration. A set of allocation rules determines the carbon allocated to canopy growth, as well as to litter input into soils (for details see: Arneth et al., 2010b; Cramer et al., 1999; Prentice et al., 2007; Sitch et al., 2008).

Modelling tree-grass coexistence is important to correctly simulate the gradient from closed forests, through woodlands and savannas to grasslands, and has been shown to be a challenge for many DGVMs (Cramer et al., 1999; Scheiter and Higgins, 2009). Whether this is due to the fact that many DGVMs have no or only a highly simplified representation of fire and other episodic events, or whether it reflects the need for a more generally improved representation of resource competition for light, nutrients and water, or both, is unresolved. Meta-analyses of African savannas, as well as fire-exclusion experiments have demonstrated that across the continent's mesic climate regions forests could emerge but for the existence of episodic disturbance events that reduce tree cover below its climatic maximum. The annual rainfall limits that have been determined to indicate climate support of forests differ, however, between analyses (Bond et al., 2005; Bucini and Hanan, 2007; Sankaran et al., 2005; Staver et al., 2011). Bond et al. (2005) and Lehsten et al. (2011) demonstrated a largely improved agreement between estimates of African vegetation cover and DGVM output when fire disturbance was included in the simulations, especially in regions with intermediate amounts of annual precipitation. In both cases inclusion of fire reduced the simulated

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



extent of the tropical forest biome considerably and grasses and shrubs became more competitive in areas that are climatically suitable for dense tree cover. A statistical generalized addition model, based on remotely sensed information of fire frequency, tree cover and precipitation yielded a similar conclusion (Staver et al., 2011).

A number of regionally (Scheiter and Higgins, 2009) and globally (Hickler et al., 2008; Smith et al., 2001) applicable DGVMs have begun to incorporate gap-model dynamics (Purves and Pacala, 2008) with a much more explicit representation of plant growth dynamics than in traditional DGVMs, for example now even resolving individuals or age cohorts in canopies. These types of models allow an improved analysis of the ecological effects of fire that is based on fire effects on vegetation structure and subsequent re-growth (Arneth et al., 2010a; Scheiter and Higgins, 2009), and the interactions with climate and the carbon cycle in Africa. It is important to note that today's modelling and data-assimilation efforts do not exclude other disturbance (e.g. herbivory, human land use) or resource effects (nutrients, water) as additional drivers to determine the relative proportion of forests and savannas, but these are as yet difficult to assess on regional and continental scales.

Only very few DGVM modelling studies have explored the African carbon cycle and its response to climate, CO<sub>2</sub> and land use change drivers. This is true despite the fact that (1) savannas and tropical forests are highly productive ecosystems (Grace et al., 2006; Williams et al., 2007) and (2) African carbon cycle dynamics thus play a potentially significant role for the global terrestrial carbon uptake and its interannual variability. Likewise, only limited observations exist to evaluate the quality of model output either on local (e.g. eddy flux data) or continental (e.g. comparison with inversion results) level (Kutsch et al., 2008, 2011; Merbold et al., 2009; Veenendaal et al., 2004; Weber et al., 2009; Williams et al., 2007; Ziehn et al., 2010).

Applying the model ORCHIDEE with dynamic vegetation disabled, Ciais et al. (2009) simulated a source-sink shift in the continental C balance from +0.14 PgCyr<sup>-1</sup> in the 1980s to -0.13 PgCyr<sup>-1</sup> in the 1990s, and interpreted this as the model's response to increasing CO<sub>2</sub> concentrations, increasing precipitation in some regions

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





period 1990–2009, although both the sign and magnitude of the flux varies among DGVMs. DGVMs simulate GPP, NPP and NBP over the period 1990–2010 of  $28.16 \pm 5.94 \text{ PgCyr}^{-1}$ ,  $13.27 \pm 3.25 \text{ PgCyr}^{-1}$  and  $0.41 \pm 0.31 \text{ PgCyr}^{-1}$ , respectively.

### 3.6 Fluxes resulting from land-use change (mainly deforestation)

In this section we provide a new estimate of carbon emission from land use changes in Africa. These are mainly deforestation, although we attempt to estimate also forest degradation and the role of agro-forestry systems. Annual emissions of carbon from land use (e.g. timber harvest) and land-use change (e.g. conversion of forest to cropland) were determined as described in Houghton and Hackler (2006) except that the calculations were updated here with the most recent FAO data on changes in forest area (FAO, 2010). Briefly, the analysis considered four sub-regions within sub-Saharan Africa: Western, Eastern, Central, and Southern Africa. Land uses included permanent croplands (cultivated lands), pastures, tree plantations, industrial wood harvest, and shifting cultivation (a rotational land use with alternating cropping and fallow, in which trees regrow until being cleared again for the cropping period). Fuelwood harvest was not explicitly included in the analysis, but it was implicitly included, in part, because both shifting cultivation and the conversion of forests to croplands and pastures result in large emissions of carbon from burning, some of which take place via collection and burning of fuelwood. Deforestation rates between 1980 and 2010 (FAO, 2010) expanded either cropland or pasture areas based on data from FAOSTAT (2009). If net forest area decreased by more than the net expansion of croplands and pastures, the difference was assumed to be deforestation for shifting cultivation (Houghton and Hackler, 2006). Between 1900 and 1980 rates of land-use change were obtained from various historical accounts (Houghton and Hackler, 2006). A bookkeeping model tracked changes in the carbon density ( $\text{MgC ha}^{-1}$ ) of vegetation and soils in response to land use. We assumed that following cultivation, native soils lost 25 % of the organic carbon in the top meter. Unmanaged ecosystems neither lost nor accumulated carbon, and environmental conditions (i.e.  $\text{CO}_2$  concentrations and climate) were assumed not

to affect carbon dynamics. Forest areas and biomass densities at the beginning of the analysis (1850) were chosen so that the areas and average biomass densities in 2010 matched those given by FAO (2010).

Over the last 100 yr, forest areas declined as a result of conversion to agricultural lands, and carbon densities declined as a result of wood harvest (degradation). For all of Sub-Saharan Africa the net emissions of carbon from land use and land-cover change averaged  $320 \pm 50 \text{ TgCyr}^{-1}$  over the period 1990–2009. The rate since 1980 was nearly constant for the entire region, but emissions increased slightly in Eastern, Western, and Southern Africa and decreased in Central Africa (Table 6). The largest emissions were from Western Africa ( $107 \text{ TgCyr}^{-1}$ ) and the lowest were in Eastern and Central Africa ( $\sim 60 \text{ TgCyr}^{-1}$ ). The land uses contributing to these emissions (in descending order of magnitude) were expansion of croplands, shifting cultivation, pastures, and industrial wood harvest (Table 6). The establishment of tree plantations was not a significant sink in any region. Again, the major drivers varied among sub-regions. Expanding croplands were most important in Western Africa; shifting cultivation was most important in Central Africa; and industrial wood harvest was most important in Central and Southern Africa. The emissions of carbon from land use and land-cover change in Sub-Saharan Africa over the period 1990–2009 account for about a third of global emissions from land use (Friedlingstein et al., 2010). An independent study of shifting cultivation supports the importance of Central Africa (Silva et al., 2011), and the expansion of industrial logging in Central Africa was recently documented (Laporte et al., 2007). Somewhat unusual for Africa is the observation that the areas of primary (untouched) forest, given by FAO (2010), are much smaller than the areas of primary forest determined here on the basis of land use. The total forest areas match, but, according to the FAO, most of the current forests are secondary or managed forests, while most of them are primary forests according to the rates of land use documented in this study.

This discrepancy between FAO estimates of primary forests and ours is difficult to explain at continental scale. Recent studies highlight the importance of forest degradation

## BGD

10, 8343–8413, 2013

### The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏮

⏭

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏮

⏭

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and conversion to agro-forestry systems as two potential underestimated drivers of understocking of primary forests. Most analysis reported deforestation as the land use change between forest to any other use, such as crop, pasture or even urban areas. Very few studies report the impact of selective logging on forest carbon stocks. Often the intensity of the disturbance caused by selective logging is regarded as negligible or even positive for the ecosystem and, when “sustainable”, supported by international certifications. In the past the intensity of the disturbance caused by selective logging was often quantified as the number of felled trees or type of machinery used for the construction of access roads and exploitation trails. These impacts have been assessed as minor disturbances by previous studies (Deckker and De Graaf, 2003). We have studied the impact of selective logging in African tropical forests on biomass and biodiversity losses (Cazzolla Gatti et al., 2013). The study sites selected were tropical forest areas located in West Africa along the border between Ghana and Ivory Coast (Bia National Park and surrounding areas) and Sierra Leone and Liberia (Gola National Park), and in Central Africa, within the Congo river basin, on the border between Cameroon and Central African Republic (Lobèkè National Park and surrounding areas). It was observed that the levels of biomass in forests subjected to selective logging are significantly lower than those of primary forests (even if there is a removal of very few trees per hectare) and that this reduction is more evident 30 yr after logging than after 20. The results showed a loss of carbon between 20 and 51 % of the initial primary forest value after 20 yr from selective logging and 24–72 % after a period of 30 yr. These figures highlight the significant loss of carbon even without changes in the main dominant land use (forest).

Another source of uncertainty is the role of conversion of primary forests to plantations. We have studied the impact of forest conversion to plantation on carbon stocks in a tropical rainforest area of West Ghana (Grieco et al., 2013). The study underlines that the land use change leads in all the cases to a significant loss of C stock when passing from forest to any of the other plantations analyzed (coconut, rubber, oil palm, cocoa, mixed).



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



According to Gibbs et al. (2010) nearly 60 % of new agricultural land in Africa was derived from intact forests. Africa had a net forest loss of 3.4 Mha annually for the period 2000–2010 (FRA 2010); if we assume that 60 % of this areas were converted into agricultural land we estimate 2.0 Mha yearly converted from forest to agriculture.

If we assume that 12.4 % of agricultural land in Africa is agroforestry with tree cover percentage > 30 % (Zomer et al., 2009), this results in a rough estimation of 253 000 ha of agroforestry land derived from deforestation of pristine forests in Africa every year. Considering a mean annual C loss per hectare of  $4.1 \pm 3.8$  Mg C due to conversion of forest into tree crop plantations (Grieco et al., 2013), this would cause an annual loss of around 1.0 Tg C in Africa.

These results show that, although deforestation remains the main driver of carbon losses from land use change, forest degradation and conversion, though lower and highly uncertain, are likely to increase carbon losses and need to be taken into account particularly considering the future economic development of Africa.

### 3.7 Wild fires and domestic biomass burning

Fires compete with heterotrophic respiration and herbivory to consume carbon that has been sequestered by photosynthesis. The amount of carbon combusted through “wild” fires (i.e. excluding the burning of fuelwood or residues, but including managed burning of natural vegetation) depends on the fire frequency, the amount of biomass available, what fraction of biomass that is exposed to fire is actually combusted (the combustion completeness) and the post-fire tree mortality and regrowth rates. In most African ecosystems, especially those that see frequent fires, fires mostly consume the herbaceous vegetation while a thick bark protects the trees. In addition to these fires that occur in the landscape, fire is used domestically for cooking and heating purposes. We will refer to these fires as domestic fires.

In general, fires are not a net source of CO<sub>2</sub> over decadal timescales unless fire is used in the deforestation process or if areas see an increase in the fire frequency. Humans ignite most fires in Africa (Archibald et al., 2010), and climate limits the



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



occurrence of fires in arid regions (due to a lack of fuel build-up) and in areas that see frequent rainfall even in the dry season (van Wilgen and Scholes, 1997; van der Werf et al., 2008). In between these extremes, the importance of climate is relatively minor. Archibald et al. (2011), for example, found that while climate could explain about 40 % of the variability in interannual variability in burned area in protected areas, outside these parks it could not account for much of the variability, highlighting the human control over fire regimes. This strong human component has not necessarily led to an increase in emissions over the past decades as often suggested; human interference with the landscape has led to a more patchy landscape where fuel beds are not continuous (Archibald et al., 2010), while more frequent grazing may have led to less fuel in general.

A recent review of emissions from all types of fire in Africa (wildfires, domestic fuel-wood and fire associated with land use change) is provided by Scholes et al. (2011). The authors concluded, based on a synthesis of many sources, that the total annual consumption of biomass is  $1.4 \pm 0.2 \text{ Pg DM yr}^{-1}$  (the uncertainty is the 95 % confidence interval), which corresponds to about  $0.7 \pm 0.1 \text{ Pg C yr}^{-1}$ . They further estimate that 57 % of this is due to wildfires, mostly in savannas, 36 % is domestic wood and charcoal, 3 % is associated with land use change and 4 % is the burning of crop residues.

Currently, the most accurate burned area estimates for Africa are derived from the MODIS sensors (Giglio et al., 2009, 2010; Roy et al., 2003). The two different algorithms used indicate that total annual burned area is on average  $250 \text{ Mha yr}^{-1}$  over the past decade, almost evenly split between Africa north and Africa south of the equator. This is almost 70 % of the global total burned area. Since a fire has to consume at least 50 % of the  $500 \times 500 \text{ m}$  grid cell (but most often more) to be detected as burned area, the burned area estimates are likely an underestimate due to missing small fires, for example those in agricultural areas.

Converting burned area estimates to emissions requires the use of models that take into account the variable fuel consumption due to landscape heterogeneity and changes in meteorological or climatic conditions that impact combustion completeness.

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Over larger scales, probably the most frequently used dataset is the Global Fire Emissions Database (GFED), which combines burned area from Giglio et al. (2010) with the satellite-driven Carnegie–Ames–Stanford–Approach biogeochemical model that estimates fuel loads based on carbon input from Net Primary Production (NPP), and losses through heterotrophic respiration, herbivory, fuelwood collection, and fires with a monthly timestep on a  $0.5^\circ \times 0.5^\circ$  grid (van der Werf et al., 2010).

Total annual carbon emissions estimates for Africa according to GFED were  $1.03 \pm 0.22 \text{ Pg C yr}^{-1}$ , averaged over the period 1997–2011 (Table 7; see also: [http://www.falw.vu/~gwerf/GFED/GFED3/tables/emis\\_C\\_absolute.txt](http://www.falw.vu/~gwerf/GFED/GFED3/tables/emis_C_absolute.txt)).

About 90 % of these emissions originated from savanna and woodland burning, with a smaller role for forest and deforestation fires. Since there is relatively more woodland burning in Africa south of the equator, mean fuel consumption was somewhat higher here ( $\sim 450 \text{ g C m}^{-2}$  burned) than in northern Africa ( $\sim 380 \text{ g C m}^{-2}$  burned) and total emissions were thus also somewhat higher in Southern Hemisphere Africa. These fuel consumption estimates are higher than most field measurements (Scholes et al., 2011) and GFED may be biased high, possibly partly balancing the underestimation due to missing small fires.

Fire emissions estimates are uncertain due to uncertainties in burned area, fuel loads, fire-combustion completeness and emission factors. A promising new approach uses the fire radiative power (FRP) measured by satellite, which removes much of the uncertainty in fuel load and combustion completeness, since it directly estimates the fuel combusted. Wooster et al. (2005) showed a direct relation between time-and-space integrated FRP (yielding fire radiative energy, FRE) and biomass combusted, and both MODIS (Ellicott et al., 2009) and SEVERI (Roberts and Wooster, 2008) data has been used to calculate emissions over the African continent. These estimates are about a factor 3 lower than those from GFED, but when combined with burned area estimates yield a better correspondence with fuel consumption measurements in the field than those from GFED. At the same time, combining measurements of fire-emitted gases from several sources with inverse modeling techniques yielded better agreement

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with modeled fire emissions (AMMABB) in West Africa when the L3JRC burned area product (Tansey et al., 2008) used in this database was boosted by a factor 2 and 1.67 for the land cover classes open broadleaf deciduous forest and deciduous shrubs, respectively (Lioussé et al., 2010). This emissions estimate is about twice as high as GFED although part of the discrepancy originated from using different emission factors, so estimate carbon emissions are more similar. Using the L3JRC product, Lehsten et al. (2009) estimated emissions for Africa of  $0.7 \text{ Pg C yr}^{-1}$  with similar fuel consumption as GFED. These divergent results indicate that fire emissions estimates are still uncertain.

On a global scale, Africa accounted for 72 % of global burned area, 52 % of global carbon emissions, 44 % of  $\text{CO}_2$  emissions and 36 % of  $\text{CH}_4$  emissions (van der Werf et al., 2010; Scholes et al., 2011). The smaller contribution of carbon and trace gas emissions is related to the dominance of savanna burning with relatively low fuels and clean combustion.

### 3.8 Lateral fluxes through river flows

To estimate the land to ocean carbon flux and transport to endorheic regions we used the 21 ocean coastline ( $26\,212\,740 \text{ km}^2$ ) and 8 endorheic ( $3\,706\,207 \text{ km}^2$ ) segments with their corresponding river catchments for Africa as described by the COSCAT database (Meybeck et al., 2006). The lateral transport of carbon to the coast was estimated at the river basin scale using the Global Nutrient Export from WaterSheds (NEWS) model framework (Mayorga et al., 2010), including NEWS basin areas. The carbon species models are hybrid empirically and conceptually based models that include single and multiple linear regressions developed by the NEWS effort and Hartmann et al. (2009), and single-regression relationships assembled from the literature. Modeled dissolved and particulate organic carbon (DOC and POC) loads used here (Mayorga et al., 2010) were generated largely using drivers corresponding to the year 2000, including observed hydro-climatological forcings, though some parameters and the observed loads are based on data spanning the previous two decade.

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Total suspended sediment (TSS) exports were also estimated by NEWS. Dissolved inorganic carbon (DIC) estimates correspond to weathering-derived bicarbonate exports and do not include  $\text{CO}_2$  supersaturation; the statistical relationships developed by Hartmann et al. (2009) were adjusted in highly weathered tropical soils (ferralsols) to 25 % of the modeled values found in Hartmann et al. (2009) to account for overestimates relative to observed river exports (as suggested in Hartmann and Moosdorf, 2011); adjusted grid-cell scale exports were aggregated to the basin scale using NEWS basin definitions (Mayorga et al., 2010), then reduced by applying a NEWS-based, basin scale consumptive water removal factor from irrigation withdrawals (Mayorga et al., 2010). DIC modeled estimates represent approximately 1970–2000. Overall, carbon loads may be characterized as representing general conditions for the period 1980–2000. Carbon, sediment and water exports were aggregated from the river basin scale to corresponding COSCAT regions. It is worth pointing out that Africa is under-represented in the data sets used to generate the regressions relationships.

The total amount of water discharge is  $4285 \text{ km}^3 \text{ yr}^{-1}$  ( $0.16 \text{ myr}^{-1}$ ) to the coast and  $268 \text{ km}^2 \text{ yr}^{-1}$  ( $0.07 \text{ myr}^{-1}$ ) to endorheic regions. These water yields are approximately an half of the global average of  $0.32 \text{ myr}^{-1}$  (Raymond et al., 2013). Furthermore, although the endorheic segments make up 12.4 % of the land mass of Africa they are responsible for only 5.9 % of the water discharge. DIC, DOC, and POC fluxes to the ocean are 29.40, 17.90, and  $15.70 \text{ TgCyr}^{-1}$ , respectively, or yields of 1.12, 0.68 and  $0.60 \text{ gm}^{-2}$  of land mass draining to the ocean. These global yields are also approximately an half of the global averages. Thus the yields of Africa are low globally due to low water throughput.

The Congo River dominates the dissolved fluxes from the African continent to the ocean, making up approximately 35 % of the water and 40 % of the DOC and DIC. The percent contribution of the Congo to POC fluxes are only 14 %. Approximately 73, 26, and 1 % of the dissolved carbon fluxes are to the Atlantic, Indian, and Mediterranean/Black Seas, respectively. For particulate organic carbon, approximately 66, 32,

and 2 % of the fluxes are to the Atlantic, Indian, and Mediterranean/Black Seas, respectively.

### 3.9 N<sub>2</sub>O and CH<sub>4</sub> fluxes from biogenic sources and fires

Although, excluding water vapor, N<sub>2</sub>O and CH<sub>4</sub> are the second and third most important greenhouse gases after CO<sub>2</sub> in terms of atmospheric concentration and radiative forcing effect on climate, studies on biogenic fluxes of both gases in African ecosystems are quite scanty and scaling up and yearly extrapolation estimates are strongly affected by high uncertainty. N<sub>2</sub>O is mainly driven by availability of N and its production is favored by low oxygen tension; however, prolonged anoxic conditions lead to further N<sub>2</sub>O reduction to N<sub>2</sub>, as it occurs in wetlands. In terrestrial ecosystems, the most favorable conditions for N<sub>2</sub>O production are when soil water filled pore space is between 60 % and 90 % (Davidson, 1993), which more frequently occurs in humid tropical forests (Castaldi et al., 2012), and in savannas during the rainy season (Castaldi et al., 2006), in particular during its onset. In arid and semiarid areas conditions are generally limiting for significant N<sub>2</sub>O production. In cropland areas high N<sub>2</sub>O emissions are favored by the addition of extra N inputs (organic or mineral N) but only at suitable soil water content high N<sub>2</sub>O fluxes can be expected.

The emissions of N<sub>2</sub>O from natural ecosystems were calculated from an empirical sigmoidal relationship, (Castaldi et al., 2013), between total annual N<sub>2</sub>O emission and total annual rainfall using flux data reported by Castaldi et al. (2006) for seasonally dry ecosystems and by Castaldi et al. (2012) for humid tropics. At continental level, this source represents 42.4 % of total (Table 8) with humid tropical forests contributing to 90 % of this fraction. In these ecosystems the environmental conditions allow for high productivity and high organic matter inputs to the soil, which, favored by high soil humidity and temperature during most of the year, create a fast flow of N in the ecosystem which can support high mineralization, nitrification and denitrification rates and N<sub>2</sub>O production (Castaldi et al., 2012).

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The second most important source is represented by agricultural ecosystems (29.9 %, Table 8), where the source of extra N to the soil derives from both fertilizer use and husbandry, the latter prevailing over the former in most areas (Potter et al., 2010).

In cultivated areas derived from quite recently deforested areas, a further input of extra N originates from mineralization of the organic matter left on the ground (slash, wood) after the conversion, and by the mineralization of soil organic matter, which is stimulated by management practices and disturbance.

Several studies show that during the first years following deforestation the site emits more  $N_2O$  than the original forest (control site). This extra flux of  $N_2O$  decreases exponentially with time and within 20–40 yr, depending on the study, the deforested site baseline flux levels off stabilizing around annual  $N_2O$  values lower (around 50 %) than the original forest. An exponential decay fitting equation was applied to available data on deforestation (Castaldi et al., 2013) to calculate the background annual  $N_2O$  emission in agricultural areas for the African continent, assuming that all crop areas derived from forest clearing. To define the year of land use change yearly maps of land use were derived from the dataset on Global Cropland and Pasture Data by Ramankutty and Foley (1999). This source is estimated to be  $69 \text{ GgN}_2\text{Oyr}^{-1}$ , thus about 2.5 % of the total  $N_2O$  budget.

Indirect sources of  $N_2O$  are emissions associated with N deposition and N export from land to rivers (Table 8). The former are dominated by “fire generated N compounds” and at a minor extent by agricultural practices. A very low contribution comes from anthropogenic pollution (Dentener, 2006). N export and consequent derived  $N_2O$  emissions are subject to an enormous uncertainty associated both to N export quantification and emission coefficient estimates (Van Dreht et al., 2003; IPCC, 1997). Another relevant source is represented by direct  $N_2O$  emissions from fires (natural and agricultural), which accounts for 16.4 % of the total annual  $N_2O$  emission to the atmosphere from the African continent.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



More biological processes contribute to the continental budget of  $\text{CH}_4$ . Biogenic  $\text{CH}_4$  is produced by metabolic activity of methanotrophic bacteria which transform fermentation products, formed by other microorganisms, in  $\text{CH}_4$  when conditions are strictly anaerobic.  $\text{CH}_4$  production occurs in wetlands and saturated soils, anaerobic hotspots in unsaturated soils, mostly associated to organic input and intense mineralization activity.  $\text{CH}_4$  is also produced by enteric fermentation in the rumen and hindgut of animals, being greatest in ruminants, and in particular in domesticated ones (Crutzen et al., 1986). However, also several invertebrates host methanogenic organisms in their gut, termites being the most relevant in terms of total biomass and  $\text{CH}_4$  emission rates.  $\text{CH}_4$  is oxidated by chemoautotrophic aerobic bacteria. Most typically methanotrophic activity occurs in well aerated soils, but it also contributes to the net ecosystem exchange of  $\text{CH}_4$  in the column of water and in saturated soils, when sufficient oxygen is available. Taking into account  $\text{CH}_4$  biological sources and sinks, the African  $\text{CH}_4$  budget is dominated by emitting processes, natural wetlands and animal husbandry, being the most relevant sources (49 % and 25.5 %, respectively, Table 9). Wetlands emissions are, however, characterized by high uncertainty due to widely differing estimates from the few available studies (Table 9). Rice paddies represent a minor source in Africa due to their smaller extent compared to other tropical countries (Table 9). Tropical humid forests and pastures are a weak net  $\text{CH}_4$  sources. In the former, net  $\text{CH}_4$  emissions in lowland areas are balanced by the net  $\text{CH}_4$  sink of upland areas; whereas pastures are generally net sources of  $\text{CH}_4$  during the wettest months (Castaldi et al., 2006). Soils in seasonally dry forests and savannas are instead a net  $\text{CH}_4$  sink (Castaldi et al., 2006), which, given the wide extension of these ecosystems, counterbalances the small source of  $\text{CH}_4$  from humid forests, resulting, at continental level, in a small net  $\text{CH}_4$  sink (Table 9). Emissions from fires represent the third largest  $\text{CH}_4$  source at continental level, due to the wide surface of territory which gets burned every year, especially in arid and seasonally dry areas. Estimates of  $\text{CH}_4$  emissions from termites are highly variable depending on the proposed approach for calculations (from 2.5 to  $6.9 \text{ Tg CH}_4 \text{ yr}^{-1}$ , Table 9). The number of studies which allow for good up-scaling is



scanty, so that all proposed estimates are based on very rough extrapolations. Biogenic sources and fires overall lead to an average total emissions of  $70 \text{ Tg CH}_4 \text{ yr}^{-1}$ . In terms of  $\text{CO}_2$  equivalents, this would correspond to an average emissions of  $1.75 \text{ Pg CO}_2\text{-eq yr}^{-1}$ .

## 4 Interannual variability

The African continent is responsible for a large fraction of the global carbon cycle's interannual variability (Williams et al., 2007), as indicated by both ecosystem process models and atmospheric inversions solving for surface sources and sinks of  $\text{CO}_2$ . Nearly a quarter of the year-to-year variation in the global scale exchange of  $\text{CO}_2$  between land and atmosphere can be attributed to the African continent (Williams et al., 2007).

The major source of this variability is gross primary productivity whose variation exceeds that of ecosystem respiration (Ciais et al., 2009; Weber et al., 2009; Williams et al., 2008). Hydroclimatic fluctuations are the prevailing cause, with much of the continent being water-limited and subject to highly variable precipitation inputs (Weber et al., 2009; Williams et al., 2008). There is wide disagreement among models regarding the magnitude and spatial allocation of interannual variations in ecosystem productivity (Weber et al., 2009) (e.g. Fig. 6). Models do tend to agree that variability is high in the Sudanian to Sahelian transition zone, equatorial East Africa, and south-central to southeastern Africa. These are regions where hydroclimatic fluctuations are reported to be correlated with El Niño Southern Oscillation (ENSO) and/or the Indian Ocean Dipole (or Zonal Mode) climate oscillations (Nicholson and Kim, 1997; Reason, 2002; Reason and Rouault, 2002; Richard et al., 2000). Recent work has demonstrated their sizeable effects on carbon exchanges (Ciais et al., 2009; Williams, 2010; Williams and Hanan, 2011) (Fig. 7). While important, we note that such teleconnections explain only a small fraction of continent-wide interannual variability in water and carbon balances.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Longer-term hydroclimatic trends are another important source of variability at the regional scale, and have been noted as particularly strong in the Sudano-Sahelian zone (e.g. Nicholson, 2000; Tyson et al., 2002; Nicholson and Kim, 1997; Nicholson, 2001). This drives corresponding periods of greening or browning at decadal scales (Prince et al., 2007; Herrmann et al., 2005). Such patterns are also reflected in carbon sinks or sources and can be attributed to increases to decreases in precipitation, but have also been linked to growth enhancement from rising concentrations of carbon dioxide (Ciais et al., 2009) (Fig. 8).

Though Africa contributes a large fraction of global fire emissions, at nearly half of the global total (see Sects. 3.7), the continent's pyrogenic emissions are less interannually variable in both relative and absolute terms compared to other regions of the world, such as southeast Asia and Oceania, as well as South America (van der Werf et al., 2006).

## 5 Integrating the African GHG budget

Table 10 synthesizes the best available current estimates of carbon fluxes and pools in Africa, as well as flux data for methane and nitrous oxide. There are still large uncertainties within (and between) each of the approaches used (inventory, ecosystem fluxes, models, atmospheric inversions). For each of the different approach used we have calculated, on the basis of the best available data, the Net Carbon Balance as follow: NEP – Fires – Land use – lateral transport (DOC only). From these results we can conclude that the contribution by the African continent to the global carbon cycle range between a slight source of  $0.05 \pm 0.28 \text{ PgCyr}^{-1}$  (estimated by the atmospheric inversion, excluding fossil fuel) to a sink of  $-1.34 \pm 1.32 \text{ PgCyr}^{-1}$  (estimated by the ecosystem fluxes). The median estimate is a sink of  $-0.58 \text{ PgCyr}^{-1}$ . Interestingly three different approaches provide a quite similar GPP fluxes ( $31.7 \pm 6.6$ ,  $24.3 \pm 2.9$ ,  $28.2 \pm 5.9 \text{ PgCyr}^{-1}$  for inventory data, ecosystem fluxes and DGVM model, respectively). NPP also seems to be well constrained by both inventory and DGVM estimates

( $14.3 \pm 3.7$  and  $13.3 \pm 3.3$  Pg C yr<sup>-1</sup>, respectively). Thus the disagreements between approaches reside in the respiration fluxes (broadly defined, to include domestic and wild fire and decomposition of soil, biomass and litter stocks following land cover change).

Although total GPP, NPP seems to be reasonably constrained, respiratory fluxes show high uncertainties and reflect the difficulty of both measurements and a modeling approach. In particular the assumptions to derive respiratory fluxes by inventory are weak. This is partly due to the paucity of data and in situ measurements able to represent the full range of African ecosystems. The similar estimates for heterotrophic respiration derived from inventory and DGVMs is likely to be coincidental.

The comparison of inventories, ecosystem flux measurements and DGVM estimates of NEE with atmospheric inversions can only be carried out by correctly accounting for the lateral transport of carbon and for the land use change emissions. For this reason we have compiled relevant data on land use emissions and lateral transport (see Sects. 3.6 and 3.8, respectively) as common input data for all the approaches. While all the approaches based on inventories, ecosystem fluxes and models indicate a modest carbon sink ( $-0.74 \pm 1.19$ ,  $1.34 \pm 1.32$ ,  $-0.41 \pm 0.31$  Pg C yr<sup>-1</sup>, respectively), the atmospheric inversions, excluding fossil fuel, show a small carbon source ( $0.05 \pm 0.28$  Pg C yr<sup>-1</sup>). There are several reasons for such discrepancies, mostly related to the features of the individual approaches and their assumptions. The discrepancies are within the uncertainty associated with the various approaches. In the case of inventory, the reconstruction of carbon fluxes from NPP is based on a few datasets from savannas and some speculative exercises for tropical forests. Flux data and their spatial integration in Africa suffer from the lack of adequate representation of all ecosystem types. The likely effects of terrain and night time fluxes on the estimation of the overall annual budget (and in particular the respiration fluxes) are severe. In the paper the possibility of some overestimation of net ecosystem exchanges from flux data is also considered. DGVM may not adequately represent plant functional types in Africa and respiratory fluxes. Furthermore, the emissions by unsustainable land use, mainly fuel wood extraction leading to forest degradation may be underestimated. Atmospheric

## BGD

10, 8343–8413, 2013

### The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



inversions are affected by the lack of a reasonable network of observation sites in Africa. In addition inter-annual variability plays an important role and the overall carbon balance may easily swing from a sink to a source of carbon following climate variability and anthropogenic changes.

5 The contribution of  $\text{N}_2\text{O}$  and  $\text{CH}_4$  fluxes to the overall GHG balance of the African continent is equivalent to  $2.57 \pm 1.28 \text{ PgCO}_2\text{-eq yr}^{-1}$ .  $\text{CH}_4$  is the dominant emission ( $1.75 \pm 0.85 \text{ PgCO}_2\text{-eq yr}^{-1}$ ) followed by  $\text{N}_2\text{O}$  to a lesser extent ( $0.82 \pm 0.41 \text{ PgCO}_2\text{-eq yr}^{-1}$ ), although the latter has a larger global warming potential. If we compare these estimates with the carbon balance of the continent we can say that the role of non-  
10  $\text{CO}_2$  gases emissions is prevalent in making the continent a net source of greenhouse gases.

## 6 Key uncertainties and research gaps

After having assessed the availability of data, the key processes involved in the GHG balance of Africa, and the suitability of the various methods that have been used, some  
15 conclusions can be drawn here below:

1. There is a general lack of data and information on relevant processes at ecosystem level, able to constrain atmospheric data and DGVM models. This includes respiratory fluxes and NPP data, particularly for tropical rainforest, but also for several other ecosystems such as savannas and the woodlands transitional to forests.  
20
2. Land use emissions are increasingly well-studied, but still there are gaps relating to forest degradation and other land use that make the African continent unique, such as selective logging, biomass use for charcoal production and the effects of shifting agriculture.

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3. Interannual variability is the major feature of the continental carbon fluxes, believed to be on the order of  $\pm 0.5 \text{ Pg C yr}^{-1}$  in standard deviation accounting for as much as a quarter of the year-to-year variation in the global carbon budget.
4. It seems that climatic fluctuations (particularly as they impact the water balance) are an important source of variability at the regional scale, and have been noted as particularly strong in the Sudano-Sahelian zone. Integrating the biogeochemical cycles of carbon and water research is therefore of particular importance for understanding the dynamics of the African carbon balance.
5. Non- $\text{CO}_2$  greenhouse gas emissions are poorly studied across the various African ecosystems. In particular, little is known about how they are affected by land use and climate. The lack of such information is critical for understanding the African methane budget at global level and to provide an insight on the natural sources of nitrous oxide.
6. Since both climate and human pressures will play an important role on the future of the African region a greater effort on in situ and atmospheric observations is urgently needed and should also comprise socio-economic drivers of change. Integrated inter-disciplinary research would allow us to reduce uncertainties in the carbon and other GHG fluxes of the continent and to understand its temporal and spatial variability in response of climate and land use drivers.

## 7 Conclusions

The vast majority of the results from this study and the literature, based on different methodologies and for different time periods (see Table 10), seem to agree that Africa is probably a small sink of carbon on an annual time scale. By averaging out all the estimates this value is  $-0.61 \pm 0.58 \text{ Pg C yr}^{-1}$ . Nevertheless the emissions of  $\text{CH}_4$  and  $\text{N}_2\text{O}$  may turn Africa into a carbon source.

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



At sub-regional level there is a significant spatial variability in both carbon sources and sinks, mainly due to the biome's change and the different anthropic impact, with southern Africa as the main source and central Africa as the main sink. In particular 33 % of the global tropical forest carbon sink (and 16 % of the global terrestrial carbon sink) is estimated to be due to African tropical forests that contain around 78 Pg of above ground biomass.

Emissions from land use change in Africa are significant (around  $0.32 \pm 0.05 \text{ PgCyr}^{-1}$ ) and even higher than the fossil fuel ones; this is a unique feature among all the continents. In addition there can be significant carbon losses from land even without changes in the land use (forest), as results from the impact of selective logging.

Fires also play a significant role, with around  $1.03 \pm 0.22 \text{ PgCyr}^{-1}$  of annual carbon emissions mainly (90 %) originated by savanna and woodland burning. On a global scale, this contributes to more than half (52 %) of the global carbon emissions by fires, and more than one third (36 %) of  $\text{CH}_4$  emissions.

The African continent is responsible for a large fraction (around 25 %) of the global carbon cycle's interannual variability.

These figures on the whole, even if still highly uncertain, show the important role Africa plays in the carbon cycle at global level, both in terms of absolute values and variability.

**Acknowledgements.** The research leading to these results has received funding from the European Research Council (ERC) under grant agreement no. 247349 "Africa-GHG".

We would like to thank also the developers of the CarboScope web site (<http://www.carboscope.eu>), funded by the CNRS (Centre National de la Recherche Scientifique) from which inversion results were made available to us, and the modeling groups, in particular the Trendy modelers for having made the simulations.

## References

UN, Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat, World Population Prospects: The 2010 Revision, <http://esa.un.org/unpd/wpp/index.htm>, 2011.

- 5 Alam, S. A., Starr, M., and Clark, B. J. F.: Tree biomass and soil organic carbon densities across the Sudanese woodland savannah: a regional carbon sequestration study, *J. Arid Environ.*, 89, 67–76, 2013.

Archibald, S., Nickless, A., Govender, N., Scholes, R. J., and Lehsten, V.: Climate and the inter-annual variability of fire in southern Africa: a meta-analysis using long-term field data and satellite-derived burnt area data, *Global Ecol. Biogeogr.*, 19, 794–809, 2010.

- 10 Archibald, S., Staver, A. C., and Levin, S. A.: Evolution of human-driven fire regimes in Africa, *P. Natl. Acad. Sci. USA*, 109, 847–852, 2011.

Arneth, A., Lehsten, V., Thonicke, K., and Spessa, A.: Climate-fire interactions and savanna ecosystems: a dynamic vegetation modelling study for the African continent, in: *Ecosystem Function in Savannas: Measurement and Modeling at Landscape to Global Scales*, edited by: Hill, M. J. and Hanan, N. P., Taylor & Francis/CRC, Boca Raton, 463–478, 2010a.

- 15 Arneth, A., Sitch, S., Bondeau, A., Butterbach-Bahl, K., Foster, P., Gedney, N., de Noblet-Ducoudré, N., Prentice, I. C., Sanderson, M., Thonicke, K., Wania, R., and Zaehle, S.: From biota to chemistry and climate: towards a comprehensive description of trace gas exchange between the biosphere and atmosphere, *Biogeosciences*, 7, 121–149, doi:10.5194/bg-7-121-2010, 2010b.

20 Aselmann, I. and Crutzen, P. J.: Global distribution of natural freshwater wetlands and rice paddies, and their net primary productivity, seasonality and possible methane emissions, *J. Atmos. Chem.*, 8, 307–358, 1989.

- 25 Baccini, A., Laporte, N., Goetz, S. J., Sun, M., and Dong, H.: A first map of tropical Africa's above-ground biomass derived from satellite imagery, *Environ. Res. Lett.*, 3, 045011, doi:10.1088/1748-9326/3/4/045011, 2008.

- 30 Baccini, A., Goetz, S. J., Walker, W. S., Laporte, N. T., Sun, M., Sulla-Menashe, D., Hackler, J., Beck, P. S. A., Dubayah, R., Friedl, M. A., Samanta, S., and Houghton, R. A.: Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps, *Nature Clim. Change*, 2, 182–185, 2012.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Banker, B. C., Hludze, H. K., Alford, D. P., DeLaune, R. D., and Lindau, C. W.: Methane sources and sinks in rice paddy soils: relationship to emissions, *Agr. Ecosyst. Environ.*, 53, 243–251, 1995.
- Bartlett, K. B., Crill, P. M., Sebacher, D. I., Harriss, R. C., Wilson, J. O., and Melack, J. M.: Methane flux from the central Amazonian floodplain, *J. Geophys. Res.*, 93, 1571–1582, 1988.
- Bombelli, A., Henry, M., Castaldi, S., Adu-Bredu, S., Arneth, A., de Grandcourt, A., Grieco, E., Kutsch, W. L., Lehsten, V., Rasile, A., Reichstein, M., Tansey, K., Weber, U., and Valentini, R.: An outlook on the Sub-Saharan Africa carbon balance, *Biogeosciences*, 6, 2193–2205, doi:10.5194/bg-6-2193-2009, 2009.
- Bonan, G. B. and Levis, S.: Quantifying carbon-nitrogen feedbacks in the Community Land Model (CLM4), *Geophys. Res. Lett.*, 37, L07401, doi:10.1029/2010GL042430, 2010.
- Bond, W. J., Woodward, F. I., and Midgley, G. F.: The global distribution of ecosystems in a world without fire, *New Phytol.*, 165, 525–538, 2005.
- Bouwman, A. F., Boumans, L. J. M., and Batjes, N. H.: Modeling global annual  $\text{N}_2\text{O}$  and  $\text{NO}$  emissions from fertilized fields, *Global Biogeochem. Cy.*, 16, 1080–1090, 2002.
- Brown, S. and Gaston, G.: Use of forest inventories and geographic information systems to estimate biomass density of tropical forests: Application to tropical Africa, *Environ. Monitor. Assess.*, 38, 157–168, 1995.
- Bousquet, P., Ciais, P., Peylin, P., Ramonet, M., and Monfray, P.: Inverse modeling of annual atmospheric  $\text{CO}_2$  sources and sinks 1. Method and control inversion, *J. Geophys. Res.-Atmos.*, 104, 26161–26178, 1999.
- Bucini, G. and Hanan, N. P.: A continental-scale analysis of tree cover in African savannas, *Global Ecol. Biogeogr.*, 16, 593–605, 2007.
- Canadell, J. G., Raupach, M. R., and Houghton, R. A.: Anthropogenic  $\text{CO}_2$  emissions in Africa, *Biogeosciences*, 6, 463–468, doi:10.5194/bg-6-463-2009, 2009a.
- Canadell, J. G., Ciais, P., Dhakal, S., Le Quéré, C., Patwardhan, A., and Raupach, M. R.: The human perturbation of the carbon cycle, UNESCO-UNEP-SCOPE Policy Brief, Paris, 6 pp., 2009b.
- Canadell, J. G., Ciais, P., Gurney, K., Le Quéré, C., Piao, S., Raupach, M. R., and Sabine, C. L.: An international effort to quantify regional carbon fluxes, *EOS*, 92, 81–82, 2011.
- Castaldi, S., Ermice, A., and Strumia, S.: Fluxes of  $\text{N}_2\text{O}$  and  $\text{CH}_4$  from soils of savannas and seasonally-dry ecosystems, *J. Biogeogr.*, 33, 401–415, 2006.



- Castaldi, S., Bertolini, T., Valente, A., Chiti, T., and Valentini, R.: Nitrous oxide emissions from soil of an African rain forest in Ghana, *Biogeosciences Discuss.*, 9, 16565–16588, doi:10.5194/bgd-9-16565-2012, 2012.
- 5 Cazzolla Gatti, R., Castaldi, S., Lindsell, J. A., Coomes, D. A., Marchetti, M., Maesano, M., Di Paola, A., Paparella, F., and Valentini, R.: The impact of selective logging on biodiversity and above-ground biomass of African tropical forests, in preparation, 2013.
- Chevallier, F., Maksyutov, S., Bousquet, P., Bréon, F.-M., Saito, R., Yoshida, Y., and Yokota, T.: On the accuracy of the CO<sub>2</sub> surface fluxes to be estimated from the GOSAT observations, *Geophys. Res. Lett.*, 36, L19807, doi:10.1029/2009GL040108, 2009.
- 10 Chevallier, F., Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P., Brunke, E. G., Ciattaglia, L., Esaki, Y., Fröhlich, M., Gomez, A., Gomez-Pelaez, A. J., Haszpra, L., Krummel, P., Langenfelds, R., Leuenberger, M., Machida, T., Maignan, F., Matsueda, H., Morguí, J. A., Mukai, H., Nakazawa, T., Peylin, P., Ramonet, M., Rivier, L., Sawa, Y., Schmidt, M., Steele, P., Vay, S. A., Vermeulen, A. T., Wofsy, S., and Worthy, D.: CO<sub>2</sub> surface fluxes at grid point scale estimated from a global 21 yr reanalysis of atmospheric measurements, *J. Geophys. Res.*, 115, D21307, doi:10.1029/2010JD013887, 2010.
- 15 Ciais, P., Piao, S.-L., Cadule, P., Friedlingstein, P., and Chédin, A.: Variability and recent trends in the African terrestrial carbon balance, *Biogeosciences*, 6, 1935–1948, doi:10.5194/bg-6-1935-2009, 2009.
- 20 Ciais, P., Bombelli, A., Williams, M., Piao, S. L., Chave, J., Ryan, C. M., Henry, M., Brender, P., and Valentini, R.: The carbon balance of Africa: synthesis of recent research studies, *Philos. T. Roy. Soc. A*, 269, 2038–2057, 2011.
- Collatz, G. J., Ball, J. T., Grivet, C., and Berry, J. A.: Physiological and Environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer, *Agr. Forest Meteorol.*, 54, 107–136, 1991.
- 25 Collatz, G. J., Ribascarbo, M., and Berry, J. A.: Coupled photosynthesis-stomatal conductance model for leaves of C<sub>4</sub> plants, *Aust. J. Plant. Physiol.*, 19, 519–538, 1992.
- Cox, P. M.: Description of the “TRIFFID” dynamic global vegetation model, Hadley Centre Technical Note 24, Hadley Centre, Met Office, London Road, Bracknell, Berks, RG122SY, UK, 17 pp., 2001.
- 30 Cramer, W., Kicklighter, D. W., Bondeau, A., Moore III, B., Churkina, G., Nemry, B., Ruimy, A., Schloss, A. L., and the participants of the Potsdam NPP Model Intercomparison: Comparing

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

global models of terrestrial net primary productivity (NPP): overview and key results, *Glob. Change Biol.*, 5, 1–15, 1999.

Crutzen, P. J., Aselman, I., and Seiler, W.: Methane production by domestic animals, wild ruminants, other herbivorous fauna, and humans, *Tellus B*, 38, 271–284, 1986.

5 Davidson, E. A.: Soil water content and the ratio of nitrous to nitric oxide emitted from soil, in: *Biogeochemistry of Global Change: radiatively active trace gases*, edited by: Orem-land, R. S., Chapman & Hall, London, 369–386, 1993.

Davis, S. J., Peters, G. P., and Caldeira, K.: The supply chain of CO<sub>2</sub> emissions, *P. Natl. Acad. Sci. USA*, 108, 18554–18559, doi:10.1073/pnas.1107409108, 2011.

10 Deckker, M. and de Graaf, N. R.: Pioneer and climax tree regeneration following selective logging with silviculture in Suriname, *Forest Ecol. Manag.*, 172, 183–190, 2003.

Delmas, R. A., Tathy, J. P., and Cross, B.: Atmospheric methane budget in Africa, *J. Atmos. Chem.*, 14, 395–409, 1992.

Dentener, F., Drevet, J., Lamarque, J. F., Bey, I., Eickhout, B., Fiore, A. M., Hauglustaine, D.,  
15 Horowitz, L. W., Krol, M., Kulshrestha, U. C., Lawrence, M., Galy-Lacaux, C., Rast, S., Shindell, D., Stevenson, D., Van Noije, T., Atherton, C., Bell, N., Bergman, D., Butler, T., Cofala, J., Collins, B., Doherty, R., Ellingsen, K., Galloway, J., Gauss, M., Montanaro, V., Müller, J. F., Pitari, G., Rodriguez, J., Sanderson, M., Solomon, F., Strahan, S., Schultz, M., Sudo, K., Szopa, S., and Wild, O.: Nitrogen and sulfur deposition on regional and global scales: a multi-model evaluation, *Global Biogeochem. Cy.*, 20, GB4003, doi:10.1029/2005GB002672, 2006.

Devol, A. H., Richey, J. E., Clark, W. A., and King, S. L.: Methane emissions to the troposphere from the Amazon floodplain, *J. Geophys. Res.*, 93, 1583–1592, 1988.

Ellicott, E., Vermote, E., Giglio, L., and Roberts, G.: Estimating biomass consumed from fire using MODIS FRE, *Geophys. Res. Lett.*, 36, L13401, doi:10.1029/2009GL038581, 2009.

25 FAO: Global Forest Resources Assessment 2005, Main Report, Progress Towards Sustainable Forest Management, FAO Forestry paper 147, Food and Agriculture Organization of the United Nations, Rome, Italy, 320 pp., 2006.

FAO: Global Forest Resources Assessment 2010, FAO Forestry Paper 163, Food and Agriculture Organization of the United Nations, Rome, 340 pp., available at: <http://www.fao.org/forestry/fra/fra2010/en/>, 2010.

30 FAOSTAT: available at: <http://faostat.fao.org/site/377/default.aspx#ancor>, last access: 11/09, 2009.

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Farquhar, G. D., von Caemmerer, S., and Berry, J. A.: A biochemical model of photosynthetic CO<sub>2</sub> assimilation in leaves of C<sub>3</sub> species, *Planta*, 149, 78–90, 1980.
- Friedlingstein, P., Houghton, R. A., Marland, G., Hackler, J., Boden, T. A., Conway, T. J., Canadell, J. G., Raupach, M. R., Ciais, P., and Le Quéré, C.: Update on CO<sub>2</sub> emissions, *Nat. Geosci.*, 3, 811–812, 2010.
- Gibbs, H. K., Ruesch, A. S., Achard, F., Clayton, M. K., Holmgren, P., Ramankutty, N., and Foley, J. A.: Tropical forests were the primary sources of new agricultural land in the 1980s and 1990s, *P. Natl. Acad. Sci. USA*, 107, 16732–16737, 2010.
- Giglio, L., Loboda, T., Roy, D. P., Quayle, B., and Justice, C. O.: An active-fire based burned area mapping algorithm for the MODIS sensor, *Remote Sens Environ*, 113, 408–420, 2009.
- Giglio, L., Randerson, J. T., van der Werf, G. R., Kasibhatla, P. S., Collatz, G. J., Morton, D. C., and DeFries, R. S.: Assessing variability and long-term trends in burned area by merging multiple satellite fire products, *Biogeosciences*, 7, 1171–1186, doi:10.5194/bg-7-1171-2010, 2010.
- Gilmanov, T. G., Aires, L., Beileli, L., Barcza, Z., Baron, V. S., Beringer, J., Billesbach, D., Bonal, D., Bradford, J., Ceschia, E., Cook, D., Corradi, C., Frank, A., Gianelle, D., Gimeno, C., Gruenwald, T., Guo, H., Hanan, N., Haszpra, L., Heilman, J., Jacobs, A., Jones, M. B., Johnson, D. A., Kiely, G., Li, S., Magliulo, V., Moors, E., Nagy, Z., Nasyrov, M., Owensby, C., Pinter, K., Pio, C., Reichstein, M., Sanz, M. J., Scott, R., Soussana, J. F., Stoy, P. C., Svejcar, T., Tuba, Z., and Zhou, G.: Productivity, respiration, and light-response parameters of world grassland and agro-ecosystems derived from flux-tower measurements, *Rangeland Ecol. Manag.*, 63, 16–39, doi:10.2111/REM-D-09-00072.1, 2010.
- Glenday, J.: Carbon storage and emissions offset potential in an African dry forest, the Arabuklo-Sokoke Forest, Kenya, *Environ. Monit. Assess.*, 142, 85–95, 2008.
- Grace, J., San Jose, J., Meir, P., Miranda, H. S., and Montes, R. A.: Productivity and carbon fluxes of tropical savannas, *J. Biogeogr.*, 33, 387–400, 2006.
- Grieco, E., Chiti, T., Perugini, L., and Valentini, R.: Deforestation and land use change in Jomoro District, Ghana, impacts on carbon stocks and perspectives of REDD+, in preparation, 2013.
- Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fan, S., Fung, I. Y., Gloor, M., Heimann, M., Higuchi, K., John, J., Maki, T., Maksyutov, S., Masarie, K., Peylin, P., Prather, M., Pak, B. C., Randerson, J., Sarmiento, J., Taguchi, S., Takahashi, T., and Yuen, C.-W.: Towards robust regional estimates of CO<sub>2</sub> sources and sinks using atmospheric transport models, *Nature*, 415, 626–630, 2002.

- Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Pak, B. C., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y. H., Ciais, P., Fung, I. Y., Heimann, M., John, J., Maki, T., Maksyutov, S., Peylin, P., Prather, M., and Taguchi, S.: Transcom 3 CO<sub>2</sub> inversion intercomparison: 1. Annual mean control results and sensitivity to transport and prior flux information, *Tellus B*, 55, 555–579, 2003.
- Hadi, A., Inubushi, K., Furukawa, Y., Purnomo, E., Rasmadi, M., and Tsuruta, H.: Greenhouse gas emissions from tropical peatlands of Kalimantan, Indonesia, *Nutr. Cycl. Agroecosys.*, 71, 73–80, 2005.
- Hartmann, J., Jansen, N., Durr, H. H., Kempe, S., and Kohler, P.: Global CO<sub>2</sub> consumption by chemical weathering: what is the contribution of highly active weathering regions?, *Global Planet. Change*, 69, 185–194, 2009.
- Hartmann, J., and Moosdorf, N.: Chemical weathering rates of silicate-dominated lithological classes and associated liberation rates of phosphorus on the Japanese Archipelago - implications for global scale analysis, *Chem. Geol.*, 287, 125–157, doi:10.1016/j.chemgeo.2010.12.004, 2011.
- Henry, M.: Carbon Stocks and Dynamics in Sub Saharan Africa, PhD thesis, University of Tuscia, 2010.
- Herrmann, S. M., Anyamba, A., and Tucker, C. J.: Recent trends in vegetation dynamics in the African Sahel and their relationship to climate, *Global Environ. Chang.*, 15, 394–404, 2005.
- Hickler, T., Smith, B., Prentice, I. C., Mjöfors, K., Miller, P., Arneth, A., and Sykes, M.: CO<sub>2</sub> fertilization in temperate FACE experiments not representative of boreal and tropical forests, *Glob. Change Biol.*, 14, 1–12, 2008.
- Hickman, J. E., Havlikova, M., Kroeze, C., and Palm, C. A.: Current and future nitrous oxide emissions from African agriculture, *Curr. Opin. Environ. Sustain.*, 3, 370–378, 2011.
- Houghton, R. A. and Hackler, J. L.: Emissions of carbon from land use change in sub-Saharan Africa, *J. Geophys. Res.*, 111, G02003, doi:10.1029/2005JG000076, 2006.
- IFAD: Rural Poverty Report 2001 – The Challenge of Ending Rural Poverty, International Fund for Agricultural Development, Rome, 266 pp., 2001.
- IPCC: Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, edited by: Houghton, J. T., Meira Filho, L. G., Lim, B., Tranton, K., Mamaty, I., Bonduki, Y., Griggs, D. J., and Callander B. A., Intergovernmental Panel on Climate Change, Meteorological Office, Bracknell, United Kingdom, Vol. 3: Greenhouse Gas Inventory Reference Manual, 482 pp, 1997.

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



IPCC: Changes in Atmospheric Constituents and Radiative Forcing, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment, Cambridge University Press, Cambridge, UK and New York, NY, USA, 129–234, 2007.

IPCC: AR4 WGII, Inter-Relationships Between Adaptation and Mitigation, Chapter 18, 2007b. Jung, M., Reichstein, M., and Bondeau, A.: Towards global empirical upscaling of FLUXNET eddy covariance observations: validation of a model tree ensemble approach using a biosphere model, *Biogeosciences*, 6, 2001–2013, doi:10.5194/bg-6-2001-2009, 2009.

Jung, M., Reichstein, M., Margolis, H., Cescatti, A., Richardson, A., Arain, A., Arneth, A., Bernhofer, C., Bonal, D., Chen, J., Gianelle, D., Gobron, N., Kiely, G., Kutsch, W., Lasslop, G., Law, B., Lindroth, A., Merbold, L., Montagnani, L., Moors, E., Papale, D., Sottocornola, M., Vaccari, F. P., and Williams, C.: Global patterns of land-atmosphere fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and meteorological observations, *J. Geophys. Res.*, 116, G00J07, doi:10.1029/2010JG001566, 2011.

Kesselmeier, J., Ciccioli, P., Kuhn, U., Stefani, P., Biesenthal, T., Rottenberger, S., Wolf, A., Vitullo, M., Valentini, R., Nobre, A., Kabat, P., and Andreae, M. O.: Volatile organic compound emissions in relation to plant carbon fixation and the terrestrial carbon budget, *Global Biogeochem. Cy.*, 16, 1126, doi:10.1029/2001GB001813, 2002.

Kirschke, S., Bousquet, P., Ciais, P., Saunio, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Fraser, P. J., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Krummel, P. B., Lamarque, J., Langenfelds, R. L., Le Quéré, C., Naik, V., O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., van der Werf, G. R., Voulgarakis, A., van Weele, M., Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of methane sources and sinks: budgets and variations, *Nat. Geosci.*, submitted, 2013.

Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S., and Prentice, I. C.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, *Global Biogeochem. Cy.*, 19, GB1015, doi:10.1029/2003GB002199, 2005.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kutsch, W. L., Hanan, N., Scholes, B., McHugh, I., Kubheka, W., Eckhardt, H., and Williams, C.: Response of carbon fluxes to water relations in a savanna ecosystem in South Africa, *Biogeosciences*, 5, 1797–1808, doi:10.5194/bg-5-1797-2008, 2008.
- Kutsch, W. L., Merbold, L., Ziegler, W., Mukelabai, M. M., Muchinda, M., Kolle, O., and Scholes, R. J.: The charcoal trap: Miombo forests and the energy needs of people, *Carbon Balance and Management*, 6:5, doi:10.1186/1750-0680-6-5, 2011.
- Kuze, A., Suto, H., Nakajima, M., and Hamazaki, T.: Thermal and near infrared sensor for carbon observation Fourier-transform spectrometer on the Greenhouse Gases Observing Satellite for greenhouse gases monitoring, *Appl. Optics*, 48, 6716–6733, 2009.
- Laporte, N. T., Stabach, J. A., Grosch, R., Lin, T. S., and Goetz, S. J.: Expansion of industrial logging in Central Africa, *Science*, 316, 1451, doi:10.1126/science.11410572007, 2007.
- Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z.-L., Levis, S., Sakaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization improvements and functional and structural advances in version 4 of the Community Land Model, *J. Adv. Modeling Earth Systems*, 3, 2011MS000045, doi:10.1029/2011MS000045, <http://www.cesm.ucar.edu/models/cesm1.0/clm/lawrence.clm4.james.2011.pdf>, 2011.
- Lehsten, V., Tansey, K., Balzter, H., Thonicke, K., Spessa, A., Weber, U., Smith, B., and Arneth, A.: Estimating carbon emissions from African wildfires, *Biogeosciences*, 6, 349–360, doi:10.5194/bg-6-349-2009, 2009.
- Lehsten, V., Arneth, A., Thonicke, K., and Spessa, A.: Tree-Grass coexistence in Savannas: testing two mechanisms, *J. Veg. Sci.*, in review, 2013.
- Levy, P. E., Cannell, M. G. R., and Friend, A. D.: Modelling the impact of future changes in climate, CO<sub>2</sub> concentration and land use on natural ecosystems and the terrestrial carbon sink, *Global Environ. Chang.*, 14, 21–30, 2004.
- Lewis, S. L., Lopez-Gonzalez, G., Sonke, B., Affum-Baffoe, K., Baker, T. R., Ojo, L. O., Phillips, O. L., Reitsma, J. M., White, L., Comiskey, J. A., Djuikouo, M. N., Ewango, C. E. N., Feldpausch, T. R., Hamilton, A. C., Gloor, M., Hart, T., Hladik, A., Lloyd, J., Lovett, J. C., Makana, J. R., Malhi, Y., Mbago, F. M., Ndangalasi, H. J., Peacock, J., Peh, K. S. H., Sheil, D., Sunderland, T., Swaine, M. D., Taplin, J., Taylor, D., Thomas, S. C., Votere, R., and Woll, H.: Increasing carbon storage in intact African tropical forests, *Nature*, 457, 1003–1006, 2009.
- Liousse, C., Guillaume, B., Grégoire, J. M., Mallet, M., Galy, C., Pont, V., Akpo, A., Bedou, M., Castéra, P., Dungall, L., Gardrat, E., Granier, C., Konaré, A., Malavelle, F., Mariscal, A.,



# The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Meybeck, M., Durr, H. H., and Vorosmarty, C. J.: Global coastal segmentation and its river catchment contributors: a new look at land-ocean linkage, *Global Biogeochem. Cy.*, 20, GB1S90, doi:10.1029/2005GB002540, 2006.

Mitchard, E. T. A., Saatchi, S. S., Woodhouse, I. H., Feldpausch, T. R., Lewis, S. L., Sonké, B., Rowland, C., and Meir, P.: Measuring biomass changes due to woody encroachment and deforestation/degradation in a forest-savanna boundary region of central Africa using multi-temporal L-band radar backscatter, *Remote Sens. Environ.*, 115, 2861–2873. doi:10.1016/j.rse.2010.02.022, 2011.

Mitchard, E. T. A., Saatchi, S. S., White, L. J. T., Abernethy, K. A., Jeffery, K. J., Lewis, S. L., Collins, M., Lefsky, M. A., Leal, M. E., Woodhouse, I. H., and Meir, P.: Mapping tropical forest biomass with radar and spaceborne LiDAR in Lopé National Park, Gabon: overcoming problems of high biomass and persistent cloud, *Biogeosciences*, 9, 179–191, doi:10.5194/bg-9-179-2012, 2012.

Nahlik, A. M. and Mitsch, W. J.: Methane emissions from tropical freshwater wetlands located in different climatic zones of Costa Rica, *Glob. Change Biol.*, 17, 1321–1334, 2011.

Nicholson, S. E.: The nature of rainfall variability over Africa on time scales of decades to millenia, *Global Planet. Change*, 26, 137–158, 2000.

Nicholson, S. E.: Climatic and environmental change in Africa during the last two centuries, *Clim. Res.*, 17, 123–144, 2001.

Nicholson, S. E. and Kim, J.: The relationship of the El Niño Southern Oscillation to African rainfall, *Int. J. Climatol.*, 17, 117–135, 1997.

Papale, D. and Valentini, R.: A new assessment of European forests carbon exchanges by eddy fluxes and artificial neural network spatialization, *Glob. Change Biol.*, 9, 525–535, 2003.

Patt, A. G. and Winkler, J.: Applying Climate Information in Africa: An Assessment of Current Knowledge, Written in support of NOAA, Boston University and the International Institute for Applied Systems Analysis (IIASA), 2007.

Peters, W., Krol, M. C., van der Werf, G. R., Houweling, S., Jones, C. D., Hughes, J., Schaefer, K., Masarie, K., Jacobson, A., Miller, J. B., Cho, C. H., Ramonet, M., Schmidt, M., Ciattaglia, L., Apadula, F., Heltai, D., Meinhardt, F., Di Sarra, A. G., Piacentino, S., Sferlazzo, D., Aalto, T., Hatakka, J., Ström, J., Haszpra, L., Meijer, H. A. J., Jordan, A., Rodó, X., Morgui, J. A., Vermeulen, A. T., Rozanski, K., Zimnoch, M., Manning, A., Leuenberger, M., Uglietti, C., Dolman, A. J., Ciais, P., Heimann, M., and Tans, P. P.: Seven years of recent



- European net terrestrial carbon dioxide exchange constrained by atmospheric observations, *Glob. Change Biol.*, 16, 1317–1337, 2009.
- Peylin, P., Bousquet, P., Le Queré, C., Sitch, S., Friedlingstein, P., McKinley, G., Gruber, N., Rayner, P., and Ciais, P.: Multiple constraints on regional CO<sub>2</sub> flux variations over land and oceans, *Global Biogeochem. Cy.*, 19, GB1011, doi:10.1029/2003GB002214, 2005.
- Phillips, O. L., Aragao, L., Lewis, S. L., Fisher, J. B., Lloyd, J., Lopez-Gonzalez, G., Malhi, Y., Monteagudo, A., Peacock, J., Quesada, C. A., van der Heijden, G., Almeida, S., Amaral, I., Arroyo, L., Aymard, G., Baker, T. R., Banki, O., Blanc, L., Bonal, D., Brando, P., Chave, J., de Oliveira, A. C. A., Cardozo, N. D., Czimczik, C. I., Feldpausch, T. R., Freitas, M. A., Gloor, E., Higuchi, N., Jimenez, E., Lloyd, G., Meir, P., Mendoza, C., Morel, A., Neill, D. A., Nepstad, D., Patino, S., Penuela, M. C., Prieto, A., Ramirez, F., Schwarz, M., Silva, J., Silveira, M., Thomas, A. S., ter Steege, H., Stropp, J., Vasquez, R., Zelazowski, P., Davila, E. A., Andelman, S., Andrade, A., Chao, K. J., Erwin, T., Di Fiore, A., Honorio, E., Keeling, H., Killeen, T. J., Laurance, W. F., Cruz, A. P., Pitman, N. C. A., Vargas, P. N., Ramirez-Angulo, H., Rudas, A., Salamao, R., Silva, N., Terborgh, J., and Torres-Lezama, A.: Drought sensitivity of the Amazon forest, *Science*, 323, 1344–1347, 2009.
- Poulter, B., Frank, D. C., Hodson, E. L., and Zimmermann, N. E.: Impacts of land cover and climate data selection on understanding terrestrial carbon dynamics and the CO<sub>2</sub> airborne fraction, *Biogeosciences*, 8, 2027–2036, doi:10.5194/bg-8-2027-2011, 2011.
- Prentice, I. C., Bondeau, A., Cramer, W., Harrison, S. P., Hickler, T., Lucht, W., Sitch, S., Smith, B., and Sykes, M. T.: Dynamic global vegetation modelling: quantifying terrestrial ecosystem responses to large-scale environmental change, in: *Terrestrial Ecosystems in a Changing World*, edited by: Canadell, J. G., Pataki, D. E., and Pitelka, L. F., Springer, Heidelberg, Berlin, 175–192, 2007.
- Prince, S. D., Wessels, K. J., Tucker, C. J., and Nicholson, S. E.: Desertification in the Sahel: a reinterpretation of a reinterpretation, *Glob. Change Biol.*, 13, 1308–1313, 2007.
- Purves, D. and Pacala, S.: Predictive models of forest dynamics, *Science*, 320, 1452–1453, 2008.
- Raymond, P. A., Hartmann, J., Sobek, S., Lauerwald, R., McDonald, C., Hoover, M., Butman, D., Striegl, R., Mayorga, E., Humborg, C., Kortelainen, P., Durr, H., Meybeck, M., Ciais, P., and Guth, P.: Global Carbon Dioxide Emissions from Inland Waters, *Nature*, in review, 2013.
- Reason, C. J. C.: Sensitivity of the southern African circulation to dipole sea-surface temperature patterns in the south Indian Ocean, *Int. J. Climatol.*, 22, 377–393, 2002.

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Reason, C. J. C. and Rouault, M.: ENSO-like decadal variability and South African rainfall, *Geophys. Res. Lett.*, 29, 1638–1642, 2002.
- Richard, Y., Trzaska, S., Roucou, P., and Rouault, M.: Modification of the Southern African rainfall variability/ENSO relationship since the late 1960s, *Clim. Dynam.*, 16, 883–895, 2000.
- 5 Roberts, G. J. and Wooster, M. J.: Fire detection and fire characterization over africa using meteosat SEVIRI, *IEEE T. Geosci. Remote*, 46, 1200–1218, 2008.
- Rödenbeck, C., Houweling, S., Gloor, M., and Heimann, M.: CO<sub>2</sub> flux history 1982–2001 inferred from atmospheric data using a global inversion of atmospheric transport, *Atmos. Chem. Phys.*, 3, 1919–1964, doi:10.5194/acp-3-1919-2003, 2003.
- 10 Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T. A., Salas, W., Zutta, B., Buermann, W., Lewis, S., Hagen, S., Petrova, S., White, L., Silman, M., and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents, *P. Natl. Acad. Sci. USA*, 108, 9899–9904, doi:10.1073/pnas.1019576108, 2011.
- Sanderson, M. G.: Biomass of termites and their emissions of methane and carbon dioxide: a global database, *Global Biogeochem. Cy.*, 4, 543–557, 1997.
- 15 Sankaran, M., Hanan, N. P., Scholes, R. J., Ratnam, J., Augustine, D. J., Cade, B. S., Gignoux, J., Higgins, S. I., Le Roux, X., Ludwig, F., Ardo, J., Banyikwa, F., Bronn, A., Bucini, G., Caylor, K. K., Coughenour, M. B., Diouf, A., Ekaya, W., Feral, C. J., February, E. C., Frost, P. G. H., Hiernaux, P., Hrabar, H., Metzger, K. L., Prins, H. H. T., Ringrose, S., Sea, W., Tews, J., Worden, J., and Zambatis, N.: Determinants of woody cover in African savannas, *Nature*, 438, 846–849, 2005.
- 20 Saunders, M. J., Kansiime, F., and Jones, M. B.: Agricultural encroachment: implications for carbon sequestration in tropical African wetlands, *Glob. Change Biol.*, 18, 1312–1321, 2012.
- Scheiter, S. and Higgins, S. I.: Impacts of climate change on the vegetation of Africa: an adaptive dynamic vegetation modelling approach (aDGVM), *Glob. Change Biol.*, 15, 2224–2246, 2009.
- 25 Scholes, R. J., Archibald, S. and von Maltitz, G.: Emissions from fire in Sub-Saharan Africa: the magnitude of sources, their variability and uncertainty, *Global Environ. Res.*, 15, 53–63, 2011.
- 30 Silva, J. M. N., Carreiras, J. M. B., Rosa, I., and Pereira, J. M. C.: Greenhouse gas emissions from shifting cultivation in the tropics, including uncertainty and sensitivity analysis, *J. Geophys. Res.*, 116, D20304, doi:10.1029/2011JD016056, 2011.



## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ulate climate-carbon cycle feedbacks: results from an atmosphere-ocean general circulation model, *Biogeosciences*, 6, 2099–2120, doi:10.5194/bg-6-2099-2009, 2009.

Tyson, P. D., Cooper, G. R. J., and McCarthy, T. S.: Millennial to multi-decadal variability in the climate of southern Africa, *Int. J. Climatol.*, 22, 1105–1117, 2002.

5 United Nations Statistical Division (UNSTAT): United Nations Demographic Yearbook 2008, United Nations Statistical Division (UNSTAT), New York, United States, 2010.

United Nations, Department of Economic and Social Affairs: Population Division, World Population prospect: The 2010 Revision, CD-ROM Edition, 2011.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arel-  
lano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004,  
10 *Atmos. Chem. Phys.*, 6, 3423–3441, doi:10.5194/acp-6-3423-2006, 2006.

van der Werf, G., Randerson, J., Giglio, L., Gobron, N. and Dolman, H.: Climate controls on the variability of fires in the tropics and subtropics, *Global Biogeochem. Cy.*, 22, GB3028, doi:10.1029/2007GB003122, 2008.

15 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

Van Drecht, G., Bouwman, A. F., Knoop, J. M., Beusen, A. H. W., and Meinardi, C. R.: Global modeling of the fate of nitrogen from point and nonpoint sources in soils, groundwater and surface water, *Global Biogeochem. Cy.*, 17, 1115, doi:10.1029/2003GB002060, 2003.

Van Wilgen, B. and Scholes, R. J.: The vegetation and fire regimes of Southern Hemisphere Africa, in: *Fire in Southern African Savannas: Ecological and Atmospheric Perspectives*, edited by: Andreae, M., Goldammer, J., and Lindsay, K., Witwatersrand University Press, Johannesburg, 27–46, 1997.

25 Veenendaal, E. M., Kolle, O., and Lloyd, J.: Seasonal variation in energy fluxes and carbon dioxide exchange for a broad-leaved semi-arid savanna (Mopane woodland) in southern Africa, *Glob. Change Biol.*, 10, 318–328, 2004.

Wang, S. Y. and Huang, D. J.: Assessment of greenhouse gas emissions from poultry enteric fermentation, *Asian Austral. J. Anim.*, 18, 873–878, 2005.

30 Weber, U., Jung, M., Reichstein, M., Beer, C., Braakhekke, M. C., Lehsten, V., Ghent, D., Kaduk, J., Viovy, N., Ciais, P., Gobron, N., and Rödenbeck, C.: The interannual variability

ity of Africa's ecosystem productivity: a multi-model analysis, *Biogeosciences*, 6, 285–295, doi:10.5194/bg-6-285-2009, 2009.

Williams, C. A.: Integration of remote sensing and modeling to understand carbon fluxes and climate interactions in Africa, in: *Ecosystem Function in Savannas: Measurement and Modeling at Landscape to Global Scales*, edited by: Hill, M. J. and Hanan, N., CRC Press, Boca Raton, FL, chap. 23, 327–346, 2010.

Williams, C. A. and Hanan, N. P.: ENSO and IOD teleconnections for African ecosystems: evidence of destructive interference between climate oscillations, *Biogeosciences*, 8, 27–40, doi:10.5194/bg-8-27-2011, 2011.

Williams, C. A., Hanan, N. P., Neff, J. C., Scholes, R. J., Berry, J. A., Denning, A. S., and Baker, D. F., Africa and the global carbon cycle, *Carbon Balance and Management*, 2, 1–13, 2007.

Williams, C. A., Hanan, N. P., Baker, I., Collatz, G. J., Berry, J., and Denning, A. S.: Interannual variability of photosynthesis across Africa and its attribution, *J. Geophys. Res.*, 113, G04015, doi:10.1029/2008JG000718, 2008.

Woodward, F. I. and Lomas, M. R.: Vegetation-dynamics – simulating responses to climate change, *Biol. Rev.*, 79, 643–670, 2004.

Woodward, F. I., Smith, T. M., and Emanuel, W. R.: A global land primary productivity and phytogeography model, *Global Biogeochem. Cy.*, 9, 471–490, 1995.

Wooster, M. J., Roberts, G., Perry, G. L. W., and Kaufman, Y. J.: Retrieval of biomass combustion rates and totals from fire radiative power observations: FRP derivation and calibration relationships between biomass consumption and fire radiative energy release, *J. Geophys. Res.*, 110, D24311, doi:10.1029/2005JD006318, 2005.

Yu, K. W., Faulkner, S. P., and Baldwin, M. J.: Effect of hydrological conditions on nitrous oxide, methane, and carbon dioxide dynamics in a bottomland hardwood forest and its implication for soil carbon sequestration, *Glob. Change Biol.*, 14, 1–15, 2008.

Zaehle, S. and Friend, A. D.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation and sensitivity to parameter estimates, *Global Biogeochem. Cy.*, 24, GB1005, doi:10.1029/2009GB003521, 2010.

Zeng, N.: Glacial-interglacial atmospheric CO<sub>2</sub> change – the glacial burial hypothesis, *Adv. Atmos. Sci.*, 20, 677–673, 2003.

Zeng, N., Mariotti, A., and Wetzels, P.: Terrestrial mechanisms of interannual CO<sub>2</sub> variability, *Global Biogeochem. Cy.*, 19, GB1016, doi:10.1029/2004GB002273, 2005.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ziehn, T., Scholze, M., and Knorr, W.: Regionalisation of the key carbon storage parameter within the carbon cycle data assimilation system (CCDAS), EGU General Assembly, Vienna, Austria, 2010.

- 5 Zomer, R. J., Trabucco, A., Coe, R., and Place, F.: Trees on Farm: Analysis of Global Extent and Geographical Patterns of Agroforestry, ICRAF Working Paper no. 89, World Agroforestry Centre, Nairobi, Kenya, 2009.

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The full greenhouse gases budget of Africa

R. Valentini et al.

**Table 1.** GHGs emission sources for five different African Regions.

Regions	Emission Sources (Pg CO <sub>2</sub> -eq)						
	Anthropogenic fossil fuel emissions	Per capita* emissions (MgCO <sub>2</sub> -eq)	Agriculture	LUCF	Total Emissions	Removals	Net Emissions
North Africa	0.3437	2.1	0.1198	0.0506	0.5141	0.0384	0.4757
Eastern Africa	0.1565	0.8	0.2794	0.2090	0.6448	0.4256	0.2193
Western Africa	0.2888	1.3	0.3070	0.3855	0.9813	0.5128	0.4685
Central Africa	0.02906	0.37	0.06683	0.45926	0.55514	1.18067	−0.62552
Southern Africa	0.43165	8.48	0.05483	0.01085	0.49734	0.08083	0.41651
Total	1.2497		0.8279	1.1152	3.1928	2.2383	0.9544
Total by Bombelli et al.	0.938		0.465	1.495	2.860	2.505	0.359

\* Per capita values are estimated on the base of UN World population prospect, the 2010 revision.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏮](#)
[⏭](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




## The full greenhouse gases budget of Africa

R. Valentini et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Table 2.** Different estimates of land cover across various products derived by space observation (GlobCover) and inventories (FAO).

	GlobCover2009 (Mha)	GlobCover2006 (Mha)	GLC2000 (Mha)	FAO FRA (Mha)	Range (% of median)
Croplands (including mosaics of grasslands and forests)	468.705	499.298	477.281		6.4
Forests (including mosaics of forests and woodlands)	740.561	836.816	638.152	674.419	26.8
Shrublands/grasslands (including savannas and sparse vegetation)	638.930	768.972	880.877		14.8
Flooded vegetation	43.833	51.946	3.410		110.7
Bare soil	991.490	1061.068	1047.134		6.6
Water bodies	28.189	28.866	28.298		2.4
Others	2.961	3.411	5.868		85.2
TOTAL	2914.670	3250.378	3081.021		10.9

**Table 3.** Biomass statistics and extent according to GlobCover 2009 land cover classes.

Globcover class name	LCCS class definition biomass (Mg ha <sup>-1</sup> )	Mean deviation (Mg ha <sup>-1</sup> )	Standard of variation	Coefficient extent (km <sup>2</sup> )	Area
20	Mosaic cropland (50–70 %)/vegetation (grassland/shrubland/forest) (20–50 %)	2.25	12.78	5.58	1 193 039
30	Mosaic vegetation (grassland/shrubland/forest) (50–70 %)/cropland (20–50 %)	39.92	71.02	1.77	2 449 681
40	Closed to open (> 15 %) broadleaved evergreen and/or semi-deciduous forest (> 5 m)	200.43	95.54	0.47	1 879 612
50	Closed (> 40 %) broadleaved deciduous forest (> 5 m)	79.21	68.61	0.86	1 166 758
60	Open (15–40 %) broadleaved deciduous forest/woodland (> 5 m)	35.83	43.25	1.20	2 434 440
70	Closed (> 40 %) needle-leaved evergreen forest (> 5 m)	19.87	47.45	2.38	251
90	Open (15–40 %) needle-leaved deciduous or evergreen forest (> 5 m)	71.96	70.09	0.97	34 894
100	Closed to open (> 15 %) mixed broadleaved and needleleaved forest	8.14	25.61	3.14	662
110	Mosaic forest or shrubland (50–70 %) and grassland (20–50 %)	3.08	13.72	4.45	1 363 470
120	Mosaic grassland (50–70 %) and forest or shrubland (20–50 %)	21.82	35.82	1.64	344 184
130	Closed to open (> 15 %) (broadleaved or needleleaved, evergreen or deciduous) shrubland (< 5 m)	15.58	32.60	2.09	2 186 892
140	Closed to open (> 15 %) herbaceous vegetation (grassland, savannas or lichens/mosses)	0.89	9.08	10.2	1 857 738
150	Sparse (< 15 %) vegetation	0.02	1.28	64	476 781
160	Closed (> 40 %) broadleaved forest regularly flooded, fresh water	192.70	101.09	0.52	363 910
170	Closed (> 40 %) broadleaved semi-deciduous and/or evergreen forest regularly flooded, saline water	25.74	56.22	2.18	19 140
180	Closed to open (> 15 %) grassland or shrubland or woody vegetation on regularly flooded or waterlogged soil, fresh, brackish or saline water	22.91	37.41	1.63	88 089

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





# The full greenhouse gases budget of Africa

R. Valentini et al.

**Table 5.** DGVM average carbon pools (Pg C) and fluxes (PgCyr<sup>-1</sup>) over the period 1990–2009, NBP is defined as NBP = NPP – RH – wildfire Flux (\* note LPJ has a grazing term in the NBP calculation).

DGVM	VegC	SoilC	NPP	GPP	Wildfire	NBP
CLM4CN	96.27	87.48	9.25	25.33	0.60	0.16
Hyland	181.44	240.95	20.46	40.91	–	1.09
LPJ	82.16	94.93	11.18	24.91	0.64	0.24
LPJ-GUESS	92.14	103.36	12.21	20.61	1.47	0.38
OCN	106.03	169.78	11.26	25.92	–	0.30
ORCHIDEE	98.69	236.52	15.77	32.24	–	0.60
SDGVM	66.70	193.48	12.70	24.07	1.27	0.44
TRIFFID	127.68	120.56	13.89	28.95	–	0.50
VEGAS	129.08	259.53	12.7	30.46	–	–0.01
Mean Model	108.91	167.40	13.27	28.16	1.00	0.41
St. Dev.	33.67	68.28	3.25	5.94	0.44	0.31

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 6.** Sources (+) and sinks (–) of carbon ( $\text{TgCyr}^{-1}$ ) from different types of land use in regions of Sub-Saharan Africa (average annual net flux).

	Period	Croplands	Pasture	Afforest.	Indust. harvest	Fuelwood harvest	Shifting cult.	Total
All sub-Saharan Africa	1980–1989	101	25	–3	23	0	167	314
	1990–1999	154	65	–2	32	0	67	317
	2000–2009	164	34	–2	38	0	89	323
	1990–2009	159	49	–2	35	0	78	320
East Africa	1980–1989	26	10	–1	3	0	11	47
	1990–1999	27	25	–1	5	0	6	62
	2000–2009	45	6	0	6	0	2	58
	1990–2009	36	16	–1	5	0	4	60
West Africa	1980–1989	42	9	0	9	0	18	77
	1990–1999	93	3	0	11	0	0	108
	2000–2009	93	1	–1	13	0	1	106
	1990–2009	93	2	–1	12	0	0	107
Southern Africa	1980–1989	17	7	–1	4	0	43	71
	1990–1999	31	35	–1	7	0	10	82
	2000–2009	22	26	0	8	0	40	96
	1990–2009	27	31	0	8	0	25	89
Central Africa	1980–1989	17	0	0	7	0	95	119
	1990–1999	3	1	0	9	0	51	64
	2000–2009	4	1	0	10	0	47	62
	1990–2009	3	1	0	10	0	49	63

# The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 7.** Fire emissions characteristics for Africa based on 1997–2011 GFED data.

	North of equator	South of equator
Mean annual emissions ( $\text{PgCyr}^{-1}$ )	$0.469 \pm 0.094$	$0.562 \pm 0.112$
Coefficient of variation (%)	15	9
Range ( $\text{PgC}$ )	0.221	0.199
Fuel consumption ( $\text{gCm}^{-2}$ )	371	446

# The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 8.** Emissions of  $\text{N}_2\text{O}$  associated to biogenic sources and fires in the African continent.

	$\text{N}_2\text{O}$ emissions at continental level ( $\text{Gg N}_2\text{O yr}^{-1}$ )	Contribution of single sources to total continental emissions (%)
Soil Emissions		
Natural ecosystems	$1173.2^a \pm (0.8-5.8)^g$	42.4 %
Agricultural practices and husbandry	$827.1^b \pm (2.2-3.9)^g$	29.9 %
Deforestation effect on soil (background flux)	$69.0^c \pm 0.67$	2.5 %
Other direct and indirect sources		
N depositions	$123.8^d$	4.5 %
N leaching derived	$117^e (9-566)^h$	4.2 %
Emissions from fire	$455^f$	16.4 %
Total emissions $\text{Tg N}_2\text{O yr}^{-1}$	2.76 (0.01–0.57)	
$\text{Tg CO}_2\text{-eq}$	824	

<sup>a</sup> Data derived from an empirical relationship between annual total emissions (64 ecosystems spanning from rain forest to seasonally dry ecosystems) and total annual rainfall (Thompson et al., 2013);

<sup>b</sup> modeled for Sub-Saharan Africa after Bouwman et al., 2002 (Castaldi et al., 2013), N input in agroecosystems derived by Potter et al. (2010), Edgar 4.1 data for natural emissions from countries or North Africa, based on IPCC methodology (Hickman et al., 2011);

<sup>c</sup> empirical equation base on published data (Thompson et al., 2013);

<sup>d</sup> N inputs derived from Dentener et al. (2006) and EFs from IPCC (2007);

<sup>e</sup> N export from Van Dreht et al. (2003) and IPCC (2007) EFs;

<sup>f</sup> Edgar 4.1;

<sup>g</sup> calculated error range;

<sup>h</sup> error range as reported by IPCC (2007).



# The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 9.** CH<sub>4</sub> fluxes from biogenic processes and fires. Total surfaces derived from Globcover.

	CH <sub>4</sub> fluxes at continental level Tg CH <sub>4</sub> yr <sup>-1</sup>	Contribution of single sources to total continental emissions (%)
<b>Net soil CH<sub>4</sub> exchange</b>		
Tropical humid forests	0.27 (0.16) <sup>a</sup>	
Seasonally dry forests	-0.21 (0.42) <sup>b</sup>	
Savannas	-1.43 (2.86) <sup>b</sup>	
Pastures	0.1 (0.74) <sup>b</sup>	
Croplands	-0.19 (0.38) <sup>b</sup>	
<b>Total soil Tg CH<sub>4</sub> yr<sup>-1</sup></b>	<b>-1.46 (3.01)</b>	<b>-2.08</b>
CH <sub>4</sub> emissions from forested wetlands	18.8 (7.2–26.4) <sup>c</sup>	26.85
CH <sub>4</sub> emissions from unforestet wetlands	15.6 (9.6–68) <sup>c</sup>	22.28
CH <sub>4</sub> emissions from rice paddy fields	4.4 (2.9–9.9) <sup>d</sup>	6.28
CH <sub>4</sub> emissions from termites	2.5 <sup>h</sup> (0.9 <sup>h</sup> )–6.9 <sup>e</sup>	3.78–9.80
CH <sub>4</sub> emissions from animal husbandry and wildlife	17.83 <sup>f</sup>	25.46
CH <sub>4</sub> emissions from fires	8 (5–11) <sup>g</sup>	11.42
<b>Total Tg CH<sub>4</sub> yr<sup>-1</sup></b>	<b>65.65 (62.24)</b>	
<b>Pg CO<sub>2</sub>-eq yr<sup>-1</sup></b>	<b>1.64</b>	

<sup>a</sup> EFs from Castaldi, personal communication, 2013., Yu et al. (2008);

<sup>b</sup> EFs from Castaldi et al. (2006);

<sup>c</sup> EFs median of yearly ecosystem emissions from Bartlett et al. (1988), Matthews and Fung (1987), Devol et al. (1988), Smith et al. (2000), Melack et al. (2004), Hadi et al. (2005), Nahlik e Mistch (2011), in brackets scale up based on 25° and 75° percentile of available data, land cover from <https://secure.worldwildlife.org>;

<sup>d</sup> median and percentiles EFs derived from Aselmann and Crutzen (1989), Banker et al. (1995), Hadi et al. (2005), total hectares from FAOSTAT (2010) data;

<sup>e</sup> EFs from Sanderson (1997) Globcover classes 20–70, 90–140, uncertainty is in the range of 1 %;

<sup>f</sup> populations from FAOSTAT (2010), EFs from Crutzen et al. (1986) and Wang and Huang (2005);

<sup>g</sup> Delmas et al. (1992);

<sup>h</sup> based on the method proposed in Kirschke et al. (2013).

**Table 10.** Synthesis of current estimates of GHG and carbon fluxes and pools in Africa. All these values do not include fossil fuel emissions. Reference periods are indicated in the footnotes.

	Inventory data	Ecosystem Fluxes <sup>a</sup>	Models (DGVM) <sup>b</sup>	Atmospheric inversions <sup>c</sup>
<b>Area(Mha)</b>				
Forest	708 ± 89 <sup>d</sup>			
Savannas	769 ± 109 <sup>d</sup>			
Crops	477 ± 14 <sup>d</sup>			
<b>Stocks (PgC)</b>				
Forest Biomass	63 ± 24 <sup>d</sup>			
<b>Carbon Fluxes(PgCyr<sup>-1</sup>)</b>				
Gross Primary Production	31.7 ± 6.6	24.3 ± 2.9 <sup>a</sup>	28.2 ± 5.9 <sup>b</sup>	
Forest	18.8 ± 5.3 <sup>e</sup>			
Savannas	12.9 ± 1.9 <sup>e</sup>			
Net Primary Production	14.3 ± 3.7		13.3 ± 3.3 <sup>b</sup>	
Forests	8.5 ± 2.4 <sup>f</sup>			
Savannas	5.8 ± 2.8 <sup>g</sup>			
Autothropic respiration	17.4 ± 7.6 <sup>h</sup>			
Forests	11.1 ± 6.7 <sup>i</sup>			
Savannas	6.3 ± 10.0			
Heterotrophic respiration	12.0 ± 11.3		12.1 ± 1.1 <sup>b</sup>	
Forests	7.1 ± 4.7 <sup>i</sup>			
Savannas	4.8 ± 10.3 <sup>j</sup>			
Total respiration	29.3 ± 13.6	19.0 ± 1.4 <sup>a</sup>		
<b>Net Ecosystem Production</b>	-2.3 ± 1.2	-2.9 ± 1.3 <sup>a</sup>	-1.2 ± 3.5 <sup>b</sup>	
<b>Land Use Emissions</b>	0.51 ± 0.10			
Deforestation (PgCyr <sup>-1</sup> )	0.32 ± 0.05 <sup>k</sup>			
Forest Degradation (PgCyr <sup>-1</sup> )	0.19 ± 0.09 <sup>f</sup>			
<b>Wildfires emissions (PgCyr<sup>-1</sup>)</b>	1.03 ± 0.22 <sup>l</sup>			

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 10.** Continued.

	Inventory data	Ecosystem Fluxes <sup>a</sup>	Models (DGVM) <sup>b</sup>	Atmospheric inversions <sup>c</sup>
<b>Lateral transport (PgCyr<sup>-1</sup>)</b>				
Carbon in trade	0.000008 <sup>m</sup>			
DIC	0.029 <sup>n</sup>			
DOC	0.018 <sup>n</sup>			
POC	0.016 <sup>n</sup>			
<b>Net Biome Production (PgCyr<sup>-1</sup>)</b>	$-0.74 \pm 1.19$	$-1.34 \pm 1.32^a$	$-0.41 \pm 0.31^b$	$0.05 \pm 0.28^c$
<b>Net Biome Production (PgCO<sub>2</sub>yr<sup>-1</sup>)</b>	$-2.66 \pm 4.28$	$-4.82 \pm 4.75$	$-1.48 \pm 1.12$	$0.18 \pm 1.01$
<b>Other GHG fluxes</b>				
CH <sub>4</sub> fluxes (TgCH <sub>4</sub> yr <sup>-1</sup> )	$70.03 \pm 35^o$			
CH <sub>4</sub> fluxes (PgCO <sub>2</sub> -eq)	$1.75 \pm 0.85^o$			
N <sub>2</sub> O fluxes (TgN <sub>2</sub> Oyr <sup>-1</sup> )	$2.77 \pm 1.38^o$			
N <sub>2</sub> O fluxes (PgCO <sub>2</sub> -eqyr <sup>-1</sup> )	$0.82 \pm 0.41^o$			
Total other GHG fluxes	$2.57 \pm 1.28$			
NMHC (PgCyr <sup>-1</sup> )	$0.16 \pm 0.04^p$			
NMHC (PgCO <sub>2</sub> yr <sup>-1</sup> )	$0.59 \pm 0.12$			

<sup>a</sup> Reference period: 1982–2008.

<sup>b</sup> Reference period: 1990–2010.

<sup>c</sup> Reference period: 1996–2004. Inversion systems: Jena.s96, LSC2.var, LSCE.an, and Ctracker.EU.

<sup>d</sup> Median value and error estimated as 95 and 0.5 percentile.

<sup>e</sup> Derived by NPP/0.45 error derived by uncertainty in area estimation.

<sup>f</sup> Estimated by this study.

<sup>g</sup> Derived by Bombelli et al. (2009) and Lehsten et al. (2009).

<sup>h</sup> Derived by GPP-NPP.

<sup>i</sup> Derived by Malhi et al. (2009).

<sup>j</sup> Derived by Gilmanov et al. (2010).

<sup>k</sup> Reference period: 1990–2009.

<sup>l</sup> Reference period: 1997–2011.

<sup>m</sup> Derived by Davis et al. (2011).

<sup>n</sup> Reference period: 1980–2000.

<sup>o</sup> the time period of input data was centered on 2000, based on availability of most input data, which spanned from the middle of 1990s' to 2009.

<sup>p</sup> Derived by Kesselmeier et al. (2002).

**BGD**

10, 8343–8413, 2013

## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 1.** Histograms of biomass for different land cover aggregated classes (see Table 4).

## BGD

10, 8343–8413, 2013

### The full greenhouse gases budget of Africa

R. Valentini et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



## The full greenhouse gases budget of Africa

R. Valentini et al.



**Fig. 2.** Year to year variations in the net CO<sub>2</sub> flux of Africa seen by five recent inversions. The CO<sub>2</sub> flux includes land use CO<sub>2</sub> emissions and other ecosystem CO<sub>2</sub> fluxes, and fossil CO<sub>2</sub> emissions as well (<http://www.carboscope.eu/>).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 3.** (a) Distribution of mean CO<sub>2</sub> fluxes averaged across the five recent inversions analyzed. (b) shows the spread of the inversions (defined from the estimator of the standard deviation). Inversions have a different spatial resolution, and the fluxes have been interpolated on a common grid. Negative values are atmospheric CO<sub>2</sub> sinks (<http://www.carboscope.eu/>).

# The full greenhouse gases budget of Africa

R. Valentini et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Fig. 4.** Average Net Primary Productivity (NPP) for each DGVM and model mean over the period 1990–2009, units,  $\text{kg C m}^{-2} \text{yr}^{-1}$ .



# The full greenhouse gases budget of Africa

R. Valentini et al.




**Fig. 5.** Average Net Biome Productivity (NBP) for each DGVM and model mean over the period 1990–2009, units,  $\text{kg C m}^{-2} \text{yr}^{-1}$ .

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏮](#)
[⏭](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


# The full greenhouse gases budget of Africa

R. Valentini et al.



**Fig. 6.** Standardized interannual variability of gross primary productivity, defined according to Weber et al., 2009 as  $\frac{IAV_i - \overline{IAV}}{\sigma_{IAV}}$ , from five modeling exercises. SiB3 results were reported in Williams et al. (2008) and cover for 1982–2003 and the remaining four were reported in Weber et al. (2009) spanning 1982–2008, reproduced with permission. Values greater than 0 indicate above average interannual variability.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






**Fig. 7.** Spatial distribution across Africa of the slope of least-squares linear regressions of SiB3 modeled **(a, b)** annual net photosynthesis ( $P_{\text{net}}$ ) or CRU/TRMM **(c, d)** annual rainfall on climate indices. Each column corresponds to analyses with a particular index with the left column **(a, c)** for the multivariate ENSO index, and the right column **(b, d)** for the Indian Ocean Dipole Mode Index (IOD). Results are based on Williams et al. (2008).

# The full greenhouse gases budget of Africa

R. Valentini et al.



**Fig. 8.** Spatial distributions of ORCHIDEE modeled trends in annual GPP, TER, and NBP from 1980 to 2002 and of their main driving factor. Reproduced with permission from Ciais et al. (2009).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

