

A U-Pb and Rb-Sr isotope study of a migmatite area Scoresby Sund, East Greenland

Doctoral Thesis

Author(s):

Oberli, Felix

Publication date:

1976

Permanent link:

<https://doi.org/10.3929/ethz-a-000101334>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Dissertation No. 5702

A U-PB AND RB-SR ISOTOPE STUDY OF A MIGMATITE AREA

SCORESBY SUND, EAST GREENLAND

A Dissertation submitted
to the
SWISS FEDERAL INSTITUTE OF TECHNOLOGY
ZURICH

for the degree
Doctor of Natural Sciences

by

FELIX OBERLI

dipl.sc.nat.ETH

born on December 10, 1946

from Rüderswil BE and Adliswil ZH, Switzerland

Accepted on the recommendation of

Prof. Dr. R.H. Steiger, Zurich

Prof. Dr. P. Signer, Zurich

1976

Abstract

A 2000 m thick massive granodioritic sheet and its migmatized metasedimentary cover, situated in the southern part of the East Greenland fold belt, have been chosen for a study of isotope systems in migmatites. Rb-Sr whole-rock and zircon U-Pb data obtained from various granitic and metasedimentary rock phases show a pronounced scatter indicative of complex open-system behaviour. The disturbances of these isotopic systems are believed to result from a polymetamorphic history of the area including a high-grade Caledonian event. The latter is dated by a Pb-Pb monazite age of 434 ± 5 m.y. as well as by a Rb-Sr mineral isochron age of 410 ± 8 m.y. A concordant U-Pb monazite age of 423 ± 4 m.y. marks the intrusion of post-kinematic granitic dikes.

The data scatter on a U-Pb concordia diagram resulting from conventionally separated zircon fractions of a complex migmatitic rock leads to ambiguous or even wrong age information. However, bulk zircon separates produce a linear data pattern and permit - in combination with U and Pb concentration data - to assign a Caledonian age to the migmatization.

The thesis is supplemented by a statistical investigation of mass spectrometer data.