

Multiquadric Pre-Wavelets on Non- Equally Spaced Centres

Report**Author(s):**

Buhmann, Martin Dietrich

Publication date:

1994-07

Permanent link:

<https://doi.org/10.3929/ethz-a-004284079>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Originally published in:

SAM Research Report 1994-06

Multiquadric Pre-Wavelets on Non-Equally Spaced Centres

M.D. Buhmann

Research Report No. 94-06

July 1994

Seminar für Angewandte mathematik

Eidgenössische Technische Hochschule

CH-8092 Zürich

Switzerland

Multiquadric Pre-Wavelets on Non-Equally Spaced Centres

M.D. Buhmann

Seminar für Angewandte mathematik
Eidgenössische Technische Hochschule
CH-8092 Zürich
Switzerland

Research Report No. 94-06 July 1994

Abstract

In this paper, we identify univariate prewavelets on spaces spanned by translates of multiquadric functions and other radial basis functions with *non-equally spaced* centres (or “knots”). Although the multiquadric function and its relations are our prime examples, the theory is sufficiently broad to admit prewavelets from other radial basis function spaces as well.

AMS(MOS) Subject Classification: 41A15, 41A30, 42C15, 65D15

2 June 1994, revised 18 October 1994

1. Introduction.

The theme of this work is the construction of univariate prewavelets that are generated by translates of a radial basis function, where the translates are defined by a sequence of non-equally spaced centres. We first explain why radial basis functions are useful for the construction of prewavelets.

The purpose of wavelet and prewavelet decompositions of univariate square-integrable functions is to decompose those functions both into their different frequency components, much like Fourier analysis does, as well as with respect to time, which is not possible with standard Fourier analysis. Every such decomposition of a given square-integrable function f , say, into a time/frequency series starts with establishing an initial approximation f_k to f from a linear space V_k that is an element of a nested sequence of spaces

$$V_0 \subset V_1 \subset V_2 \subset \cdots \subset L^2(\mathbb{R}) \quad (1.1)$$

from which the (pre)wavelets are taken. One requires that

$$\overline{\bigcup_{j=0}^{\infty} V_j} = L^2(\mathbb{R}), \quad (1.2)$$

so that every $f \in L^2(\mathbb{R})$ can be approximated arbitrarily closely in the $L^2(\mathbb{R})$ norm by an $f_k \in V_k$ if k is sufficiently large. Then, orthogonal complements $W_j \subset V_{j+1}$ of V_j , $j \in \mathbb{Z}_+$ are studied. In other words, one seeks spaces $W_j \subset V_{j+1}$ such that $W_j \perp V_j$ and $V_j \oplus W_j = V_{j+1}$. In fact, the sequence of spaces (1.1) is usually biinfinite, but for the results of this paper it is more appropriate to restrict it to an infinite sequence.

By constructing the W_j in the outlined fashion, we get an orthogonal decomposition of f_k into an $f_0 \in V_0$ plus a series of $g_\ell \in W_\ell$, $\ell = 0, 1, \dots, k-1$. Therefore a decomposition of any square-integrable f into mutually orthogonal components, corresponding to the different frequency components in Fourier analysis is found. This can be viewed as a spatial orthogonal decomposition

$$L^2(\mathbb{R}) = V_0 + W_0 + W_1 + W_2 + \cdots. \quad (1.3)$$

The *prewavelets* are functions that generate such W_j . It is preferable that those prewavelets have local support or that they decay, so that the expansion of f 's components in each W_j is a localized decomposition (with respect to time). The prewavelets are called wavelets if they are *orthonormal bases* of the W_j s, but we are only dealing with prewavelets here. It is always the function $f_k \in V_k$, which may for instance have been obtained by quasi-interpolation (e.g., Buhmann, 1993c), that is actually decomposed computationally by the so-called fast wavelet transform. Therefore it is

important to the effectiveness of the decomposition that the spaces V_k allow f_k to be a sufficiently accurate approximation to f , even if k is moderate. This is the point where the efficacy of radial basis functions can be exploited.

We want to explain this point further. Radial basis function methods are generally known to be useful and accurate for the approximation of functions; cf., e.g., the review paper by the author (1993a) which also includes an extensive list of references to other people's work. The key idea is to approximate from a space spanned by translates of a single function ϱ , usually of global support, where the translates take the form $\varrho(|\cdot - x_j|)$ and the x_j are given "centres". They were first introduced as interpolatory schemes but, because they were found to provide high quality approximants, they are now used for many different approximation tools. In this paper, we submit another one in the shape of prewavelets. Radial function methods are easy to implement, and so is the approach proposed here, as we shall see, although special considerations are needed for the matrix computations involved and evaluation of approximants if the amount of data is large (Powell, 1992). Among all radial functions currently in use, the multiquadric radial function $\varrho(r) = \sqrt{r^2 + \gamma^2}$ is probably best understood, both theoretically and from a practical point of view, and also it is the one most frequently used, partly by virtue of the variable real parameter γ . Therefore, we shall focus on it in this article.

Up to now, the most frequent choice for the aforementioned V_j are spline spaces with equally spaced knots (Chui and Wang, 1992, for example), i.e. spaces of piecewise polynomials with knotspacing 2^{-j} for the V_j . These spaces are spanned by B-splines which have compact support. This works well except that the resulting approximations are always of limited smoothness. However, if one takes spaces V_j spanned by the multiquadric functions centred at $2^{-j}k$, i.e. $\sqrt{(\cdot - 2^{-j}k)^2 + \gamma^2}$, $k \in \mathbb{Z}$, then the known theory of radial functions (Buhmann, 1993c, for example) provides essentially the same approximation order as splines and infinite differentiability as well. Sometimes better (spectral) approximation orders are obtained too (Buhmann and Dyn, 1991), but this depends strongly on f 's smoothness. The price of these advantages is giving up compact support of the basis functions, but there are generating functions for the V_j from radial basis function spaces that diminish very fast toward infinity. Such generating functions in the univariate multiquadric setting with scattered centre points are found in (Powell, 1990), so that a theory of prewavelets on non-equally spaced data may be established for radial basis functions. Therein lies the purpose of this work. We therefore study prewavelets that stem from nested spaces V_j spanned by a radial basis function, using in particular the multiquadric function and allowing the centres to be scattered. A very general account of generating prewavelets from *shift-invariant*

spaces (including radial function spaces with gridded centres) is given by de Boor, DeVore and Ron (1992) but their technique cannot be applied to scattered data. The main focus of that paper lies in compactly supported prewavelets (mainly from box-spline spaces) which do not exist in spaces spanned by radial functions, other than $\phi(r) = r^{2k+1}$, $k \in \mathbb{Z}_+$, in one dimension.

We will show how to find prewavelets $\psi_j \in V_1$ that are orthogonal to V_0 and generate

$$W_0 = \left\{ \sum_{j=-\infty}^{\infty} c_j \psi_j \mid c = \{c_j\}_{j=-\infty}^{\infty} \in \ell^2(\mathbb{Z}) \right\}, \quad (1.4)$$

with $W_0 \perp V_0$, where V_0 and V_1 are spaces generated by non-equally spaced translates of a radial basis function such as the multiquadric function $\varrho(r) = \sqrt{r^2 + \gamma^2}$. Here, $V_0 \subset V_1$ is obtained by demanding that these two spaces be produced by way of two sets of scattered centres, one a subset of the other. The “finer” set generates V_1 , of course. Our principal example is the multiquadric radial function but the setup in this work is sufficiently general to admit other choices of radial functions.

Once it is demonstrated how V_0 , V_1 and the prewavelets that generate W_0 are found, the same principles can be followed to establish a whole sequence of nested subspaces V_j of $L^2(\mathbb{R})$ and to decompose the space as in (1.3). Incidentally, wavelets can be obtained from those prewavelets by a standard orthonormalization procedure.

Prior work to this paper are the papers about radial basis function prewavelets due to (Buhmann, 1993b), (Chui, Ward and Stöckler, 1994) and (Micchelli, Rabut and Utreras, 1992) which treat the case of equally spaced centres only (some of the preliminary results in the first paper mentioned also apply to scattered centres, and, on the other hand, the results in the third paper apply to the multidimensional setting too).

There now follows a section with preliminaries where the spaces V_0 and V_1 are defined. Thereafter, there is a section containing the principal results of this paper, where the prewavelets are explicitly constructed and some of their fundamental properties established. Several examples to which our analysis applies are stated at the end of that section as well.

2. Preliminaries: The spaces V_0 and V_1 .

Let two sequences of real numbers

$$\mathbf{x} = \left\{ \cdots < x_{-1} < x_0 < x_1 < x_2 < \cdots \right\}$$

and

$$\boldsymbol{\tau} = \left\{ \cdots < \tau_{-1} < \tau_0 < \tau_1 < \tau_2 < \cdots \right\}$$

with $\tau_{2i-1} = x_i$, $i \in \mathbb{Z}$, be given. We assume that these points satisfy the boundedness conditions

$$\kappa := \sup_{i \in \mathbb{Z}} (\tau_{i+1} - \tau_i) < \infty, \quad \delta := \inf_{i \in \mathbb{Z}} (\tau_{i+1} - \tau_i) > 0.$$

We let $\{B_j^c\}_{j=-\infty}^{\infty}$ and $\{B_j^f\}_{j=-\infty}^{\infty}$ be the sequences of B-splines of degree n on the knot-sequences \mathbf{x} and $\underline{\tau}$, with supports $[x_j, x_{j+n+1}]$ and $[\tau_j, \tau_{j+n+1}]$, respectively, normalized to form a partition of unity. Let $\phi \in C(\mathbb{R})$ with $\phi(x) \geq 0$, $x \in \mathbb{R}$,

$$\phi(x) = O(|x|^{-n-1-\varepsilon}), \quad x \rightarrow \pm\infty, \quad \text{for some positive } \varepsilon, \quad (2.1)$$

$\hat{\phi}(t) > 0$, $t \in \mathbb{R}$, and $\int_{-\infty}^{\infty} \phi = 1$ be given. For instance, we may take $\phi(t) = \frac{1}{2}\varrho''(t)$ where $\varrho(t)$ is the multiquadric function $\sqrt{t^2 + \gamma^2}$, $n = 1$ and $\varepsilon = 1$. We consider the functions

$$\left. \begin{aligned} C_j &:= B_j^c * \phi, \quad j \in \mathbb{Z}, \\ F_j &:= B_j^f * \phi, \quad j \in \mathbb{Z}. \end{aligned} \right\} \quad (2.2)$$

In (2.2), $*$ denotes convolution. We immediately record the two important identities

$$\left. \begin{aligned} \sum_{j=-\infty}^{\infty} |F_j(x)| &= \sum_{j=-\infty}^{\infty} F_j(x) \equiv 1, \\ \sum_{j=-\infty}^{\infty} |C_j(x)| &= \sum_{j=-\infty}^{\infty} C_j(x) \equiv 1. \end{aligned} \right\} \quad (2.3)$$

They are a result of the fact that the B-splines form a partition of unity and that the integral of ϕ is one. With the example of ϕ as suggested above and $n = 1$, C_j and F_j are second divided differences of the multiquadric function with respect to the sequences \mathbf{x} and $\underline{\tau}$, respectively (Powell, 1990). We define

$$\left. \begin{aligned} V_0 &:= \left\{ \sum_{j=-\infty}^{\infty} c_j C_j \mid c = \{c_j\}_{j=-\infty}^{\infty} \in \ell^2(\mathbb{Z}) \right\}, \\ V_1 &:= \left\{ \sum_{j=-\infty}^{\infty} c_j F_j \mid c = \{c_j\}_{j=-\infty}^{\infty} \in \ell^2(\mathbb{Z}) \right\}. \end{aligned} \right\}$$

We note that if ϕ is the $\underline{\delta}$ -distribution, then we get the spline spaces considered in (Buhmann and Micchelli, 1992). We shall now prove two elementary facts about these spaces. In the next section, we shall proceed to constructing the prewavelets that generate the orthogonal complement W_0 as in (1.4) of V_0 in V_1 . We begin with

Proposition 1. *The space V_0 is contained in the space V_1 .*

Proof. This proof has already been presented in (Buhmann, 1993b), but we repeat it here for the convenience of the reader. Let $c = \{c_j\}_{j=-\infty}^{\infty} \in \ell^2(\mathbb{Z})$. Hence the function $f \in V_0$ defined pointwise by

$$f(x) = \sum_{j=-\infty}^{\infty} c_j \int_{-\infty}^{\infty} B_j^c(y) \phi(x-y) dy, \quad x \in \mathbb{R}, \quad (2.4)$$

is defined by an absolutely convergent series and bounded in modulus by $\|c\|_{\infty}$, because $\phi(x-y) \geq 0$ and the B-splines form a partition of unity. Now recall that there exists a positive constant λ that depends only on n (de Boor, 1976, p. 37) so that

$$\lambda \|c\|_p \leq \left\| \sum_{j=-\infty}^{\infty} c_j \left(\frac{n+1}{x_{j+n+1} - x_j} \right)^{1/p} B_j^c \right\|_p \leq \|c\|_p, \quad 1 \leq p \leq \infty, \quad (2.5)$$

where $\|\cdot\|_p$ denotes the $\ell^p(\mathbb{Z})$ or the $L^p(\mathbb{R})$ norm as is appropriate. If only $p = \infty$ is admitted into (2.5), λ is bounded below by $2^{-n-2} \sqrt{\pi n} / (2n+1)$ (this is a recent result due to Scherer; de Boor, private communication). It follows from (2.5) with $p = 2$ and from the Cauchy–Schwarz inequality that we may exchange summation and integration in (2.4). It is the same as

$$f(x) = \int_{-\infty}^{\infty} \sum_{j=-\infty}^{\infty} c_j B_j^c(y) \phi(x-y) dy$$

which can be re-written by the Curry–Schoenberg theorem (de Boor, 1976, p. 36) as

$$f(x) = \int_{-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \tilde{c}_j B_j^f(y) \phi(x-y) dy$$

with $\tilde{c} \in \ell^2(\mathbb{Z})$. Hence, reverting back to the form of f with the sum outside the integral, $f \in V_1$, as required. ■

It will be convenient in the sequel to write the spaces V_0 and V_1 as $\mathcal{S}_{\mathbf{x},n,\phi}$ and $\mathcal{S}_{\underline{\tau},n,\phi}$, respectively, so that $\mathcal{S}_{\mathbf{x},n,\underline{\delta}}$, for example, is the space of splines of degree n on the knot sequence \mathbf{x} that are in $L^2(\mathbb{R})$ (by (2.5) with $p = 2$).

Of course it is relevant to our discussion that (1.2) can be satisfied in the present set-up. Indeed, we have

Theorem 2. *Let $\{\underline{\tau}^k\}_{k=0}^{\infty}$ be a sequence of nested knot sequences in \mathbb{R} , i.e. $\underline{\tau}^0 \subset \underline{\tau}^1 \subset \dots$, where the nesting is in the same sense as in $\mathbf{x} \subset \underline{\tau}$, that satisfy*

$$\kappa_k := \sup_{j \in \mathbb{Z}} (\tau_{j+1}^k - \tau_j^k) \rightarrow 0, \quad k \rightarrow \infty,$$

and become dense in \mathbb{R} for $k \rightarrow \infty$. Here $\underline{\tau}^k = \{\tau_j^k\}_{j=-\infty}^{\infty}$. Let $V_k = \mathcal{S}_{\underline{\tau}^k, n, \phi}$. Then

$$\overline{\bigcup_{k=0}^{\infty} V_k} = L^2(\mathbb{R}). \quad (2.6)$$

Proof. As the Fourier transform is an isometrical isomorphism $L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$, we may prove (2.6) in the Fourier domain. Let $\hat{h} \in C_c(\mathbb{R})$, $C_c(\mathbb{R})$ denoting the space of compactly supported continuous functions on the reals. This is dense in $L^2(\mathbb{R})$. We need to show that \hat{h} can be approximated arbitrarily closely in $L^2(\mathbb{R})$ by a function of the form $\sum_{j=-\infty}^{\infty} c_j \widehat{B_j^{\mathbf{f}, [k]}} \hat{\phi}$ if k is large enough, where $c = \{c_j\}_{j=-\infty}^{\infty} \in \ell^2(\mathbb{Z})$ and where the $\{B_j^{\mathbf{f}, [k]}\}_{j=-\infty}^{\infty}$ are the B-splines in $\mathcal{S}_{\underline{\tau}^k, n, \delta}$. Since $\hat{\phi}(t) > 0$, $\hat{h}/\hat{\phi} \in L^2(\mathbb{R})$ and it can be approximated arbitrarily closely by $\sum_{j=-\infty}^{\infty} c_j \widehat{B_j^{\mathbf{f}, [k]}}$ if k is large enough. This shows that the theorem is true. \blacksquare

3. A prewavelet on non-equally spaced centres.

In this section we shall construct prewavelets on non-equally spaced centres with the aid of certain “fundamental functions” of splines of degree $2n + 1$. Precisely, suppose that there exist $d^\ell = \{d_k^\ell\}_{k=-\infty}^{\infty} \in \ell^1(\mathbb{Z}) \setminus \{0\}$, $\ell \in \mathbb{Z}$, such that

$$\Psi_\ell(x) = \sum_{k=-\infty}^{\infty} d_k^\ell \int_{-\infty}^{\infty} B_k^{\mathbf{f}, 2n+1}(x-y)\phi(y) dy, \quad x \in \mathbb{R}, \quad (3.1)$$

where $B_k^{\mathbf{f}, 2n+1} \in \mathcal{S}_{\underline{\tau}, 2n+1, \delta}$ are the B-splines with support $[\tau_k, \tau_{k+2n+2}]$, gives the identity

$$\int_{-\infty}^{\infty} \phi(y - \tau_{j+n+1}) \Psi_\ell(y) dy = \begin{cases} 1 & \text{if } \ell = j \text{ and } n \text{ odd,} \\ 1 & \text{if } \ell = j + 1 \text{ and } n \text{ even,} \\ 0 & \text{otherwise,} \end{cases} \quad j, \ell \in \mathbb{Z}. \quad (3.2)$$

We call these Ψ_ℓ “fundamental functions” because their construction is related to the use of fundamental functions of interpolation for constructing prewavelets in (Chui and Wang, 1992) and in (Buhmann and Micchelli, 1992). We define $\psi_\ell := \Psi_{2\ell}^{(n+1)}$. This is in V_1 because $(n+1)$ -st derivatives of B-splines of degree $2n + 1$ are expressible as finite linear combinations of B-splines of degree n (see Schumaker, 1981, p. 121, for instance, but recall that the B-splines we use are normalized). In order for the differentiation to result in an absolutely convergent series here we need that $\tau_{i+1} - \tau_i$ is bounded away from zero by $\delta > 0$ uniformly in i , because the formula for the derivative of a B-spline involves division by differences of τ_i s. We have already required that this be true.

We assert that the desired orthogonality conditions $\psi_\ell \perp V_0$ hold, i.e. that

$$\int_{-\infty}^{\infty} \psi_\ell(x) C_k(x) dx = 0, \quad k, \ell \in \mathbb{Z}.$$

Together with (1.4), they imply $W_0 \perp V_0$. This orthogonality is settled by a simple argument using integration by parts: Suppose (3.2) holds for a suitable $d^\ell \in \ell^1(\mathbb{Z})$. Let, for any integer ℓ , s_ℓ be defined by

$$s_\ell(x) := \frac{d^{n+1}}{dx^{n+1}} \sum_{k=-\infty}^{\infty} d_k^\ell B_k^{\mathbf{f}, 2n+1}(x), \quad x \in \mathbb{R}. \quad (3.3)$$

We get, using the fact that C_j and $\Psi_{2\ell}$ and all their derivatives up to degree $n+1$ are absolutely integrable and vanish at infinity, due to (2.1) and the B-splines' compact support,

$$\begin{aligned} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} B_j^c(x-y)\phi(y) dy \int_{-\infty}^{\infty} s_{2\ell}(x-y)\phi(y) dy dx \\ = (-1)^{n+1} \int_{-\infty}^{\infty} \sum_{k=-\infty}^{\infty} \mu_k^j \phi(x-x_k) \Psi_{2\ell}(x) dx = 0, \quad j \in \mathbb{Z}, \end{aligned} \quad (3.4)$$

because the $(n+1)$ -st derivative of B_j^c is a finite collection of delta functions $\underline{\delta}(\cdot - x_k)$, $k \in \mathbb{Z}$, with real coefficients μ_k^j . Before we state our principal theorem, we introduce some notation, namely

$$\Phi(\theta) := \int_{-\infty}^{\infty} \phi(y)\phi(y-\theta) dy, \quad \theta \in \mathbb{R},$$

in other words $\Phi = \phi * \phi(-\cdot)$. This is a continuous function because of (2.1).

Theorem 3. *The functions (3.1) exist as desired and they are such that $\psi_\ell \in V_1$ and $V_1 = V_0 \oplus W_0$, if the three sufficient conditions*

$$\sup_{j \in \mathbb{Z}} \frac{\tau_{j+2n+2} - \tau_j}{2n+2} \sup_{t \in (-\infty, \infty)} \sum_{k=-\infty}^{\infty} \phi(t - \tau_k) < 2 \inf_{k \in \mathbb{Z}} \frac{\tau_{k+2n+2} - \tau_k}{2n+2} \inf_{\theta \in (\tau_k - \tau_{k+n+1}, \tau_{k+2n+2} - \tau_{k+n+1})} \Phi(\theta), \quad (3.5)$$

$$\sup_{k \in \mathbb{Z}} \frac{\tau_{2k+2n+1} - \tau_{2k-1}}{n+1} < 2 \min \left\{ 1, \inf_{k \in \mathbb{Z}} \frac{(\tau_{2k+2n+1} - \tau_{2k-1})^2}{(n+1)^2} \inf_{\theta \in (0, \tau_{2k+2n+1} - \tau_{2k-1})} \Phi(\theta) \right\} \quad (3.6)$$

and

$$\inf_{k \in \mathbb{Z}} \frac{\tau_{k+2n+2} - \tau_k}{2n+2} \inf_{\theta \in (\tau_k - \tau_{k+n+1}, \tau_{k+2n+2} - \tau_{k+n+1})} \Phi(\theta) > \frac{1}{2} \quad (3.7)$$

hold. In other words, the ψ_ℓ are prewavelets, since they generate $W_0 \perp V_0$ with $V_1 = V_0 \oplus W_0$.

These prewavelets satisfy the summability properties

$$\sum_{j=-\infty}^{\infty} |\psi_\ell(x - \tau_j)| \leq \text{const.} < \infty \quad \text{uniformly in } x \in \mathbb{R} \text{ and } \ell \in \mathbb{Z} \quad (3.8)$$

and

$$\sum_{\ell=-\infty}^{\infty} |\psi_\ell(x)| \leq \text{const.} < \infty \quad \text{uniformly in } x \in \mathbb{R}. \quad (3.9)$$

It follows from our proof of the theorem that (3.5), (3.6) and (3.7) may be replaced by three *weaker* but harder to verify conditions. (That is the reason why we have opted to mention them now rather than in the statement of the theorem.) In order to state them, it is convenient to adopt yet another notation, viz.

$$D_{jk} := \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \phi(y - \tau_{k+n+1}) B_j^{\mathbf{f}, 2n+1}(y - z) \phi(z) dz dy, \quad j, k \in \mathbb{Z}. \quad (3.10)$$

This is well-defined because of (2.1). The new conditions are

$$\sup_{j \in \mathbb{Z}} \sum_{k=-\infty}^{\infty} D_{jk} < 2 \inf_{k \in \mathbb{Z}} D_{kk}, \quad (3.11)$$

$$\sup_{k \in \mathbb{Z}} \frac{\tau_{2k+2n+1} - \tau_{2k-1}}{n+1} < 2 \min \left\{ 1, \inf_{k \in \mathbb{Z}} \frac{1}{2\pi} \int_{-\infty}^{\infty} |\hat{\phi}(y)|^2 |\widehat{B}_k^{\mathbf{c}}(y)|^2 dy \right\} \quad (3.12)$$

and

$$\inf_{k \in \mathbb{Z}} D_{kk} > \frac{1}{2}, \quad (3.13)$$

respectively. Note that there are sets $\underline{\tau}$ that satisfy these conditions, because, for instance, any small enough perturbation of $\underline{\tau} = \frac{1}{2}\mathbb{Z}$ does for the multiquadric function $\phi = \frac{1}{2}\varrho''$, $\varrho(r) = \sqrt{r^2 + \gamma^2}$, $n = 1$, if γ is small enough. For that choice, the difference between the right and the left-hand sides of (3.11) is $\approx \frac{1}{3}$ for small γ , and so is the difference between the right and the left-hand sides of (3.12). The left-hand side of (3.13) is $\approx \frac{2}{3}$ when γ is small.

Proof. Condition (3.2) means that the d^ℓ are the ℓ -th row of the inverse of the bi-infinite matrix $D = \{D_{jk}\}_{j,k=-\infty}^{\infty}$ when n is odd, otherwise it is the inverse of D shifted by one. This matrix D is a *continuous* linear operator $D : \ell^p(\mathbb{Z}) \rightarrow \ell^p(\mathbb{Z})$ for $p \in \{1, \infty\}$ (thus for all $p \in [1, \infty]$ by the Riesz convexity theorem) as we shall see now. Indeed, the matrix D has $\ell^1(\mathbb{Z})$ norm one because the B-splines form a partition of unity and because it was a condition on ϕ that its integral be one. Further, the matrix D has bounded $\ell^\infty(\mathbb{Z})$ norm. This is a consequence of Schumaker's (1981) formula on p. 128 for the zero moment of a (normalized) B-spline, namely

$$\|D\|_\infty \leq \sup_{j \in \mathbb{Z}} \frac{\tau_{j+2n+2} - \tau_j}{2n+2} \sup_{t \in (-\infty, \infty)} \sum_{k=-\infty}^{\infty} \phi(t - \tau_k). \quad (3.14)$$

The expression on the right-hand side of (3.14) is finite because $\kappa < \infty$ and (2.1), together with the boundedness condition $\tau_{k+1} - \tau_k \geq \delta$, $k \in \mathbb{Z}$. Thus, under the conditions stated in the theorem, $\|D\|_\infty < \infty$. If (3.11) instead of (3.5) is demanded, the $\ell^\infty(\mathbb{Z})$ boundedness of the operator D becomes, in fact, trivial.

D may be inverted in $\ell^\infty(\mathbb{Z})$ by a Neumann series approach, because $\|D - I\|_\infty < 1$, as we shall now derive from (3.5) (or (3.11)). (Here I is the identity matrix on $\mathbb{Z} \times \mathbb{Z}$.) To explain this point in detail, observe that there is a $\theta \in (\tau_k - \tau_{k+n+1}, \tau_{k+2n+2} - \tau_{k+n+1})$ such that

$$\begin{aligned} D_{kk} &= \int_{-\infty}^{\infty} \Phi(z) B_k^{\mathbf{f}, 2n+1}(\tau_{k+n+1} + z) dz \\ &= \int_{\tau_k - \tau_{k+n+1}}^{\tau_{k+2n+2} - \tau_{k+n+1}} \Phi(z) B_k^{\mathbf{f}, 2n+1}(\tau_{k+n+1} + z) dy \\ &= \Phi(\theta) \int_{\tau_k}^{\tau_{k+2n+2}} B_k^{\mathbf{f}, 2n+1}(z) dz \\ &= \frac{\tau_{k+2n+2} - \tau_k}{2n+2} \Phi(\theta). \end{aligned}$$

Nonnegativity of D 's entries and $\|D\|_1 = 1$ imply $D_{kk} \leq 1$ and so

$$\|D - I\|_\infty \leq \|D\|_\infty + 1 - 2 \inf_{k \in \mathbb{Z}} D_{kk}. \quad (3.15)$$

The penultimate display, (3.14), (3.15) and (3.5) give the desired estimate $\|D - I\|_\infty < 1$. Therefore D is a homeomorphism on $\ell^\infty(\mathbb{Z})$, whence the coefficients d_k^ℓ exist with $\sup_{\ell \in \mathbb{Z}} \|d^\ell\|_1 < \infty$. That the weaker requirement (3.11) may replace (3.5), with D 's invertibility by Neumann series remaining intact, can be seen from (3.15).

It is a direct consequence of $d^\ell \in \ell^1(\mathbb{Z})$ and our conditions in the statement of the theorem that

$$\sum_{j=-\infty}^{\infty} |\Psi_\ell(x - \tau_j)| \leq \sup_{t \in (-\infty, \infty)} \sum_{j=-\infty}^{\infty} \phi(t - \tau_j) \sup_{\ell \in \mathbb{Z}} \|d^\ell\|_1 \sup_{k \in \mathbb{Z}} \frac{\tau_{k+2n+2} - \tau_k}{2n+2}. \quad (3.16)$$

The right-hand side is finite because the right-hand side of (3.14) is. In order to deduce (3.8) from (3.16), one uses the fact that $(n+1)$ -st derivatives of B-splines of degree $2n+1$ are finite linear combinations of those of degree n (Schumaker, 1981, p. 121). Precisely, the left-hand side of (3.8) is bounded above by

$$\frac{1}{2} \sup_{t \in (-\infty, \infty)} \sum_{j=-\infty}^{\infty} \phi(t - \tau_j) \sup_{\ell \in \mathbb{Z}} \|d^\ell\|_1 \sup_{k \in \mathbb{Z}} \frac{\tau_{k+n+1} - \tau_k}{n+1} \left(\frac{1}{2}\delta\right)^{-n-1} \quad (3.17)$$

according to Schumaker's differentiation formula. If (2.1) holds also for derivatives of ϕ of order at most $n+1$, then the bound (3.17) may be replaced by

$$\sup_{t \in (-\infty, \infty)} \sum_{j=-\infty}^{\infty} \phi^{(n+1)}(t - \tau_j) \sup_{\ell \in \mathbb{Z}} \|d^\ell\|_1 \sup_{k \in \mathbb{Z}} \frac{\tau_{k+2n+2} - \tau_k}{2n+2},$$

which is a bound that does not contain the increasing factor δ^{-n-1} as the spacing of the τ_k 's decreases.

We have to show (3.9). By the same arguments as in the penultimate paragraph,

$$\|D - I\|_1 \leq \|D\|_1 + 1 - 2 \inf_{k \in \mathbb{Z}} D_{kk}.$$

Hence, under our assumption (3.7) or, in its weaker form, (3.13), $\|D - I\|_1 < 1$. So the coefficients d_k^ℓ are also absolutely summable over $\ell \in \mathbb{Z}$. Using the fact that the B-splines $\{B_k^{\mathbf{f}, 2n+1}\}_{k=-\infty}^\infty$ form a partition of unity, we may conclude from $d_k = \{d_k^\ell\}_{\ell=-\infty}^\infty \in \ell^1(\mathbb{Z})$, $k \in \mathbb{Z}$, and (2.1) that

$$\sum_{\ell=-\infty}^\infty |\Psi_\ell(x)| \leq \sup_{k \in \mathbb{Z}} \|d_k\|_1. \quad (3.18)$$

This inequality implies (3.9) because Schumaker's differentiation formula for B-splines supplies once more the explicit bound

$$\sum_{\ell=-\infty}^\infty |\psi_\ell(x)| \leq \frac{1}{2} \sup_{k \in \mathbb{Z}} \|d_k\|_1 \left(\frac{1}{2}\delta\right)^{-n-1}. \quad (3.19)$$

A remark as the one after (3.17) applies here as well: in this case, (3.19) may be replaced by

$$\sum_{\ell=-\infty}^\infty |\psi_\ell(x)| \leq \sup_{k \in \mathbb{Z}} \|d_k\|_1 \int_{-\infty}^\infty \phi^{(n+1)}.$$

The claim $V_1 = V_0 \oplus W_0$ is established as follows. Recall that every $g \in V_1$ is expressible as an infinite linear combination of F_j s with square-summable coefficients. We require a decomposition of each $F_j \in V_1$ first. To this end, we seek coefficients B_{jk} , $j, k \in \mathbb{Z}$, such that the orthogonality conditions

$$\eta_j := F_j - \sum_{k=-\infty}^\infty B_{jk} C_k \perp C_\ell, \quad \ell \in \mathbb{Z}, \quad (3.20)$$

hold for all integers j . In other words, we require that

$$\sum_{k=-\infty}^\infty B_{jk} \langle C_k, C_\ell \rangle = \langle F_j, C_\ell \rangle, \quad \ell \in \mathbb{Z}, \quad (3.21)$$

where $\langle \cdot, \cdot \rangle$ is the standard $L^2(\mathbb{R})$ inner product. Once such B_{jk} are found, we shall expand η_j in terms of the ψ_k s to reach the desired conclusion $F_j \in V_0 \oplus W_0$.

As the entries of the symmetric matrix $\chi = \{\langle C_k, C_\ell \rangle\}_{k, \ell \in \mathbb{Z}}$ are nonnegative and the B-splines form a partition of unity, the matrix has $\ell^1(\mathbb{Z})$ and $\ell^\infty(\mathbb{Z})$ norm $\sup_{k \in \mathbb{Z}} (\tau_{2k+2n+1} - \tau_{2k-1}) / (n+1)$. This is finite because κ is. More importantly it is true that, under condition (3.12) or (3.6), $\|I - \chi\|_1 = \|I - \chi\|_\infty < 1$. The specific argument is as follows: Firstly, we note that

$$\|I - \chi\|_p \leq \sup_{k \in \mathbb{Z}} \frac{\tau_{2k+2n+1} - \tau_{2k-1}}{n+1} + \max\{-1, 1 - 2 \inf_{k \in \mathbb{Z}} \langle C_k, C_k \rangle\}, \quad p = 1, \infty, \quad (3.22)$$

and then

$$\begin{aligned}
\langle C_k, C_k \rangle &= \int_{-\infty}^{\infty} B_k^c(z) \int_{-\infty}^{\infty} \Phi(x) B_k^c(x+z) dx dz \\
&= \int_{\tau_{2k-1}}^{\tau_{2k+2n+1}} B_k^c(z) \int_{-\infty}^{\infty} \Phi(x) B_k^c(x+z) dx dz \\
&= \frac{(\tau_{2k+2n+1} - \tau_{2k-1})^2}{(n+1)^2} \Phi(\theta)
\end{aligned}$$

for a suitable $\theta \in (0, \tau_{2k+2n+1} - \tau_{2k-1})$. Thus, according to the conditions in the statement of our theorem, the right-hand side of (3.22) is less than one. Hence χ is a homeomorphism with inverse χ^{-1} in $\ell^1(\mathbb{Z})$ and in $\ell^\infty(\mathbb{Z})$. That the weaker requirement (3.12) is appropriate follows from the last display and from

$$\langle C_k, C_k \rangle = \frac{1}{2\pi} \int_{-\infty}^{\infty} |\hat{\phi}(y)|^2 |\widehat{B}_k^c(y)|^2 dy$$

by the Parseval–Plancherel identity. (The latter may be applied by (2.5) [for $p = 2$] and the integrability of ϕ .)

The desired coefficients are $B_{jk} := \sum_{\ell \in \mathbb{Z}} \langle F_j, C_\ell \rangle \chi_{\ell k}^{-1}$; we denote the matrix $\{B_{jk}\}_{j,k=-\infty}^{\infty}$ by B . The matrix B is well-defined and $\|B\|_1, \|B\|_\infty$ are finite because of (2.3) and χ 's invertibility. Hence, in particular, $\eta_j \in V_1$ by Proposition 1 and Theorem 275 on p. 198 of (Hardy, Littlewood and Pólya, 1934).

To advance the proof, we require

Lemma 4. *Let $\eta_j, j \in \mathbb{Z}$, be defined by (3.20). There exists, for each j , H_j such that $H_j^{(n+1)} = \eta_j$.*

Moreover,

$$\sup_{j \in \mathbb{Z}} \sum_{k=-\infty}^{\infty} |\langle \phi(\cdot - \tau_{2k}), H_j \rangle| < \infty \quad (3.23)$$

and

$$\sup_{k \in \mathbb{Z}} \sum_{j=-\infty}^{\infty} |\langle \phi(\cdot - \tau_{2k}), H_j \rangle| < \infty. \quad (3.24)$$

Proof. We note first that there is such an H_j for each integer j . We can provide it explicitly:

$$H_j(x) = \frac{1}{n!} \int_{-\infty}^{\infty} (x-t)_+^n \eta_j(t) dt, \quad x \in \mathbb{R}, \quad (3.25)$$

where $(\cdot)_+^n$ is a truncated power. It follows from η_j 's definition, $\|B\|_\infty < \infty$ and the upper bound (2.1) that H_j is well-defined. We observe

$$\langle H_j, \phi(\cdot - \tau_{2\ell-1}) \rangle = 0, \quad \ell \in \mathbb{Z}. \quad (3.26)$$

In order to prove (3.26), we note that, by integration by parts and (3.20),

$$\begin{aligned} 0 &= \langle \eta_j, C_\ell \rangle \\ &= (-1)^{n+1} \left\langle H_j, \int_{-\infty}^{\infty} B_\ell^{c(n+1)}(y) \phi(\cdot - y) dy \right\rangle \\ &= \left\langle H_j, \sum_{k=-\infty}^{\infty} \mu_k^\ell \phi(\cdot - \tau_{2k-1}) \right\rangle, \end{aligned}$$

where the $\{\mu_k^\ell\}_{k=-\infty}^{\infty}$ are fixed multiples of the coefficients of the $(n+1)$ -st order divided difference operator based on the knots associated with B_ℓ^c 's support. The last display implies (3.26) as $\langle H_j, \phi(\cdot - \tau_k) \rangle$ is bounded but not constant, whereas there are no nontrivial bounded sequences (except constants) in the kernel of the aforementioned difference operator.

Therefore, by taking an $(n+1)$ -st divided difference of $\langle \phi(\cdot - y), H_j \rangle$ with respect to y at the $n+2$ points $\tau_{2k-1}, \tau_{2k}, \tau_{2k+1}, \tau_{2k+3}, \tau_{2k+5}, \dots, \tau_{2k+2n-1}$, and by the Peano kernel theorem,

$$|\langle \phi(\cdot - \tau_{2k}), H_j \rangle| \leq C \left| \left\langle \phi(\cdot - \tilde{\tau}_{2k}) [\tau_{2k+2n-1} - \tau_{2k-1}]^{n+1}, |\eta_j| \right\rangle \right| \quad (3.27)$$

for a suitable positive constant C . Here, $\tilde{\tau}_{2k} \in (\tau_{2k-1}, \tau_{2k+2n-1})$. If we sum (3.27) over $j \in \mathbb{Z}$, the result is bounded independently of k , because $\sum_{j=-\infty}^{\infty} |\eta_j|$ is uniformly bounded due to (2.1), the compact support of the B-splines and $\|B\|_1 < \infty$. By the same token, if (3.27) is summed over k , the result is again bounded, independently of j , since $\eta_j \in L^1(\mathbb{R})$ and $\kappa < \infty$. \blacksquare

We continue with the proof of Theorem 3 and recall that it is still our aim to expand each η_j in terms of the ψ_k which proves $F_j \in V_0 \oplus W_0$. Equipped with Lemma 4, our goal is precisely to show the following lemma:

Lemma 5. *Let $\eta_j, j \in \mathbb{Z}$, be defined by (3.20). Then*

$$\eta_j = \sum_{k=-\infty}^{\infty} A_{jk} \psi_k, \quad j \in \mathbb{Z}, \quad (3.28)$$

for coefficients A_{jk} which are absolutely summable over $j \in \mathbb{Z}$ as well as over $k \in \mathbb{Z}$.

Proof. We observe immediately that it suffices to prove instead of (3.28) that

$$\tilde{H}_j = \sum_{k=-\infty}^{\infty} A_{jk} \Psi_{2k}, \quad j \in \mathbb{Z}, \quad (3.29)$$

for suitable coefficients A_{jk} and for an \tilde{H}_j with $\tilde{H}_j^{(n+1)} = \eta_j$, because we may differentiate on both sides of this identity $(n+1)$ -times. We claim that, due to the form of the Ψ_k and due to properties

of H_j , the following choice is suitable:

$$\left. \begin{aligned} \tilde{H}_j &:= \sum_{k=-\infty}^{\infty} \langle \phi(\cdot - \tau_{2k+n+1}), H_j \rangle \Psi_{2k}, & n \text{ odd}, \\ \tilde{H}_j &:= \sum_{k=-\infty}^{\infty} \langle \phi(\cdot - \tau_{2k+n}), H_j \rangle \Psi_{2k}, & n \text{ even}, \end{aligned} \right\} j \in \mathbb{Z}, \quad (3.30)$$

the series being absolutely convergent because the Ψ_{2k} are uniformly bounded by $\sup_{\ell \in \mathbb{Z}} \|d^\ell\|_1$ and because of Lemma 4. Theorem 275 on p. 198 of (Hardy, Littlewood and Pólya, 1934), Lemma 4 and $\|d^\ell\|_1 < \infty$ imply that the right-hand sides of (3.30) are in V_1^{2n+1} , where we let $V_1^{2n+1} = \mathcal{S}_{\tau, 2n+1, \phi}$.

By taking inner products $\langle \cdot, \cdot \rangle$ with $\phi(\cdot - \tau_\ell)$ on both sides of (3.30), we observe that (3.2) implies for odd n

$$\langle \phi(\cdot - \tau_{2\ell+n+1}), \tilde{H}_j \rangle = \langle \phi(\cdot - \tau_{2\ell+n+1}), H_j \rangle, \quad \ell \in \mathbb{Z},$$

while (3.26) implies

$$\langle \phi(\cdot - \tau_{2\ell+n}), \tilde{H}_j \rangle = \langle \phi(\cdot - \tau_{2\ell+n}), H_j \rangle = 0, \quad \ell \in \mathbb{Z}.$$

There are analogous identities for even n .

Nonsingularity of D as an operator $\ell^2(\mathbb{Z}) \rightarrow \ell^2(\mathbb{Z})$ means that the implication

$$f \in V_1, F \in V_1^{2n+1}, F^{(n+1)} = f, \langle \phi(\cdot - \tau_\ell), F \rangle = 0 \quad \forall \ell \in \mathbb{Z} \quad \implies \quad f = 0$$

holds. Hence $\tilde{H}_j^{(n+1)} = H_j^{(n+1)} = \eta_j$, as required.

We have thus found the coefficients A_{jk} , $j, k \in \mathbb{Z}$. They are not only absolutely summable over k , but also over j because of Lemma 4. Therefore Lemma 5 is settled. \blacksquare

Now let $g \in V_1$, $g = \sum_{j=-\infty}^{\infty} c_j F_j$, for $c = \{c_j\}_{j=-\infty}^{\infty} \in \ell^2(\mathbb{Z})$. Then, according to Lemma 5,

$$g = \sum_{k=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} c_j A_{jk} \psi_k + \sum_{k=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} c_j B_{jk} C_k$$

with

$$\sum_{k=-\infty}^{\infty} \left(\sum_{j=-\infty}^{\infty} c_j A_{jk} \right)^2 < \infty, \quad \sum_{k=-\infty}^{\infty} \left(\sum_{j=-\infty}^{\infty} c_j B_{jk} \right)^2 < \infty. \quad (3.31)$$

The second inequality in (3.31) follows from the fact that $\sum_{k=-\infty}^{\infty} |B_{jk}| \leq \text{const.} < \infty$ and $\sum_{j=-\infty}^{\infty} |B_{jk}| \leq \text{const.} < \infty$, uniformly in j and k , respectively, and from Theorem 275 on p. 198 of (Hardy, Littlewood and Pólya, 1934). The first inequality in (3.31) is true by Lemma 5. \blacksquare

We remark that the assertions of Theorem 3 remain true if ϕ is no longer positive so long as it is explicitly demanded that $\|D - I\|_\infty < 1$, $\|D - I\|_1 < 1$ and $\|\chi - I\|_\infty < 1$, in order that the Neumann series argument can be applied.

Theorem 6. *Given the set-up of Theorems 2 and 3, there are prewavelets $\{\psi_{\ell k}\}_{\ell=-\infty}^{\infty}$, $k \in \mathbb{Z}_+$, in V_{k+1} and spaces W_k spanned by sums of those prewavelets with square-summable coefficients such that $V_{k+1} = V_k \oplus W_k$, $k \in \mathbb{Z}_+$, and*

$$L^2(\mathbb{R}) = V_0 \oplus \bigoplus_{k=0}^{\infty} W_k, \quad (3.32)$$

if the assumptions of Theorem 3 hold for all $\underline{\tau}^k$.

Proof. Let $f \in L^2(\mathbb{R})$ and let a positive ε be given. Due to Theorem 2, there are $k \in \mathbb{Z}_+$ and $f_k \in V_k$ such that

$$\|f - f_k\|_2^2 \leq \varepsilon.$$

Further,

$$f_{\ell+1} = f_{\ell} + g_{\ell}, \quad f_{\ell} \in V_{\ell}, \quad g_{\ell} \in W_{\ell}, \quad \forall 0 \leq \ell < k.$$

Therefore, the arbitrariness of $\varepsilon > 0$ and the orthogonality between different W_{ℓ} s imply (3.32). ■

We point out that, according to the requirements of Theorem 3, it may not be possible to construct prewavelets that span W_k for arbitrarily large k depending on $\underline{\tau}^k$, the prescribed radial function and its parameters. Therefore, in practice and when using multiquadrics for instance, γ must be chosen suitably to allow that the sufficient conditions for the existence of prewavelets with small stepsizes in Theorem 3 are met, where the stepsize should be according to the resolution we wish to obtain.

Further examples to our theory can be obtained by applying it, for example, to $(r^2 + \gamma^2)^{\frac{1}{2}\beta}$ (where $\beta \in \mathbb{Z} \setminus 2\mathbb{Z}_+$). Quickly decaying radial functions with positive Fourier transforms $\phi(r) = \exp(-\gamma^2 r^2)$, $\phi(r) = \exp(-\gamma r)$, etc. can be used too, in this case directly as ϕ without any differentiating. The essential feature that admits derivation of suitable decaying functions from the radial functions by differentiation is that $\hat{\varrho}$ be positive and integrable outside a neighbourhood of the origin, with perhaps an integer order singularity at zero. It is the purpose of the differentiation to resolve this singularity. More specifically, $\varrho(r) = \sqrt{r^2 + \gamma^2}^3$ with $\phi = \varrho^{(4)}$ or $\phi(r) = \frac{1}{r^2 + \gamma^2}$ are both suitable choices for $n = 3$ and $n = 0$, respectively, except they have to be normalized to integrate to one.

It is highly relevant to the computation of the prewavelets that, if $\phi = \varrho^{(n+1)}$ as in our initial multiquadric example for $n = 1$, the entries of the matrix D are a constant multiple of

$$\int_{-\infty}^{\infty} \varrho(y - \tau_j) \phi_{*}^k(y) dy, \quad (3.33)$$

where ϕ_*^k is a $(2n + 2)$ -nd order divided difference of the function g based on the centres that are associated with $B_k^{\mathbf{f}, 2n+1}$'s support. The formula (3.33) is derived from (3.10) by integration by parts. By the same token, the prewavelets themselves can be expressed as constant multiples of

$$\sum_{k=-\infty}^{\infty} d_k^{2\ell} \phi_*^k(y), \quad y \in \mathbb{R},$$

instead of (3.1). This observation has been used to create the image of the multiquadric wavelet for $n = 1$ that is attached to this paper. It uses integer data randomly perturbed (by at most 0.4) and $\gamma = 0.1$. The positions of the centres are indicated by crosses. So we see there a multiquadric prewavelet for scattered centres, probably for the first time in the literature. It remains to add, however, that the theory presented here is far from complete, which reflects the fact that the approach to scattered data prewavelets is substantially different from that of prewavelets on gridded data. For instance, no Riesz basis properties are shown of the prewavelets presented here nor are there concrete estimates for their asymptotic behaviour at infinity.

In order not to end on a downbeat note, we point out that it was shown in (Buhmann, 1993b) how to derive explicit decay estimates for prewavelets and decomposition coefficients A_{jk}, B_{jk} , if ϕ is a derivative of a radial basis function from the class discussed in that paper and the centres are equally spaced. In this context it is also important to note that, when the centres are equally spaced, Riesz stability for the $\ell^2(\mathbb{Z})$ norm can be shown, so that the prewavelets here form a Riesz basis for W_0 . We do not elaborate on this point, however, because the subject of this paper are prewavelets on non-equally spaced centres. (This claim is actually easily established using (2.5) and the fact that D is a homeomorphism on $\ell^2(\mathbb{Z})$ —the latter follows from D being a homeomorphism on $\ell^p(\mathbb{Z})$, $p \in \{1, \infty\}$, and from the Riesz convexity theorem.) The subject of Riesz stability will be one of the main topics in the forthcoming paper (Buhmann, 1994).

References.

1. de Boor, C. (1976) “Odd-degree spline interpolation at a biinfinite knot sequence”, in: *Approximation Theory Bonn 1976*, R. Schaback and K. Scherer, eds., Springer Lecture Notes in Mathematics 556, Springer (Berlin), 30–53.
2. de Boor, C., R.A. DeVore and A. Ron (1993) “On the construction of multivariate (pre)wavelets”, *Constr. Approx.* **9**, 123–166.
3. Buhmann, M.D. (1993a) “New developments in the theory of radial basis function interpolation”, in *Multivariate Approximation: From CAGD to Wavelets*, K. Jetter and F.I. Utreras, eds., World Scientific (Singapore), 35–75.
4. Buhmann, M.D. (1993b) “Discrete least squares approximation and pre-wavelets from radial function spaces”, *Mathemat. Proc. Cambridge Philosophical Society* **114** (1993), 533–558.
5. Buhmann, M.D. (1993c) “On quasi-interpolation with radial basis functions”, *J. Approx. Theory* **72** (1993), 103–130.
6. Buhmann, M.D. (1994) “Pre-Wavelets on Scattered Knots and from Radial Function Spaces: A Review”, in preparation for *Proceedings of the 6th Conference on the Mathematics of Surfaces*, R.R. Martin, ed., IMA Conference Proceedings Series, Oxford University Press.
7. Buhmann, M.D. and C.A. Micchelli (1992) “Spline prewavelets with non-uniform knots”, *Numerische Mathematik* **61**, 455–474.
8. Buhmann, M.D. and N. Dyn (1991) “Error estimates for multiquadric interpolation”, in *Curves and Surfaces*, P.J. Laurent, A. LeMéhauté and L.L. Schumaker, eds., Academic Press (Boston), 51–58.
9. Chui, C.K. and J.Z. Wang (1992) “A cardinal spline approach to wavelets”, *Proc. AMS.* **113**, 785–793.
10. Chui, C.K., J.D. Ward and J. Stöckler (1994) “Analytic wavelets generated by radial functions”, ms.
11. Hardy, G.H., J.E. Littlewood, J.E. and G. Pólya (1934) *Inequalities*, Cambridge University Press (Cambridge).
12. Micchelli, C.A., C. Rabut and F. Utreras (1991) “Using the refinement equation for the construction of pre-wavelets III: elliptic splines”, *Num. Alg.* **1**, 331–352.
13. Powell, M.J.D. (1990) “Univariate multiquadric interpolation: reproduction of linear polynomials”, in: *Multivariate Approximation and Interpolation*, W. Haussmann and K. Jetter, eds., Birkhäuser Verlag (Basel), 227–240.

14. Powell, M.J.D. (1992) “Tabulation of thin plate splines on a very fine two-dimensional grid”, in: *Numerical Methods of Approximation Theory*, D. Braess and L.L Schumaker, eds., Birkhäuser (Basel), 221–244.
15. Schumaker, L.L. (1981) *Spline Functions: Basic Theory*, Wiley (New York).