



Conference Poster

Substructures in cold dark matter halos

Author(s):

De Lucia, Gabriella; Kauffmann, G.; White, S.D.M.; Springel, V.; Lanzoni, B.; Stoehr, Felix; Tormen, G.; Yoshida, N.

Publication Date:

2003

Permanent Link:

<https://doi.org/10.3929/ethz-a-004582538> →

Rights / License:

[In Copyright - Non-Commercial Use Permitted](#) →

This page was generated automatically upon download from the [ETH Zurich Research Collection](#). For more information please consult the [Terms of use](#).



G. De Lucia¹, G. Kauffmann¹, S. D. M. White¹, V. Springel¹, B. Lanzoni², F. Stoehr¹, G. Tormen³, N. Yoshida⁴

¹ Max-Planck-Institut für Astrophysik, München, Germany, ² Osservatorio Astronomico di Bologna, Italy
³ Università di Padova, Italy, ⁴ Harvard-Smithsonian Center for Astrophysics, USA

Abstract: We analyse the properties of substructures within dark matter halos (subhalos) using a set of high-resolution numerical simulations (see poster "Chemical enrichment of the ICM in a hierarchical merger model", by De Lucia et al.) of the formation of structure in a Λ CDM universe.

The subhalo mass function:

The substructure mass function depends at most weakly on the mass of the parent halo, as already suggested by Moore et al. (1999).




Figure 1: Cumulative subhalo mass function for halos with different mass. Black line is for halos with mass $\sim 10^{14} M_{\odot}$, red line is for halos with mass $\sim 10^{13} M_{\odot}$, green line is for halos with mass $\sim 10^{12} M_{\odot}$ and blue line is for halos with mass $\sim 10^{11} M_{\odot}$. Dashed lines show the corresponding dark matter profiles.

The spatial distribution of substructures:

The subhalo profiles are "anti-biased" relative to the dark matter, in agreement with previous results by Ghigna et al. (2000). We also find that the radial number density profiles are steeper in low mass halos than in high mass halos. This suggests that subhalos have been more efficiently destroyed in the dense inner cores of massive clusters.





Figure 2: Radial distribution of substructures in halos of different mass. Black line is for halos with mass $\sim 10^{14} M_{\odot}$, red line is for halos with mass $\sim 10^{13} M_{\odot}$, green line is for halos with mass $\sim 10^{12} M_{\odot}$ and blue line is for halos with mass $\sim 10^{11} M_{\odot}$. Dashed lines show the corresponding dark matter profiles.

Subhalo histories:

Measuring the merger tree for each substructure we find that a significant fraction of the substructures residing in clusters at the present day were accreted at redshift $z \sim 1$.



This implies that a significant fraction of present-day 'passive' cluster galaxies should have been still outside the cluster progenitor and more active at $z \sim 1$.

Figure 3: Distribution of accretion redshifts for subhalos residing in clusters at present-day.

The mass accretion function:

Low mass subhalos are in place earlier than the more massive ones, as expected in a hierarchical cosmology. Tidal stripping is effective on short time-scales. The longer the substructure spends in a more massive halo, the larger is the destructive effect of tidal stripping. Field and cluster subhalos have remarkably similar histories suggesting that the efficiency of the tidal stripping is largely independent of the mass of the parent halo. Our results hint that substructures are constantly erased in the cluster and are being replenished by newly infalling halos.




Figure 4: Average mass accretion history for subhalos residing in clusters at redshift zero (solid lines) and subhalos in field (dashed line). The histories are normalised to the mass of the subhalo at the accretion time. The black and green line are for subhalos with mass $\sim 10^{11} M_{\odot}$; the red line is for subhalos with mass $\sim 10^{12} M_{\odot}$.

Merging histories:




Figure 4: The fraction of substructures with at least one major merger event at redshift $z_m < z$. The solid line is for substructures accreted at redshift < 0.5 , the dashed line is for substructures accreted at redshift $0.5 < z < 1.0$ and the dashed-dotted line is for substructures accreted at redshift > 1.0

About 80 % of the subhalos with mass $\sim 10^{12} M_{\odot}$ and residing in the cluster at present-day have had at least one major merger at redshift below 2. The fraction is almost independent of the accretion time. These results suggest that a large fraction of subhalos in our cluster sample will host early-type galaxies with a significant bulge component.

References:

- Ghigna S., Moore B., Governato F., Lake G., Quinn T., Stadel J., 2000, *Apl*, 544, 616
- Lanzoni B., Cappi A., Ciotti L., 2002, in "Computational astrophysics in Italy: methods and tools", SAIt Proc., preprint, astro-ph/0212131
- Moore B., Ghigna S., Governato F., Lake G., Quinn T., Tozzi P., 1999, *Apl*, 524, L19
- Tormen G., Bouchet F. R., White S. D. M., 1997, *MNRAS*, 286, 865