



Journal Article

Complete genome sequence of *Anoxybacillus flavithermus* strain 52-1A isolated from a heat-processed powdered milk concentrate

Author(s):

Tasara, Taurai; Morach, Marina; Klumpp, Jochen; Stephan, Roger

Publication Date:

2017-08

Permanent Link:

<https://doi.org/10.3929/ethz-b-000192454> →


Originally published in:

Genome Announcements 5(32), <http://doi.org/10.1128/genomeA.00800-17> →

Rights / License:

[Creative Commons Attribution 4.0 International](#) →

This page was generated automatically upon download from the [ETH Zurich Research Collection](#). For more information please consult the [Terms of use](#).



Complete Genome Sequence of *Anoxybacillus flavithermus* Strain 52-1A Isolated from a Heat-Processed Powdered Milk Concentrate

Taurai Tasara,^a Marina Morach,^a  Jochen Klumpp,^b Roger Stephan^a

Institute for Food Safety and Hygiene, Vetsuisse Faculty University of Zurich, Zurich, Switzerland^a; Institute of Food, Nutrition and Health, ETH Zurich, Zurich, Switzerland^b

ABSTRACT The thermophilic spore-forming bacterium *Anoxybacillus flavithermus* is responsible for powdered milk product spoilage, and its presence in dairy processing environments is a concern. Here, the complete genome sequence of the *A. flavithermus* strain 52-1A isolated from a heat-processed powdered milk product concentrate in Switzerland is presented.

The contamination of milk powder by spore-forming heat-resistant milk spoilage bacteria remains an important concern for the dairy industry (1). *Anoxybacillus flavithermus* is a thermophilic sporeformer that frequently occurs as a contaminant in dairy processing environments, leading to fouling of powdered milk products (1–3). Only a few *A. flavithermus* strains have had their complete genomes sequenced to date (4–6). An analysis based on this limited number of *A. flavithermus* genomes has shown that this bacterium is harboring genes that are compatible with its lifestyle adaptations, such as high growth temperatures and extreme pH conditions (7). More complete polished *A. flavithermus* genomes are clearly needed in order to enhance the understanding of the genetic and metabolic diversity as well as physiological mechanisms involved in adaptation of this bacterium to dairy production environments. We have determined the complete genome sequence of the *A. flavithermus* 52-1A strain, which was isolated from a heat-processed powdered milk product concentrate with an off flavor in Switzerland.

Genomic DNA was isolated from *A. flavithermus* 52-1A with the GenElute bacterial genomic DNA kit (Sigma, Buchs, Switzerland) and sequenced using the Pacific Biosciences single-molecule real-time (SMRT) sequencing technology at the Functional Genomics Centre of the University of Zurich. Sequencing generated 47,773 sequence reads (129-fold genome coverage) with an average length of 7,219 kb. The sequence reads were assembled *de novo* using SMRT Analysis 2.3.0 software and the HGAP3 algorithm. This sequence assembly generated a complete *A. flavithermus* genome sequence of 2.8 Mb comprising one chromosome (2,805,288 bp) and one plasmid (24,836 bp). Annotation of the genome was carried out using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (http://www.ncbi.nlm.nih.gov/genome/annotation_prok/) (8). Overall, there were 3,013 genes identified in the *A. flavithermus* 52-1A genome. Of these, 2,985 genes are on the chromosome, whereas 28 genes are plasmid encoded. The *A. flavithermus* 52-1A chromosome has a 42.2% GC content and carries 2,869 coding sequences, 219 pseudogenes, 85 tRNA genes, and 9 rRNA operons. Using the phage search tool PHAST, the *A. flavithermus* 52-1A genome was predicted to have a single prophage region located at positions 1556241 to 1583498 (27,275 bp) (9). Genome comparison with other available *A. flavithermus* genomes shows that the

Received 29 June 2017 Accepted 3 July 2017 Published 10 August 2017

Citation Tasara T, Morach M, Klumpp J, Stephan R. 2017. Complete genome sequence of *Anoxybacillus flavithermus* strain 52-1A isolated from a heat-processed powdered milk concentrate. *Genome Announc* 5:e00800-17. <https://doi.org/10.1128/genomeA.00800-17>.

Copyright © 2017 Tasara et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Roger Stephan, stephanr@fsafety.uzh.ch.

52-1A strain is most closely related to the *A. flavithermus* TNO-09.006 strain, sharing 99.28% average nucleotide identity.

Accession number(s). Genome sequences for the *A. flavithermus* 52-1A chromosome and plasmid are available in GenBank under accession numbers [CP021838](#) and [CP021839](#), respectively.

ACKNOWLEDGMENT

This work was supported by funding from the University of Zurich.

REFERENCES

- Burgess SA, Lindsay D, Flint SH. 2010. Thermophilic bacilli and their importance in dairy processing. *Int J Food Microbiol* 144:215–225. <https://doi.org/10.1016/j.ijfoodmicro.2010.09.027>.
- Heinen W, Lauwers AM, Mulders JW. 1982. *Bacillus flavothermus*, a newly isolated facultative thermophile. *Antonie Leeuwenhoek* 48:265–272. <https://doi.org/10.1007/BF00400386>.
- Zhao Y, Caspers MP, Metselaar KI, de Boer P, Roeselers G, Moezelaar R, Nierop Groot M, Montijn RC, Abee T, Kort R. 2013. Abiotic and microbiotic factors controlling biofilm formation by thermophilic sporeformers. *Appl Environ Microbiol* 79:5652–5660. <https://doi.org/10.1128/AEM.00949-13>.
- Caspers MP, Boekhorst J, Abee T, Siezen RJ, Kort R. 2013. Complete genome sequence of *Anoxybacillus flavithermus* TNO-09.006, a thermophilic spore-former associated with a dairy-processing environment. *Genome Announc* 1(1):e00010-13. <https://doi.org/10.1128/genomeA.00010-13>.
- Matsutani M, Shirakihara Y, Imada K, Yakushi T, Matsushita K. 2013. Draft genome sequence of a thermophilic member of the *Bacillaceae*, *Anoxybacillus flavithermus* strain Kn10, isolated from the Kan-nawa hot spring in Japan. *Genome Announc* 1(3):e00311-13. <https://doi.org/10.1128/genomeA.00311-13>.
- Saw JH, Mountain BW, Feng L, Omelchenko MV, Hou SB, Saito JA, Stott MB, Li D, Zhao G, Wu JL, Galperin MY, Koonin EV, Makarova KS, Wolf YI, Rigden DJ, Dunfield PF, Wang L, Alam M. 2008. Encapsulated in silica: genome, proteome and physiology of the thermophilic bacterium *Anoxybacillus flavithermus* WK1. *Genome Biol* 9:R161. <https://doi.org/10.1186/gb-2008-9-11-r161>.
- Goh KM, Gan HM, Chan KG, Chan GF, Shahar S, Chong CS, Kahar UM, Chai KP. 2014. Analysis of anoxybacillus genomes from the aspects of lifestyle adaptations, prophage diversity, and carbohydrate metabolism. *PLoS One* 9:e90549. <https://doi.org/10.1371/journal.pone.0090549>.
- Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity G, Kodira CD, Kyrpides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of standard operating procedures (SOPs) for (meta)genomic annotation. *Omic* 12:137–141. <https://doi.org/10.1089/omi.2008.0017>.
- Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool. *Nucleic Acids Res* 39:W347–W352. <https://doi.org/10.1093/nar/gkr485>.