



## Journal Article

# Phase transition dynamics of single optically trapped aqueous potassium carbonate particles

**Author(s):**

Esat, Kivanç; David, Grégory; Poulkas, Theodoros; Shein, Mikhail; Signorell, Ruth

**Publication Date:**

2018-05-07

**Permanent Link:**

<https://doi.org/10.3929/ethz-b-000268286> →

**Originally published in:**

Physical Chemistry Chemical Physics 20(17), <http://doi.org/10.1039/c8cp00599k> →

**Rights / License:**

[In Copyright - Non-Commercial Use Permitted](#) →

This page was generated automatically upon download from the [ETH Zurich Research Collection](#). For more information please consult the [Terms of use](#).

# PCCP- Electronic Supplementary Information

## Phase transition dynamics of single optically trapped aqueous potassium carbonate particles<sup>†</sup>

Kivanç Esat, Grégory David, Theodoros Poulkas, Mikhail Shein, Ruth Signorell\*

### 1 Bulk sample Raman scattering measurements

Single aqueous  $K_2CO_3$  particles are trapped with a counter propagating tweezer and dried while being interrogated with time resolved broad band light scattering (BLS) and Raman scattering measurements. Particle size and refractive index (Cauchy parameters,  $m_0(\lambda)$ ,  $m_1$ ) are determined with BLS measurements. The Raman shift is used to monitor the  $CO_3^{2-}$  symmetric stretching peak position,  $\nu_1(CO_3^{2-})$ . See the main text for the experimental details.<sup>†</sup> Measured values of  $\nu_1(CO_3^{2-})$  are compared with bulk solution measurements in order to determine the particle concentration prior to supersaturation. The bulk aqueous  $K_2CO_3$  solution saturation point is  $8.1 \pm 0.7$  mol  $kg^{-1}$  corresponding to 43% relative humidity (RH) in equilibrium.

**Table 1** Bulk solution measurements.

Molality/ mol $kg^{-1}$	$\nu_1(CO_3^{2-}) / cm^{-1}$
3	1062.4
4	1062.2
6	1061.4
7	1061.0
8	1060.2
Crystalline powders	1060.5

**Table 2** Single particle measurement during a drying experiment. The numbers in the square brackets indicate the 95 % credible interval.

RH/%	$m_0(400nm)$	$m_1$	$\nu_1(CO_3^{2-}) / cm^{-1}$
67.6	1.425 [1.417, 1.437]	0.0236 [0.0196, 0.0276]	1061.31
62.8	1.430 [1.422, 1.438]	0.0228 [0.0196, 0.0260]	1061.20
56.6	1.438 [1.421, 1.446]	0.0248 [0.0208, 0.0296]	1060.84
49.2	1.441 [1.428, 1.454]	0.0252 [0.0208, 0.0288]	1060.51
43.3	1.440 [1.431, 1.454]	0.0260 [0.0220, 0.0292]	1059.99
38.2	1.442 [1.433, 1.451]	0.0272 [0.0240, 0.0308]	1059.59
33.1	1.446 [1.437, 1.460]	0.0268 [0.0228, 0.0308]	1059.11
29.6	1.451 [1.432, 1.461]	0.0260 [0.0212, 0.0312]	1058.55
27.6	1.446 [1.432, 1.460]	0.0268 [0.0220, 0.0312]	1058.26
25.4	1.453 [1.439, 1.468]	0.0272 [0.0216, 0.0328]	1057.68
23.2	1.457 [1.442, 1.472]	0.0284 [0.0232, 0.0340]	1057.39
20.8	1.462 [1.447, 1.477]	0.0264 [0.0208, 0.0324]	1056.93
19.1	1.456 [1.441, 1.472]	0.0288 [0.0232, 0.0336]	1056.37
17.5	1.461 [1.445, 1.476]	0.0276 [0.0220, 0.0344]	1056.00
16.1	1.461 [1.445, 1.477]	0.0272 [0.0216, 0.0324]	1055.88
15.1	1.466 [1.445, 1.476]	0.0280 [0.0228, 0.0340]	1055.94
14.1	1.470 [1.454, 1.480]	0.0280 [0.0228, 0.0340]	1055.64
13.3	1.470 [1.459, 1.486]	0.0280 [0.0228, 0.0332]	1055.23
12.6	1.473 [1.462, 1.484]	0.0272 [0.0228, 0.0328]	1055.18

Laboratory of Physical Chemistry, ETH Zürich, Vladimir-Prelog-Weg 2, CH-8093, Zürich, Switzerland. E-mail: rsignorell@ethz.ch

<sup>†</sup> See DOI: 10.1039/b000000x/

## 2 Initial water uptake during deliquescence

Broad band light scattering (BLS) experiments measure the intensity of light scattered by a single particle as a function of wavelength,  $\lambda$ . Resonances give rise to a ripple structure in the BLS spectrum (see Fig.1 in the main text). The position and the intensity profile of a specific resonance mode depend on the particle shape, the particle radius  $R$  (volume equivalent diameter for non-spherical particles), the wavelength of the light  $\lambda$ , and the refractive index  $m(\lambda)$ . Since the ratio of  $R$  and  $\lambda$  determines the resonance behaviour of the light within the particle it is convenient to define the size parameter  $x = 2\pi N\lambda/R$  where  $N$  is the refractive index of the surrounding medium (here  $N=1$ ).<sup>1</sup> The shift of the resonance position  $\Delta\lambda$  indicates a change of  $R$  because the value of  $x$  at resonance remains constant. The change in  $R$  can then be calculated using Equation (1) under the assumption that the particle's shape and  $m(\lambda)$  do not change appreciably. In the equation  $\lambda_R$  is the resonance position before the deliquescence starts.

$$\Delta R \sim \frac{R\Delta\lambda}{\lambda_R} \quad (1)$$

During the deliquescence experiment shown in Fig.8 in the main text ( $9.5\% \leq \text{RH} \leq 30\%$ ) the peak positions in the BLS spectra drifts by  $\Delta\lambda/\lambda_R = 0.5\%$ . The particle size prior the efflorescence (at  $t_E - 10$  ms) is  $R = 2070.5$  [2052 2086] nm. Hence  $\Delta R$  is  $\sim 10$ nm.

### References

- 1 C. F. Bohren and D. R. Huffman, *Absorption and scattering of light by small particles*, Wiley, Weinheim, Germany, Wiley Professional Paperback edn., 1983.