

# Synthesis of 4-(Arylmethyl)proline Derivatives

**Journal Article****Author(s):**

Loosli, Simon; Foletti, Carlotta; Pappmeyer, Marcus; Wennemers, Helma

**Publication date:**

2019-03-01

**Permanent link:**

<https://doi.org/10.3929/ethz-b-000330184>

**Rights / license:**

[Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International](#)

**Originally published in:**

Synlett 30(4), <https://doi.org/10.1055/s-0037-1611672>

**Funding acknowledgement:**

178805 - Synthetic Collagen (Fortsetzung) (SNF)

# Synthesis of 4-(Arylmethyl)proline Derivatives

Simon Loosli<sup>◊</sup>Carlotta Foletti<sup>◊</sup>


Marcus Pappmeyer

Helma Wennemers\* 

ETH Zürich, Laboratory for Organic Chemistry, D-CHAB,  
Vladimir-Prelog-Weg 3, 8093 Zürich, Switzerland  
Helma.Wennemers@org.chem.ethz.ch

<sup>◊</sup> These authors contributed equally.

Published as part of the 30 Years SYNLETT – Pearl Anniversary  
Issue



Received: 20.12.2018

Accepted after revision: 16.01.2019

Published online: 25.01.2019

DOI: 10.1055/s-0037-1611672; Art ID: st-2018-b0822-l

License terms:

**Abstract** A synthesis of 4-(arylmethyl)proline by using Suzuki cross-couplings was developed. The route permits access to a variety of 4-substituted proline derivatives bearing various aryl moieties that expand the toolbox of proline analogues for studies in chemistry and biology.

**Key words** arylmethylprolines, prolines, hydroboration, Suzuki cross-coupling


Proline is the only proteogenic amino acid with a cyclic backbone, which confers to this residue a uniquely restricted conformation. Nature and scientists have used proline and its derivatives to regulate numerous processes, ranging from ion-gating and the structural integrity of skin to asymmetric catalysis.<sup>1–4</sup> The development of proline analogues and their incorporation into peptides and other compounds is therefore of great interest. Proline derivatives with different substituents at C $\gamma$  are the most common, due to their natural occurrence and the ease of functionalization of (2*S*,4*R*)-4-hydroxyproline.<sup>1,5</sup> Examples include derivatives with heteroatoms at C $\gamma$ , e.g., F, Cl, N<sub>3</sub>, NH<sub>2</sub>, or alkyl groups, e.g., Me and <sup>t</sup>Bu.<sup>1,6</sup> In contrast, derivatives with arylmethyl substituents at C $\gamma$  are less commonly utilized, possibly due to a lack of a straightforward synthetic route.

We became interested in proline derivatives bearing naphthyl moieties, for their value in the molecular recognition of RNA.<sup>7</sup> Synthetic routes have been reported for the functionalization of proline at C $\gamma$  with benzylic or indolylmethyl substituents.<sup>6a,8,9</sup> However, we had limited success


in transferring these reaction conditions, which rely on Wittig reactions of 4-oxoproline followed by hydrogenation, to larger aryl moieties (Scheme 1, top).

We therefore sought an alternative route and we envisioned Suzuki reactions between an organoborane–proline derivative and aryl halides as a strategy that might provide access to proline derivatives with various aryl groups (Scheme 1, bottom). Here, we report a general synthetic route to arylmethyl proline derivatives that permits the introduction of a broad range of aryl moieties at C $\gamma$ .


Our synthetic route relies on the hydroboration of the Boc/<sup>t</sup>Bu-protected 4-methyleneproline **5**, which was obtained from (2*S*,4*R*)-4-hydroxyproline (**1**) by slight modification of a previously published procedure (Scheme 2).<sup>10</sup> This four-step synthesis started with Boc-protection of **1**, followed by oxidation to ketone **3**, protection of the carboxylic acid as the <sup>t</sup>Bu ester in **4**, and introduction of an exocyclic methylene group by a Wittig reaction.<sup>11</sup>



**Scheme 1** Synthetic routes to 4-(arylmethyl)proline derivatives




Hydroboration of the 4-methyleneproline **5** with 9-BBN provided the organoborane **6**, which was used for the Suzuki reaction without further purification (Scheme 3, top). For the Suzuki reaction, various catalysts and conditions were explored by using 2-bromonaphthalene as a model aryl bromide. We focused in particular on catalysts that had proven valuable for cross-couplings with other amino acid derivatives (Scheme 3, bottom).<sup>12</sup> Among the tested palladium-based catalysts, reactions with PEPPSI<sup>13</sup> showed the highest conversion of **5** and 2-bromonaphthalene into the Suzuki reaction product **7a**. Under optimized conditions [5 M aq KOH, ArBr (1.3 equiv), PEPPSI(3% mol)], the 4-(2-naphthylmethyl)proline derivative **7a** was obtained in a yield of 83%. Note that 3 mol% of PEPPSI was enough to obtain these results. Because PEPPSI is more air-stable than other palladium catalysts,<sup>14</sup> this catalyst was used for all further experiments.



Reassuringly, this route also permitted the synthesis of proline derivatives bearing substituted naphthyl moieties (**7b** and **7c**) as well as phenyl (**7d**), 9-anthryl (**7e**), or pyren-1-yl (**7f**) substituents in good overall yields (60–83%; Scheme 3).<sup>15</sup> All derivatives were obtained with a diastereoselectivity of ~3:2 in favor of the *syn*-product, as determined by analysis of <sup>1</sup>H NMR NOE spectroscopy.<sup>11</sup>

Because peptide syntheses typically require Fmoc-protected amino acids, we converted **7a–c** into the respective Fmoc-amino acids **8a–c**. Simultaneous removal of the *t*Bu protecting groups in 6 M HCl in 1,4-dioxane, and subsequent Fmoc-protection afforded **8a–c** in yields of 74–89% (Scheme 4). The diastereoisomers were separated by preparative reverse-phase HPLC to obtain enantiomerically pure amino acids at a scale of up to 2.5 g.<sup>15,16</sup>



In conclusion, we have introduced a synthetic route to access proline derivatives bearing a variety of arylmethyl substituents at the  $\gamma$ -position. The products were obtained in good yields for every tested aromatic moiety. The diastereoselectivity of the hydroboration step was modest, but the diastereoisomeric products could be separated on a gram scale. Installation of a Fmoc-protecting group was straightforward. Thus, the route provides access to proline derivatives with a variety of arylmethyl moieties at  $\gamma$  that are suitably protected for solid-phase peptide synthesis. We envision these derivatives as being valuable additions to the toolkit of proline analogues for applications in chemistry and chemical biology.

## Funding Information

The authors gratefully acknowledge financial support by the Swiss National Science Foundation (grant 200020\_178805).

## Supporting Information

Supporting information for this article is available online at <https://doi.org/10.1055/s-0037-1611672>.

## References and Notes

- Shoulders, M. D.; Raines, R. T. *Annu. Rev. Biochem.* **2009**, *78*, 929.
- For examples, see: (a) Arnold, U.; Hinderaker, M. P.; Köditz, J.; Golbik, R.; Ulbrich-Hofmann, R.; Raines, R. T. *J. Am. Chem. Soc.* **2003**, *125*, 7500. (b) Lummis, S. C.; Beene, D. L.; Lee, L. W.;

- Lester, H. A.; Broadhurst, R. W.; Dougherty, D. A. *Nature* **2005**, *438*, 248. (c) Lieblich, S. A.; Fang, K. Y.; Cahn, J. K. B.; Rawson, J.; LeBon, J.; Ku, H. T.; Tirrell, D. A. *J. Am. Chem. Soc.* **2017**, *139*, 8384. (d) Metrano, A. J.; Abascal, N. C.; Mercado, B. Q.; Paulson, E. K.; Hurtley, A. E.; Miller, S. J. *J. Am. Chem. Soc.* **2017**, *139*, 492.
- (3) Liu, J.; Wang, L. *Synthesis* **2017**, 49, 960.
- (4) Lewandowski, B.; Wennemers, H. *Curr. Opin. Chem. Biol.* **2014**, *22*, 40.
- (5) (a) Remuzon, P. *Tetrahedron* **1996**, *52*, 13803. (b) Pandey, A. K.; Naduthambi, D.; Thomas, K. M.; Zondlo, N. J. *J. Am. Chem. Soc.* **2013**, *135*, 4333.
- (6) (a) Del Valle, J. R.; Goodman, M. J. *Org. Chem.* **2003**, *68*, 3923. (b) Koskinen, A. M. P.; Helaja, J.; Kumpulainen, E. T. T.; Koivisto, J.; Mansikkamäki, H.; Rissanen, K. J. *Org. Chem.* **2005**, *70*, 6447.
- (7) Foletti, C.; Kramer, R. A.; Mauser, H.; Jenal, U.; Bleicher, K. H.; Wennemers, H. *Angew. Chem. Int. Ed.* **2018**, *57*, 7729.
- (8) (a) Ezquerro, J.; Pedregal, C.; Yrretagoyena, B.; Rubio, A.; Carreño, M. C.; Escribano, A.; García Ruano, J. L. *J. Org. Chem.* **1995**, *60*, 2925. (b) Rawson, D. J.; Brugier, D.; Harrison, A.; Hough, J.; Newman, J.; Otterburn, J.; Maw, G. N.; Price, J.; Thompson, L. R.; Turnpenny, P.; Warren, A. N. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 3771.
- (9) Krapcho, J.; Turk, C.; Cushman, D. W.; Powell, J. R.; DeForrest, J. M.; Spitzmiller, E. R.; Karanewsky, D. S.; Duggan, M.; Rovnyak, G.; Schwartz, J.; Natarajan, S.; Godfrey, J. D.; Ryono, D. E.; Neubeck, R.; Atwal, K. S.; Petrillo, E. W. Jr. *J. Med. Chem.* **1988**, *31*, 1148.
- (10) (a) Herdewijn, P.; Claes, P. J.; Vanderhaeghe, H. *Can. J. Chem.* **1982**, *60*, 2903. (b) De Luca, L.; Giacomelli, G.; Porcheddu, A. *Org. Lett.* **2001**, *3*, 3041.
- (11) For details, see the Supporting Information.
- (12) For the use of the Suzuki reaction to prepare amino acids, see: (a) Campbell, A. D.; Raynham, T. M.; Taylor, R. J. K. *Tetrahedron Lett.* **1999**, *40*, 5263. (b) Sabat, M.; Johnson, C. R. *Org. Lett.* **2000**, *2*, 1089. (c) Lu, X.; Xiao, B.; Shang, R.; Liu, L. *Chin. Chem. Lett.* **2016**, *27*, 305.
- (13) O'Brien, C. J.; Kantchev, E. A. B.; Valente, C.; Hadei, N.; Chass, G. A.; Lough, A.; Hopkinson, A. C.; Organ, M. G. *Chem. Eur. J.* **2006**, *12*, 4743.
- (14) Ray, L.; Shaikh, M. M.; Ghosh, P. *Dalton Trans* **2007**, 4546.
- (15) **tert-Butyl (4S/4R)-N-(tert-Butoxycarbonyl)-4-(2-naphthylmethyl)-L-prolinate (7a); Typical Procedure**  
An oven-dried Schlenk flask was charged with methylene derivative **5** (4.0 g, 14.1 mmol, 1 equiv) under N<sub>2</sub>. A 0.5 M soln of 9-BBN in THF (31.0 mL, 15.5 mmol, 1.1 equiv) was added in one portion, and the solution was stirred vigorously at 60 °C for 6 h. The mixture was then allowed to cool to r.t. and 5 M aq. KOH (5.6 mL, 5 M, 28.0 mmol, 2 equiv) was added. The mixture was stirred for 20 min, then 2-bromonaphthalene (**7a**; 3.8 g, 18.36 mmol, 1.3 equiv) was added together with PEPPSI (287.7 mg, 423 μmol, 0.03 equiv). The mixture was stirred for a further 16 h at r.t., then H<sub>2</sub>O (120 mL) and EtOAc (120 mL) were added and the phases were separated. The aqueous phase was extracted with EtOAc (3 × 120 mL), and the organic layers were combined, washed with brine, dried (MgSO<sub>4</sub>), and concentrated. The resulting yellow-brown oil (9.9 g) was purified by column chromatography (silica gel, 0–25% EtOAc–hexane) to give a colorless oil; yield: 4.8 g (83%).  
<sup>1</sup>H NMR (500 MHz, C<sub>2</sub>Cl<sub>4</sub>D<sub>2</sub>, 60 °C): δ = 7.82–7.71 (m, 3 H), 7.59–7.52 (m, 1 H), 7.48–7.37 (m, 2 H), 7.27 (dd, J = 8.4, 1.7 Hz, 1 H), 4.26–4.02 (m, 1 H), 3.75–3.55 (m, 1 H), 3.14 (dd, J = 10.6, 9.0 Hz, 1 H), 2.89–2.76 (m, 2 H), 2.72–2.44 (m, 1 H), 2.42–1.88 (m, 1 H), 1.63 (ddd, J = 12.8, 9.5, 7.9 Hz, 1 H), 1.51–1.33 (m, 18 H). <sup>13</sup>C NMR (126 MHz, C<sub>2</sub>Cl<sub>4</sub>D<sub>2</sub>, 60 °C): δ = 172.2, 172.0, 153.6, 137.6, 137.4, 133.5, 133.5, 132.1, 128.1, 128.1, 127.6, 127.5, 127.4, 127.2, 127.1, 126.8, 126.8, 126.1, 125.4, 80.9, 80.8, 79.6, 79.5, 59.8, 59.7, 52.1, 51.6, 39.3, 39.2, 37.7, 36.7, 36.4, 28.4, 28.0, 28.0. HRMS (ESI+): m/z [M + H]<sup>+</sup> calcd C<sub>25</sub>H<sub>34</sub>NO<sub>4</sub>: 412.2482; found: 412.2485.
- (4S)- and (4R)-1-[(9H-Fluoren-9-ylmethoxy)carbonyl]-4-(2-naphthylmethyl)-L-proline (8a); Typical Procedure**  
Prolinate **7a** (4.8 g, 11.7 mmol, 1 equiv) was dissolved in a 6 M soln of HCl in 1,4-dioxane (110 mL), and the mixture was stirred for 3 h at r.t. The pH was adjusted to 8–9 with sat. aq. NaHCO<sub>3</sub>, then a soln of FmocCl (3.6 g, 14.0 mmol, 1.2 equiv) in 1,4-dioxane (50 mL) was added, and the mixture was stirred at r.t. for 2 h. Low-boiling volatiles were removed under reduced pressure, and EtOAc (50 mL) was added. The solution was acidified to pH 2–3 with 1 M HCl, and the organic phase was separated and extracted with EtOAc (3 × 100 mL). The combined organic layers were washed with brine, dried (MgSO<sub>4</sub>), and filtered. All volatiles were removed under reduced pressure, and the product was purified by column chromatography (silica gel, 0–5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> with 0.1% HCO<sub>2</sub>H) to give a white powder; yield: 4.7 g (84%). The diastereoisomers were subsequently separated by reverse-phase semipreparative HPLC [Reprosil-Gold 120 C18, 10 μm; 250 × 30 mm column, MeCN and H<sub>2</sub>O–MeCN–TFA (100:1:0.1)].
- (4S)-Diastereomer**  
[α]<sub>D</sub> –40.7 ± 0.5 (c 0.2, MeOH). TLC (silica gel, 2% MeOH in CH<sub>2</sub>Cl<sub>2</sub>): R<sub>f</sub> = 0.56. FTIR (neat): 3051, 2923, 1701, 1421, 1352, 1247, 1176, 1122, 1006, 972, 843, 739 cm<sup>-1</sup>.  
<sup>1</sup>H NMR (500 MHz, C<sub>2</sub>Cl<sub>4</sub>D<sub>2</sub>, 60 °C): δ = 7.92–7.79 (m, 3 H; Ar), 7.77–7.60 (m, 3 H; Ar), 7.59–7.44 (m, 4 H; Ar), 7.43–7.21 (m, 5 H; Ar), 4.55–4.41 (m, 2 H; CH<sub>2</sub>-Fmoc), 4.41–4.28 (m, 1 H; H<sub>α</sub>), 4.28–4.18 (m, 1 H; CH-Fmoc), 3.77–3.52 (m, 1 H; H<sub>δ</sub>), 3.25–3.16 (m, 1 H; H<sub>δ</sub>), 3.01–2.79 (m, 2 H; CH<sub>2</sub>-Naph), 2.57 (hept, J = 7.7 Hz, 1 H; H<sub>γ</sub>), 2.50–2.36 (m, 1 H; H<sub>β</sub>), 2.12–1.93 (m, 1 H; H<sub>β</sub>). <sup>13</sup>C NMR (500 MHz, C<sub>2</sub>Cl<sub>4</sub>D<sub>2</sub>, 60 °C): δ = 173.2 (CO<sub>2</sub>H), 156.4 (C=O<sub>Fmoc</sub>), 143.5 (Ar), 141.1 (Ar), 137.0 (Ar), 133.4 (Ar), 132.1 (Ar), 128.2 (Ar), 127.6 (Ar), 127.5 (Ar), 127.4 (Ar), 127.0 (Ar), 126.9 (Ar), 126.7 (Ar), 126.1 (Ar), 125.5 (Ar), 124.8 (Ar), 119.8 (Ar), 67.9 (CH<sub>2</sub>-Fmoc), 59.4 (C<sub>α</sub>), 52.2 (C<sub>δ</sub>), 47.1 (CH-Fmoc), 39.7 (C<sub>γ</sub>), 38.8 (CH<sub>2</sub>-Naph), 34.4 (C<sub>β</sub>). HRMS (ESI+): m/z [M + H]<sup>+</sup> calcd for C<sub>31</sub>H<sub>28</sub>NO<sub>4</sub>: 478.2013; found: 478.2003.
- (4R)-Diastereomer**  
[α]<sub>D</sub> –10.8 ± 0.3 (c 0.2, MeOH). TLC (silica gel, 2% MeOH in CH<sub>2</sub>Cl<sub>2</sub>): R<sub>f</sub> = 0.56. FTIR (neat): 3045, 2966, 1700, 1661, 1417, 1351, 1241, 1282, 1122, 1002, 947, 887, 737 cm<sup>-1</sup>.  
<sup>1</sup>H NMR (500 MHz, C<sub>2</sub>Cl<sub>4</sub>D<sub>2</sub>, 60 °C): δ = 7.91–7.80 (m, 3 H; Ar), 7.73 (dd, J = 7.6, 2.9 Hz, 2 H; Ar), 7.62 (s, 1 H; Ar), 7.59–7.48 (m, 4 H; Ar), 7.39 (tt, J = 7.6, 1.4 Hz, 2 H; Ar), 7.35–7.27 (m, 3 H; Ar), 4.56–4.36 (m, 3 H; H<sub>α</sub>, CH<sub>2</sub>-Fmoc), 4.32–4.19 (m, 1 H; CH-Fmoc), 3.74–3.49 (m, 1 H; H<sub>δ</sub>), 3.31–3.10 (m, 1 H; H<sub>δ</sub>), 2.97–2.80 (m, 2 H; CH<sub>2</sub>-Naph), 2.80–2.65 (m, 1 H; H<sub>γ</sub>), 2.47–1.88 (m, 2 H; H<sub>β</sub>). <sup>13</sup>C NMR (500 MHz, C<sub>2</sub>Cl<sub>4</sub>D<sub>2</sub>, 60 °C): δ = 173.8 (CO<sub>2</sub>H), 156.1 (C=O<sub>Fmoc</sub>), 143.5 (Ar), 141.1 (Ar), 136.8 (Ar), 133.4 (Ar), 132.1 (Ar), 128.2 (Ar), 127.6 (Ar), 127.5 (Ar), 127.4 (Ar), 127.0 (Ar), 127.0 (Ar), 126.8 (Ar), 126.1 (Ar), 125.5 (Ar), 124.8 (Ar), 119.8 (Ar), 67.9 (CH<sub>2</sub>-Fmoc), 59.2 (C<sub>α</sub>), 51.7 (C<sub>δ</sub>), 47.1 (CH-Fmoc), 38.9 (CH<sub>2</sub>-Naph, C<sub>γ</sub>), 34.3 (C<sub>β</sub>). HRMS (ESI+): m/z [M + H]<sup>+</sup> calcd C<sub>31</sub>H<sub>28</sub>NO<sub>4</sub>: 478.2013; found: 478.2003.
- (16) Note that the Suzuki reaction is not compatible with the use of Fmoc-protected amines. The stereochemistry at the stereogenic centers was retained during the synthesis.