

Neuronal response variability as a product of divisive normalization; neurobiological implications at a macroscale level

Journal Article**Author(s):**

Ruddy, Kathy L.; Cole, David M.; Simon, Colin; Bächinger, Marc T.

Publication date:

2020

Permanent link:

<https://doi.org/10.3929/ethz-b-000462125>

Rights / license:

[Creative Commons Attribution 4.0 International](#)

Originally published in:

HRB Open Research 3, <https://doi.org/10.12688/hrbopenres.13062.1>



CORRESPONDENCE

Neuronal response variability as a product of divisive normalization; neurobiological implications at a macroscale level [version 1; peer review: 2 approved]

Kathy L. Ruddy ^{1*}, David M. Cole^{2*}, Colin Simon ¹, Marc T. Bächinger³¹Trinity College Institute of Neuroscience and School of Psychology, Trinity College Dublin, Dublin, Dublin 2, Ireland²Interdisciplinary Program in Neuroscience,, Utah State University, Logan, Utah, USA³Neural Control of Movement Lab, Department of Health Sciences and Technology, ETH Zürich, Zurich, Switzerland

* Equal contributors

V1 First published: 04 Jun 2020, 3:34
<https://doi.org/10.12688/hrbopenres.13062.1>
Latest published: 04 Jun 2020, 3:34
<https://doi.org/10.12688/hrbopenres.13062.1>

Abstract

The occurrence of neuronal spikes recorded directly from sensory cortex is highly irregular within and between presentations of an invariant stimulus. The traditional solution has been to average responses across many trials. However, with this approach, response variability is downplayed as noise, so it is assumed that statistically controlling it will reveal the brain's true response to a stimulus. A mounting body of evidence suggests that this approach is inadequate. For example, experiments show that response variability itself varies as a function of stimulus dimensions like contrast and state dimensions like attention. In other words, response variability has structure, is therefore potentially informative and should be incorporated into models which try to explain neural encoding. In this article we provide commentary on a recently published study by Coen-Cagli and Solomon that incorporates spike variability in a quantitative model, by explaining it as a function of divisive normalization. We consider the potential role of neural oscillations in this process as a potential bridge between the current microscale findings and response variability at the mesoscale/macroscale level.

Keywords

Spiking, neuronal variability, neuronal oscillations

Open Peer Review

Reviewer Status

Invited Reviewers

1

2

version 1

04 Jun 2020

report

report

1. **Lucia Li** , Imperial College London, London, UK2. **Valentine L Marcar**, University of Zürich, University Hospital Zürich, Zürich, Switzerland

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Kathy L. Ruddy (ruddykl@tcd.ie)

Author roles: **Ruddy KL:** Conceptualization, Funding Acquisition, Investigation, Supervision, Writing – Original Draft Preparation; **Cole DM:** Investigation, Writing – Original Draft Preparation; **Simon C:** Investigation, Writing – Review & Editing; **Bächinger MT:** Investigation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Health Research Board, Ireland [HRB-EIA-2019-003]. Kathy Ruddy would also like to acknowledge grant funding from the Irish Research Council GOIPD/2017/798.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Ruddy KL *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Ruddy KL, Cole DM, Simon C and Bächinger MT. **Neuronal response variability as a product of divisive normalization; neurobiological implications at a macroscale level [version 1; peer review: 2 approved]** HRB Open Research 2020, 3 :34 <https://doi.org/10.12688/hrbopenres.13062.1>

First published: 04 Jun 2020, 3:34 <https://doi.org/10.12688/hrbopenres.13062.1>

The neural coding problem asks how richly graded stimuli are translated into action potentials. A notorious complication is that single spike trains do not follow a predictable schedule. Despite the stereotyped shape of spikes, their occurrence is highly irregular within and between stimulus presentations (Kostal *et al.*, 2007). The traditional solution has been to average responses across many trials. However with this approach, response variability is downplayed as noise, so it is assumed that statistically controlling it will reveal the brain's true response to a stimulus (Kostal *et al.*, 2007). Accordingly, suspicions have long been harbored that averaging is an incomplete answer. For example, experiments show that response variability itself varies as a function of stimulus dimensions like contrast and state dimensions like attention (Cohen & Maunsell, 2009; Hénaff *et al.*, 2019; Mitchell *et al.*, 2009; Rabinowitz *et al.*, 2015; Ruff & Cohen, 2016). In other words, response variability has structure, is therefore potentially informative and should be incorporated into models which try to explain neural encoding. Coen-Cagli & Solomon (2019) take a unique approach to both quantitatively describe spike variability, and relate it to a specific mechanism – *divisive normalization*.

In its basic formulation, divisive normalization refers to dividing the driving stimulus input by the pooled drive of an ensemble of inputs (i.e. the normalization signal). In a seminal writing, Heeger (1991) showed that divisive normalization can increase the dynamic range of primary visual cortex (V1) neurons to contrast and thereby improve contrast sensitivity under varying light conditions. Additionally, since it has been observed in multiple brain areas and circuits across species, it has been put forth as a canonical neural computation (i.e. similar to exponentiation and linear filtering (Carandini & Heeger, 2011)).

However, in the standard model of divisive normalization, the driving stimulus input and the normalization signal are deterministic variables (Coen-Cagli & Solomon, 2019, Eq. 1.1). This simplification either ignores response variability or operates as if it has been averaged out. Coen-Cagli & Solomon (2019) integrate response variability into their model by replacing them with Gaussian distributions (Eq. 1.2). Then, because this idealized mathematical expression is computationally intractable, they generated approximate expressions for spiking mean and spiking variance as predicted by their model (Eq. 1.3–1.4). These approximations then allowed them to test their model's predictions against real data. To do so they presented images of gratings of varying contrasts to anesthetized macaque monkeys while recording activity from V1 with microelectrodes. Thereby, they were able to verify that their model similarly predicted the effect of contrast on spiking mean.

The key finding of Coen-Cagli & Solomon (2019) was that normalization is inversely related to (a) spiking variability and (b) inter-neuronal correlation of variability. While offering a neurobiological mechanistic explanation of this effect was beyond the scope of their investigation, they cite inhibitory stabilization of network activity and specifically GABA_A

inhibition as potential candidates providing a normalizing role. This has prompted us to consider the potential role of neural oscillations in this process, and particularly their dependence on inhibition, as a potential bridge between the current microscale findings and response variability at the mesoscale/macroscale level.

In considering how neural oscillations may be a key linking factor, it is essential first to understand the role of oscillation for constraining (and allowing) neuronal spiking to occur. Traditionally considered as an epiphenomenon or by-product of neuronal firing, it is now conversely believed that neuronal oscillations are more causal than consequential in driving neuronal activity (eg. Thut *et al.*, 2012). Neurons are inherently sensitive to noise in the local brain environment, including fluctuations in temperature, pH, and nearby electrical potential changes. According to Buzsaki (2006), if the neuron's membrane potential was fixed close to its firing threshold most of the time, then it may risk frequent accidental discharge due to small changes in excitation from the fluctuating background noise. On the contrary, if the membrane potential was fixed far below threshold, such that it would be unaffected by background noise, the energy requirement to depolarize the cell would be much too costly on the brain's resources. Buzsaki (2006) suggests that the brain's most energetically economical solution is to swing the membrane potentials of coordinated populations of neurons up and down, providing windows of increased and decreased likelihood of firing. This centrally coordinated threshold-adjusting mechanism is carried out by the interneuron system, by providing periodic bursts of inhibition, the duration of which determines the frequency of the oscillation. For example, if the inhibition is mediated by GABA_A receptors, the emergent oscillation is in the Gamma frequency (40–100 Hz).

Thus, oscillations constrain neuronal spiking into periodic windows of varying responsiveness, and hence may serve as a determinant of response variability. As in Coen-Cagli & Solomon (2019) this may manifest as unreliable spike responses to invariant external (sensory) stimulation. In this regard, a correspondence between the micro and macroscopic level can be made, as the same neuronal oscillations have been shown to systematically modulate characteristics of responses to repeated invariant sensory stimuli measured at the behavioral level (Busch *et al.*, 2009; Mathewson *et al.*, 2009), and in muscle responses to exogenous transcranial magnetic stimulation (Ruddy *et al.*, 2018; Zarkowski *et al.*, 2006).

For example, when measuring ongoing neural oscillations simultaneously during sensory or transcranial stimulation, the influence of the underlying oscillation on the variability of resulting responses can be demonstrated at multiple levels of the nervous system. In the motor system, when using TMS to stimulate primary motor cortex, motor evoked potentials (MEPs) in corresponding muscles are larger when the pulse arrives coinciding with low amplitude alpha (8–13 Hz) waves, or high amplitude gamma (30–80 Hz) waves recorded

at the scalp (Ruddy *et al.*, 2018; Zarkowski *et al.*, 2006) despite the intensity of the stimulation remaining fixed. Further, within the alpha rhythm recorded over sensorimotor cortex (often referred to in this context as the μ rhythm), it has been shown that MEPs are modulated systematically by the phase of the wave cycle; they are larger when stimulation occurs at the trough of the wave and smaller when at the peaks (Schaworonkow *et al.*, 2019; Zrenner *et al.*, 2018). At a behavioural level, in the visual system when presented with the same (low contrast) stimulus on multiple trials, detection is higher when the underlying alpha oscillation in the visual cortex is at the trough of the wave cycle, rather than at the peak, and when the amplitude of the rhythm is generally low (Busch *et al.*, 2009; Mathewson *et al.*, 2009). Similar results have been obtained in investigations of variability in visual evoked potentials (Brandt & Jansen, 2009) and auditory evoked potentials (Rahn & Basar, 2009).

Considering the aforementioned evidence, it is clear that background fluctuations in the brain's electrical activity contribute substantially to the variation in responses recorded

following repeated presentations of an invariant stimulus. Buzsaki (2006) claims that this apparently spontaneous brain-state variability is internally coordinated and may be the essence of cognition itself.

Coen-Cagli & Solomon (2019) provide a novel approach to account for modulations in neuronal response variability in a quantitative and descriptive fashion by demonstrating the role divisive normalization could play. It is unclear at present whether neural oscillations and the process represented mathematically by divisive normalization are mutually exclusive concepts. Modelling the additional influence of a dynamic oscillating normalization signal and reconciling this with the extant descriptions of interacting neurobiological and electrophysiological mechanisms is a challenge for the future of this field. Resolving this would ultimately contribute further towards bridging across different scales of measurement in neuroscience.

Data availability

No data are associated with this article.

References

- Brandt ME, Jansen BH: **The Relationship Between Prestimulus Alpha Amplitude and Visual Evoked Potential Amplitude.** *Int J Neurosci.* 2009; **61**(3–4): 261–268.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Busch NA, Dubois J, VanRullen R: **The phase of ongoing EEG oscillations predicts visual perception.** *J Neurosci.* 2009; **29**(24): 7869–7876.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Buzsaki G: **Rhythms of the Brain.** United States. Oxford University Press. 2006.
[Publisher Full Text](#)
- Carandini M, Heeger DJ: **Normalization as a canonical neural computation.** *Nat Rev Neurosci.* 2011; **13**(1): 51–62.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Coen-Cagli R, Solomon SS: **Relating divisive normalization to neuronal response variability.** *J Neurosci.* 2019; **39**(37): 7344–7356.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Cohen MR, Maunsell JHR: **Attention improves performance primarily by reducing interneuronal correlations.** *Nat Neurosci.* 2009; **12**(12): 1594–600.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Heeger DJ: **Nonlinear model of neural responses in cat visual cortex.** In: *Computational models of visual processing* (Landy MS, Movshon JA, eds) Cambridge, MA: MIT Press. 1991; 119–133.
[Reference Source](#)
- Hénaff OJ, Boudry-Singer ZM, Meding K, *et al.*: **Representation of uncertainty in macaque visual cortex.** *bioRxiv.* 2019; **92**: 724021.
[Publisher Full Text](#)
- Kostal L, Lansky P, Rospars JP: **Neuronal coding and spiking randomness.** *Eur J Neurosci.* 2007; **26**(10): 2693–2701.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mathewson KE, Gratton G, Fabiani M, *et al.*: **To see or not to see: prestimulus alpha phase predicts visual awareness.** *J Neurosci.* 2009; **29**(9): 2725–2732.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mitchell JF, Sundberg KA, Reynolds JH: **Spatial attention decorrelates intrinsic activity fluctuations in macaque area V4.** *Neuron.* 2009; **63**(6): 879–878.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rabinowitz NC, Goris RL, Cohen M, *et al.*: **Attention stabilizes the shared gain of V4 populations.** *eLife.* 2015; **4**: e08998.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rahn E, Basar E: **Prestimulus EEG-Activity Strongly Influences the Auditory Evoked Vertex Response: A New Method for Selective Averaging.** *Int J Neurosci.* 2009; **69**(1–4): 207–220.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ruddy K, Balsters J, Mantini D, *et al.*: **Neural activity related to volitional regulation of cortical excitability.** *eLife.* 2018; **7**: e40843.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ruff DA, Cohen MR: **Stimulus dependence of correlated variability across cortical areas.** *J Neurosci.* 2016; **36**(28): 7546–7556.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Schaworonkow N, Triesch J, Ziemann U, *et al.*: **EEG-triggered TMS reveals stronger brain state-dependent modulation of motor evoked potentials at weaker stimulation intensities.** *Brain Stimul.* 2019; **12**(1): 110–118.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Thut G, Miniussi C, Gross J: **The Functional Importance of Rhythmic Activity in the Brain.** *Curr Biol.* 2012; **22**(16): R658–63.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zarkowski P, Shin CJ, Dang T, *et al.*: **EEG and the Variance of Motor Evoked Potential Amplitude.** *Clin EEG Neurosci.* 2006; **37**(3): 247–251.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zrenner C, Desideri D, Belardinelli P, *et al.*: **Real-time EEG-defined excitability states determine efficacy of TMS-induced plasticity in human motor cortex.** *Brain Stimul.* 2018; **11**(2): 374–389.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 1

Reviewer Report 24 November 2020

<https://doi.org/10.21956/hrbopenres.14158.r28246>

© 2020 Marcar V. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Valentine L Marcar

Department of Psychology, University of Zürich, University Hospital Zürich, Zürich, Switzerland

The authors comment on the work published in the Journal of Neuroscience by Coen-Cagli and Solomon on their observation that as normalisation of the neuronal response increases, variability in discharge activity decreases. The paper by Rudd and co-authors offers neural oscillation as a plausible explanation for the observation of Coen-Cagli and Solomon. Rudd and colleagues use findings in the literature involving studies on human volunteers to back their explanation.

While their arguments favouring neural oscillations and divisive normalisation as the underlying mechanism influencing response variability is clear, the link between neural oscillation and divisive normalisation is not. This issue would benefit from more extensive attention. It would also be helpful if the role of GABA_A or GABA_B-receptors in regulating neural oscillations would be dealt with.

The paper is well written, with arguments centring on divisive normalisation and neural oscillation clearly presented.

Is the rationale for commenting on the previous publication clearly described?

Yes

Are any opinions stated well-argued, clear and cogent?

Yes

Are arguments sufficiently supported by evidence from the published literature or by new data and results?

Yes

Is the conclusion balanced and justified on the basis of the presented arguments?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Brain imaging using fMRI, NIRS and ERP. Link between stimulus property, neural response and the BOLD-signal and VEP.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 26 June 2020

<https://doi.org/10.21956/hrbopenres.14158.r27503>

© 2020 Li L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Lucia Li 

Department of Brain Sciences, National Institute for Health Research, Imperial College London, London, UK

This is a commentary on a recently published research article (in JoN) that provides model and macaque validation for the theory that divisive normalisation modulates response variability.

The JoN paper finds that stronger normalisation of a neuron is associated with less variable response. The JoN paper does not posit the mechanism for this finding. The commentary here by Rudd *et al.* puts forward the theory that neural oscillations are a possible mechanism to explain this finding.

The authors use findings from the non-invasive brain stimulation and sensory detection literature to expand on the theory.

It is a well-written piece, which clearly and succinctly introduces two important concepts – divisive normalisation and neural oscillations. It also neatly summarises the paper on which it is commenting.

As the authors point out, although both appear to be mechanisms underlying variability of neuronal response, it is not clear how neural oscillations and divisive normalisation links up. It would be helpful if the authors could expand on how the two might be linked e.g. is there evidence that strongly normalised neurons are those that have resting thresholds furthest away from the depolarisation threshold? Or is the link at a higher level, where high activity in GABA interneurons coincides with when neurones are most strongly normalised? Even if there is not much hard evidence to link them, it would be helpful to know the 'state of play'.

Is the rationale for commenting on the previous publication clearly described?

Yes

Are any opinions stated well-argued, clear and cogent?

Yes

Are arguments sufficiently supported by evidence from the published literature or by new data and results?

Yes

Is the conclusion balanced and justified on the basis of the presented arguments?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Neuroscience, brain stimulation

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.
