ETH zürich

Does rated visual landscape quality match visual features? An analysis for renewable energy landscapes

Journal Article

Author(s): <u>Spielhofer, Reto</u> (1); Hunziker, Marcel; Kienast, Felix; Wissen Hayek, Ulrike; <u>Grêt-Regamey, Adrienne</u> (1)

Publication date: 2021-05

Permanent link: https://doi.org/10.3929/ethz-b-000465535

Rights / license: Creative Commons Attribution 4.0 International

Originally published in: Landscape and Urban Planning 209, <u>https://doi.org/10.1016/j.landurbplan.2020.104000</u>

Funding acknowledgement: 173808 - ENERGYSCAPE: Landscape strategy for renewable energy systems (SNF)



Contents lists available at ScienceDirect

Landscape and Urban Planning



journal homepage: www.elsevier.com/locate/landurbplan

Research Paper

Does rated visual landscape quality match visual features? An analysis for renewable energy landscapes



Reto Spielhofer^{a,*}, Marcel Hunziker^b, Felix Kienast^c, Ulrike Wissen Hayek^a, Adrienne Grêt-Regamey^a

^a Planning of Landscape and Urban Systems (PLUS), Institute for Spatial and Landscape Development, ETH Zurich, 8093 Zurich, Switzerland ^b Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Economics and Social Sciences, Social Sciences in Landscape Research Group, 8903

Birmensdorf, Switzerland

^c Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Land Change Science, Land-Use-Systems Group, 8903 Birmensdorf, Switzerland

HIGHLIGHTS

• Rated coherence and legibility correlate with preference for energy landscapes.

- Rated visual landscape qualities and measured visual features are correlated.
- Landscape preference decreases with increasing renewable energy systems.

• Flatlands revealed low preference regardless of share of renewable energy systems.

• Rated landscape coherence is an index of place-technology-fit in energy landscapes.

ARTICLE INFO	A B S T R A C T
<i>Keywords:</i> Renewable energy systems Visual impact assessment Landscape perception Visual features	Finding the "right" sites for developing renewable energy systems (RES) is one of the major challenges in planning strategies for energy transitions. The visibility aspects of such infrastructure are important factors that explain local opposition. Classical visibility and viewshed analyses of RES disregard people's perceptions and estimations of new infrastructure. To address this void, we demonstrate an approach that combines rated visual landscape qualities with measured visual features. In doing so, we established visual stimuli with systematically controlled visual impact scenarios featuring the use of renewables in different landscape types. The study investigated how ratings of landscape qualities are affected by landscape qualities. Finally, we intended to improve the understand of how rated landscape qualities lead to preferences for different RES visual impact scenarios. Our results showed that rated coherence is strongly influenced by renewable energy infrastructure, whereas complexity ratings are affected mainly by variations in landscape types. These findings let us to conclude that the visual understanding and visual connectedness between energy systems and surrounding landscapes are core drivers of people's visual preferences for landscapes altered with RES. Considering landscape qualities within impact assessments of RES can augment our grasp of how the visual character of a landscape changes through renewable energy infrastructure.

1. Introduction

To reach global climate targets and close the energy gaps resulting from phasing out nuclear or fossil energy production, countries increasingly focus on renewable energy resources such as wind, hydropower, solar or biomass (Mathiesen, Lund, & Karlsson, 2011; Wüstenhagen, Wolsink, & Bürer, 2007). In 2016, the production of wind and solar energy amounted to a global volume of only 1.5 Mio. GWh, whereas coal remained the major source of global electricity production (9 Mio. GWh, (World Energy Council, 2019) In the last three decades, the worldwide electricity demand has doubled from 11'000 TWh to 23'000 TWh and is expected to grow annually by up to 1% during the

https://doi.org/10.1016/j.landurbplan.2020.104000

Received 12 May 2020; Received in revised form 29 September 2020; Accepted 17 November 2020 Available online 25 January 2021

0169-2046/© 2020 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

^{*} Corresponding author at: Institute for Spatial and Landscape Development, Planning of Landscape and Urban Systems – PLUS, Hönggerberg HIL H 52.2, Stefano-Franscini-Platz 5, 8093 Zürich, Switzerland.

E-mail address: spreto@ethz.ch (R. Spielhofer).

next decades (World Energy Council, 2019). Addressing growing energy demand combined with an enhanced focus on renewable energy, requires an increasing amount of land for energy production. Let us take Switzerland, which served as the study site in this work, a technical--economic production potential of approx. 14 TWh from wind energy and approx. 58 TWh of photovoltaic energy (Cattin et al., 2012) are estimated. Today, Switzerland annually produces 0.16 TWh and 2.2 TWh of electricity from wind and solar energy, respectively (Swiss Federal Office of Energy, 2019). According to the goals indicated in the "Swiss Energy Strategy 2050" the production of electricity from renewable energy should amount to 11.4 TWh in 2035 and 24.2 TWh in 2050. With these developments come fundamental transformations that are expected to reshape ecosystem services at the landscape scale (e.g., biodiversity, recreation, food production) (Nadaï and van der Horst, 2010; Plieninger et al., 2016). Several services have been shown to conflict with the production of renewable energy (e.g., Hastik et al., 2015; Kienast, Frick, van Strien, & Hunziker, 2017; Santangeli et al., 2016). Exemplarily for Switzerland, Kienast et al. (2017) estimate that on 88% of the potential locations for wind energy production and on 32% of the potential areas for solar panels, compete with one or more other services. Therefore, the conflict of wind (Warren, Lumsden, O'Dowd, & Birnie, 2005; Wolsink, 2007) and photovoltaic energy infrastructure (Chiabrando, Fabrizio, & Garnero, 2009; Zoellner, Schweizer-Ries, & Wemheuer, 2008) with landscape aesthetics is an important factor for the decreased social acceptance of such projects. Over the last decades, studies have revealed that, in general, people visually prefer natural landscapes over built environments (e.g., Kaplan and Kaplan, 1989; Ulrich, 1986; Ulrich et al., 1991). Given that RES are artificial installations, studies conclude that adding RES to landscapes generally diminish visual landscape quality ratings (Zoellner et al., 2008). Specifically, in landscapes perceived as having high visual quality, the presence of RES strongly reduces visual quality compared to areas of low quality (Betakova, Vojar, & Sklenicka, 2015; Lothian, 2008; Molnarova et al., 2012). Although standardized visual impact assessments (VIA) are widely used in practice to quantify RES visual impact, these methods cannot illuminate why people visually prefer certain landscapes altered with RES over others. Correspondingly, the majority of VIAs quantify the visibility of RES instead of assessing RES siting-induced changes in the visual quality of landscapes (Wolsink, 2018). Inquiring into the visual qualities of RES-altered landscapes can elevate our understanding of the preference forming process for such landscapes. The insights accordingly derived can also help landscape planners and designers involve people in decision making on RES projects.

The overarching goal of this study was to investigate how people perceive RES-driven visual changes in different landscapes. All the goals pursued in this work, along with two concepts that constituted the theoretical framework are detailed in the succeeding sub-sections.

1.1. Perceiving and rating landscape qualities

Landscape perception is defined as "the seeing of [landscape] qualities" (Coeterier, 1996). This strong link between landscape and perception is underlined by the idea that a landscape is such only if it is perceived by people (Council of Europe, 2000). "Perception" orders the sensory input (e.g., seeing), matches it with mental concepts and integrates objects into a whole (Coeterier, 1996; Ware, 2011). Although landscape perception differs between genders, ages and social or cultural background (Stamps, 1999; Strumse, 1996; Zheng, Zhang, & Chen, 2011), there is a high consensus in human perception and preferences for specific landscapes (Bell, 2012; Hägerhäll et al., 2018; Kaplan and Kaplan, 1989; Ulrich, 1986; van den Berg and Koole, 2006). Two main paradigms explain the landscape preferences of people. The first is the paradigm constituted by "cultural theories," which assume that landscape preference is strongly influenced by social and cultural characteristics (Bell, 2012). These theories see familiarity with the landscape or knowledge about the ecological values as key factors for preferences (Gobster, 1999). The second

paradigm is that comprising "evolution-based theories," which explain landscape preferences as reflections of landscape qualities that were favorable for survival in early ages (Kaplan and Kaplan, 1989; Steg, 2013p. 38-46). From both paradigms evolved methods that transform physical landscape structures into psychological dimensions of landscape qualities (Bell, 2012; Coeterier, 1996; 2013: 38-46p. 38-46). Evolutionary theories encompass prospect-Refuge theory (Appleton, 1996), savanna Theory (Orians, 1980), and the preference matrix (Kaplan and Kaplan, 1989). The preference matrix, which is one of the most frequently used theories for assessing landscape qualities, distinguishes psychological dimensions of landscape preferences along two axes. The first encompasses two basic informational needs: "understanding" and "exploring". The former refers to the need of humans to make sense of a scene, whereas the latter depicts the need to gather more information (Kaplan, 1987; Kaplan and Kaplan, 1989). The second axis discriminates between the immediacy with which information is processed (Kaplan, 1987; Steg, 2013), that is, completed immediately or within further mental steps (inferred) (Kaplan, 1987). The two axes span the four visual qualities "complexity, coherence, mystery and legibility" (Kaplan and Kaplan, 1989). Complexity is the immediate exploration and assessment of how many different visual elements are distinguishable and how much is going on in a scene (Kaplan, 1987). Coherence reflects immediate understanding and refers to a sense of order or context that guides a view and thus enables people to understand landscapes (Kaplan, 1987). Uniform textures and shapes increase redundancy and advance the distinction of specific areas within a scene, thereby strengthening one's sense of coherence (Kaplan, 1987). Mystery represents an inferred exploration that captures hidden elements by assessing how much can hypothetically exist if one can further walk into a scene (Stamps, 2004). Exemplarily, the partial coverage of elements with foliage or changes in topography motivate further exploration of a landscape and, thus, increases mystery (Kaplan, 1987; Kaplan and Kaplan, 1989). Finally, legibility pertains to inferred understanding and describes the readability and possibility with which to navigate through a landscape (Stamps, 2004). Landmarks, such as mountains or lakes, influence the legibility of a landscape and enhances its orientation (Kaplan, 1987).

Lothian (2008) and Betakova et al. (2015) investigated the qualities of landscapes with and without RES. The authors concluded that in highquality landscapes, RES strongly diminishes perceived landscape quality compared with low-quality areas. Pasqualetti (2000) specified that the artificiality of RES "rudely" interrupts peoples' understanding of slowgrowing nature and might consequently influence perceived permanence. To permanence, disturbance is an antagonist, whose presence indicates a lack of coherence (Tveit, Ode, & Fry, 2006). Permanence refers to a sense of continuity, which is an important factor for establishing place identity and place attachment (Twigger-Ross and Uzzell, 1996). In relation to wind turbines, Devine-Wright and Howes (2010) asserted that opposition to such projects is related to the disruption of place attachment. Johansson and Laike (2007) found that perceived unity most strongly predicts and explains opposition against wind turbines. Unity reflects 'How well the various components in the environment fit, and function together' (Johansson and Laike, 2007). Thus, unity and coherence are closely related, since both refer to a sense of hanging together (Kaplan, 1987) and the comprehension that "the whole is more than the sum of its part" (Kuiper, 1998; Tveit et al., 2006). In addition, Stanton (2016) conclude that the sense of hanging together and relations between object is important for people to judge scale of wind turbines.

This work treated "*landscape qualities*" as the perceived visual qualities of RES-altered landscapes. As indices of these perceptions, ratings ascribed to coherence, complexity, mystery, and legibility were defined in accordance with the preference matrix (Kaplan & Kaplan 1989). However, relying exclusively on visual landscape qualities in rating might be a critical shortcoming given the questionable reliability claimed by some researchers (Palmer, 2000; Stamps, 2004). Stamps (2004), for instance, conducted a *meta*-analysis and found low reliability and replicability for the preference matrix. Similarly, Palmer (2000)



Fig. 1. Overview of the study goals.

showed that assessing visual landscape qualities using the matrix requires a substantial number of raters to obtain the minimum reliability score of 0.7. <u>Stamps (2004)</u> recommended that measures grounded on mathematical principles (i.e., entropy or visible extent) be used instead of individual landscape quality ratings.

1.2. Relating visual features to landscape qualities

As described above, there is strong evidence that the detection of visual features such as edges, shapes, colors, textures and patterns plays an important role in people's visual perception and, eventually, landscape preference (Bell, 2012p. 179). We emphasize, however, that this is only one-albeit important-entry point in examining landscape experience. Other entry points are place attachment and place meaning. We provide a short overview of quantitative studies that underscored relating visual features to preference ratings (Berman et al., 2014; Dramstad, Tveit, Fjellstad, & Fry, 2006; Hunziker and Kienast, 1999; Ibarra et al., 2017; Kardan et al., 2015; Ode, Fry, Tveit, Messager, & Miller, 2009; Valtchanov and Ellard, 2015). Color tone (hue) and saturation diversity (Kardan et al., 2015), spatial frequency (Valtchanov and Ellard, 2015) and fractality (Forsythe, Nadal, Sheehy, Cela-Conde, & Sawey, 2011; Hagerhall et al., 2008) are significant predictors of landscape preferences. Additionally, the image compression size revealed as an accurate proxy for image content complexity (Tuch, Bargas-Avila, Opwis, & Wilhelm, 2009), it might also function as such for landscape images. A work worth considering is that of Rosenholtz, Li, and Nakano (2007), who developed two indices for visual saliency and redundancy, namely, feature congestion and subband entropy.

Although the authors did not specifically test these with landscape images, both measurements address degree of visual organization and redundancy and may therefore be associated with perceived complexity or coherence. Empirically, Kuper (2017) found a direct correlation between rated complexity and measured entropy with perspective images of landscapes. In contrast to measuring the visual quality of landscapes using perspective images, landscape metrics were developed probe into ecological processes derived from analyses of landscape patterns on land cover or land use maps (Cushman, McGarigal, & Neel, 2008; O'Neill et al., 1988). On the basis of information theory (Shannon, 1948) and fractal geometry (Mandelbrot, 1983), studies related landscape metrics to landscape preferences (Frank, Fürst, Koschke, Witt, & Makeschin, 2013; Fry, Tveit, Ode, & Velarde, 2009; Hunziker and Kienast, 1999; Ode et al., 2009; Palmer, 2004). Specifically, the number of patches on land cover maps, were shown to be significantly correlated with preference estimates of corresponding perspectives (Dramstad et al., 2006; Ode et al., 2009). Further, edge density (Palmer, 2004), Shannon's diversity and the shape index (Frank et al., 2013) are correlated with landscape preferences. Hunziker and Kienast (1999) applied landscape metrics to perspective images in order to analyze gray-tone patches and patterns. Their results showed significant correlations of landscape preferences with Simpson's diversity and evenness as well as the interspersion index. Both interspersion and evenness might serve as proxies for coherence seeing as the increasing homogeneity of landscape patches enhances the sense of coherence (Palmer, 2004).

Visual features such as color, shape, texture, size, and contrast are principal components in visual impact assessments (VIA) of RES. These features have been used intensively to determine the contrasting effects



Fig. 2. Distribution of the landscape types used in this study. Black rectangle indicates the location of the vista in each landscape type.

of RES with surrounding landscapes (e.g., Bishop and Miller, 2007; Brahms and Peters, 2012; Torres Sibille, Cloquell-Ballester, Cloquell-Ballester, & Darton, 2009a; 2009b). Empirical explorations indicated that contrasting effects measured via color differences between wind turbines and backgrounds are strong predictors of perceived visual impact (Bishop and Miller, 2007; Lothian, 2008). The number of wind turbines and their distance to observers are strongly correlated with perceived visual landscape quality (Betakova et al., 2015). A more specific explanation was provided by Johansson and Laike (2007), who disentangled visual perceptions regarding wind turbines and found that contrasting effects directly influence perceived unity and, subsequently, the intention to oppose wind turbine projects.

In the present research, we treated the term "visual features" as pertaining to numerically assessed visual parameters of images containing RES–landscape scenarios. Although many studies investigated the visual effects of RESs, little is known about how visual features (i.e., colors, shapes, and textures) relate to the perceived visual qualities of RESaltered landscapes.

1.3. Goals and hypothesis

Stimulated by the theories and empirical findings described in Subsections 1.1 and 1.2, we defined three goals for our study (Fig. 1).

Goal 1. To understand how RES infrastructure influences landscape quality ratings, that is, ratings involving Kaplan and Kaplan's (1989) dimensions: With previous research (e.g., Johansson and Laike, 2007; Pasqualetti, 2000) as basis, we expected landscape qualities to be affected by RESs. In particular, we expected that in natural landscapes, the sense of permanence, as defined by Pasqualetti (2000), is disrupted by RESs and thus potentially influences the rating of other landscape qualities.

Goal 2: To delve into the relationship between landscape quality ratings and measured visual features for RES-altered landscapes: Motivated by Kuper (2017), who discovered a correlation between stated complexity and designed entropy, we assumed that additional relationships exist between visual features and visual landscape qualities in the context of RES-altered scenes.

Goal 3: To determine how landscape quality ratings and visual features correlate with the overall preference for RES-landscape scenarios: Studies showed associations between measured visual features and landscape preferences (e.g., Hunziker and Kienast, 1999; Ibarra et al., 2017; Kardan et al., 2015) and drew connections between rated visual landscape qualities and preferences scores (Kaplan, 1987; Kuper, 2017; van der Jagt, Craig, Anable, Brewer, & Pearson, 2014). Along the same lines, we hypothesized that both visual landscape qualities and visual features are related to preference for RES-altered landscapes.

2. Methods

To investigate landscape qualities and visual features in the context of RES, we elaborated visual stimuli with different landscape types. Each landscape type was altered with two levels of RES visibility. Thus, all 14 stimuli contain the two visual variables '*landscape types*' (LANDSCAPE) and '*visibility of renewable energy systems*' (RES_VISIBILITY).

2.1. Visual stimuli

Landscape types (LANDSCAPE)

Firstly, we selected typical Swiss landscape types following the widely used biogeographic regionalization of Switzerland (Gonseth, Wohlgemuth, Sansonnens, & Buttler, 2001) (Fig. 2). We removed the landscape types of '*large city centers*' because of planning restrictions for wind turbines in settlement areas. Additionally, we did not integrate landscapes, which contain lakes, due to the fact that water has a strong influence on landscape quality and preference ratings (Ibarra et al., 2017; Tveit et al., 2006). The final seven landscape types show

considerable potentials for the production of renewable energy, though at widely varying environmental costs (Kienast et al., 2017). However, according to the Swiss Energy Strategy 2050 (Swiss Federal Office of Energy, 2012) all of them are - in principle - valid sites to generate the required energy amount to phase out the nuclear power production. The final seven selected landscape types vary considerably in their topography and their land use. They cover the northern areas of flat plateaus primarily used for settlements (PLAT URB) or agricultural production (PLAT_AGRI). Additionally, we considered the Jura ridges (JURA) and the hilly and less densely populated, northern pre-alpine areas (PRE_-ALPS). We then divided the alpine areas with steep terrains into the large inner-alpine valleys, with a relatively high population density (ALP URB), the alpine landscapes used for tourism (ALP TOUR), and the near-natural alpine regions (ALP). During a workshop 25 experts from the fields of landscape planning, wind and photovoltaic project development, as well as employees of the national energy and spatial planning authorities evaluated the seven landscape types. For each landscape type the experts rated several specific views from pedestrian perspective (vistas), according to the potential for representing a future energy landscape, encompassing wind turbines and photovoltaic panels. To visualize the vistas, we used landscape models based on light detection and ranging (LiDAR) data (XYZ). LiDAR in combination with image data represents the landscape characteristics (i.e., color, vegetation, topography) highly detailed. In addition, LiDAR data comes along with three-dimensional geographical coordinates and thus it is possible to site RES accurately in the models. Hence, we combined terrestrial (RIEGL VZ-1000) and airborne LiDAR data to colored 3D visualizations for all seven landscape types. Fig. 2 shows the locations of the vistas (rectangle) and the distribution of the landscape types. The final visualizations show a 160° field of view for each vista.

Visibility of RES (RES_VISIBILITY)

In a second step, the middle 53.3° of each landscape visualization was altered with two scenarios of RES visual impact. Although we used VIA to develop the scenarios, we conservatively call this variable visibility because we neglect other aspects of visual impact (Wolsink, 2018). Both scenarios contain a concurrent visibility of wind and photovoltaic infrastructure. To control for the visibility of wind turbines and photovoltaic panels in each landscape, we made use of two objective visual impact assessments. Namely the 'objective aesthetic impact of solar power plants index' (OAISPP, Appendix A.1: Torres-Sibille, Cloquell-Ballester, Cloquell-Ballester, & Artacho Ramírez, 2009b) and the 'visual impact parameter for wind turbines' (VIWT, Appendix A.2; Brahms and Peters, 2012). The OAISPP combines four measures (i.e., visibility, color, fractality, and contrast, Appendix A.1) for the photovoltaic panels and ranges from 0 (no visual impact) to 1 (strong visual impact). We did not include a climatology coefficient as proposed by Torres-Sibille, Cloquell-Ballester, Cloquell-Ballester, and Artacho Ramírez (2009b) because atmospheric conditions were held constant across all landscapes. The VIWT considers the number of wind turbines, partial visibility, and distance from the observer and ranges from 0 (i.e., no visual impact) to (theoretically) infinity (Appendix A.2). In the high visibility scenarios, we show ten, and in the low visibility scenario three wind turbines. After the initial placement, we iteratively adjusted the exact locations of the turbines in order to reach a minimal VIWT of 1 in all landscape types (Appendix A.3). Subsequently, we placed the three wind turbines in the low visibility scenario in order to reach approximately 70% lower VIWT compared to the high scenario. For the photovoltaic infrastructure, we sited the panels clearly visible in the fore-, and mid-, and background of all landscapes so that OAISPP range between 0.13 and 0.18 in the high and between 0.02 and 0.04 in the low visibility scenarios (Appendix A.4). This procedure allowed control for the variable RES_VISIBILITY in two ways. Firstly, we could clearly distinguish between the LOW and the HIGH level of RES visibility in each landscape. Secondly, we kept the visibility of RES more or less constant between all landscape types within the LOW or the HIGH scenarios of RES visibility. Fig. 3 illustrates the two levels of RES_VISIBILITY for all seven landscape types. We provide the full field of view (160°) in the data repository of this article.

Landscape and Urban Planning 209 (2021) 104000



Fig. 3. Visualization of the middle 53.3° field of view for each landscape type with the two levels of RES VISIBILITY.

Subsequently, we animated the visualizations with rotating wind turbine blades and moving clouds. We kept motion of rotor blades, atmospheric conditions, lightning and seasonality constant between all stimuli, since these parameters influence the visibility of RES (Apostol, Palmer, Pasqualetti, Smardon, & Sullivan, 2016p. 180). Finally, we rendered 30-second panoramic videos from pedestrian perspective with a resolution of 5760 by 1080 pixels (see additional data).



Fig. 4. Example of an experimental proceeding in the MVAL for one participant, presenting randomly six stimuli from the total amount of 14 stimuli.

2.2. Procedure

We collected the data for each participant individually in a laboratory lasting for app. 45 min. All participants reported being native German speakers, physically and psychologically healthy, with normal or corrected-to-normal vision and hearing. The study was approved by the ethics committee of ETH Zurich. Participants were compensated 30 Swiss franc after finishing the session. To keep experimental conditions (i.e. light, sound) constant, we used a Mobile Visual Acoustic Laboratory (MVAL; Manyoky et al., 2016). Within the MVAL we projected the stimuli on three 106° angled screens (7.1 m \times 0.6 m). The entire experiment was performed with the software 'Experiments in Virtual Environments' (EVE; Grübel et al., 2017). This software allows the experimenter to interactively present questions to participants, while showing still images, videos or virtual realities. Particularly, we used this software to record physiological responses on landscape changes caused by RES (reported in Spielhofer et al., 2021). After signing the declaration of consent, the participants were familiarized with laboratory conditions and the keyboard control with a training trial. The training trial consisted of two 30-second scenes (videos of a moving blue circle and a moving red cube), presented to participants sequentially. Following the training stimuli pair, we asked participants to use the arrow keys on the keyboard in order to select their preferred stimulus. The main experiment was divided into two parts (Fig. 3). In the first part, participants completed three testing trials. Each trial consisted of two subsequently presented landscape-RES scenarios, followed by the participants choice of their preferred scenario. In addition to the preference, we measured participants' physiological arousal response, which is reported in XYZ. The three testing trials were separated by 20-second intervals consisting of a black cross on a gray background. We paired the stimuli so that one trial consisted of two 'HIGH' RES visibility videos, one trial consisted of a 'HIGH' and a 'LOW' RES visibility video, and one trial consisted of two 'LOW' visibility videos. The order of these trial types and the landscapes that composed each trial type were randomized and counterbalanced across participants, except that a participant never saw the same landscape more than once. In the second part of the session (Fig. 4), we showed the same six stimuli from part one again in a new randomized order. Participants rated the landscape qualities for each of the six stimuli based on the landscape preference matrix (Kaplan and Kaplan, 1989). We closed the session with a post questionnaire to record participants' socio demographics.

2.3. Measures

In this section, we present the questions to assess people's preference and the rated landscape qualities for the landscape-RES scenarios. Further, we show the visual features, measured on each stimulus.

Preferences

To assess people's landscape preference some studies used pairwise comparisons of stimuli (e.g., Courcoux and Semenou, 1997) other studies used rating (e.g., Schirpke, Tappeiner, Tasser, & Tappeiner, 2019). Particularly, for subjective ratings, the pairwise comparison method has been shown as useful due to the simplicity of the assessment (Courcoux and Semenou, 1997) and no substantial differences to ordinal rating scales revealed (Hunziker and Kienast, 1999; Stamps, 2004). Thus, we used pairwise comparisons in this study, to assess landscape preferences. In part 1 (Fig. 3) of the experiment, each participant saw three times two scenarios sequentially and choose their preferred scenario based on the question 'Which landscape do you like better' (original germ.: 'Welche Landschaft gefällt Ihnen besser?').

Rated landscape qualities

To rate the landscape qualities, we used the four dimensions complexity, coherence, mystery and legibility from the preference matrix (Kaplan and Kaplan, 1989). We assessed the four landscape qualities with a set of nine questions and report the original German question and

Table 1

Models to calculate effect of fixed effects on dependent variables landscape qualities or CHOICE with Likelihood-Ratio tests.

	Dependent variable	Fixed factors	Random factors
Linear mixed	l-effects model (ln	ner)	
three- fixed- factor model	Landscape qualities	LANDSCAPE + VISIBILITY+ (LANDSCAPE * VISIBILITY)	SUBJ_ID
two-fixed- factor model	Landscape qualities	LANDSCAPE + VISIBILITY	SUBJ_ID
one-fixed- factor	Landscape qualities	VISIBILITY OR LANDSCAPE OR GENDER OR AGE OR HOME LANDSCAPE	SUBJ_ID
Null model	Landscape qualities	1	SUBJ_ID
Generalized	linear model with	binomial regression (glm)	
three- fixed- factor model	CHOICE	LANDSCAPE + VISIBILITY+ (LANDSCAPE * VISIBILITY)	No random effects included
two-fixed- factor model	CHOICE	LANDSCAPE + VISIBILITY	
one-fixed- factor	CHOICE	VISIBILITY OR LANDSCAPE OR GENDER OR AGE OR HOME_LANDSCAPE	
Null model	CHOICE	1	

corresponding English translation in Appendix B.1. Thus, within part two of the experiment (Fig. 3), each participant answered all nine questions for each of the six scenarios with a 5-point Likert scale. The questions, scale and the wording were adopted from (Kienast et al., 2015). With slightly different wording but similar terms these questions were used in other studies (Singh, Todd Donavan, Mishra, & Little, 2008; e.g., van der Jagt et al., 2014).

Visual features

Based on the middle view (1920 by 1080 pixels), of the first video frame we computed 20 visual features for each of the 14 stimuli. To select the visual features, we refer to measures which have been found as relevant in the context of landscape perception and preferences (Appendix B.2). Firstly, we applied pixel- based statistics, where information from each single pixel is extracted and summed up to the total amount of pixels of the image. These statistics mainly cover color aspects (hue, sat, bright), visual information density (fc, se, bytes) and spatial frequency (sf_low, sf_high). Secondly, we used landscape metrics with color classified images as input. We classified the images with 12 centers by the use of a k-nearest neighbor classifier (Hunziker & Kienast, 1999). Based on the 12 class RGB reduced images (Appendix B.3), all landscape metrics were calculated in R-Studio (V.3.5.5) on landscape level with the R-package 'landscape metrics' (Hesselbarth, Sciaini, With, Wiegand, & Nowosad, 2019; McGarigal, Cushman, & Ene, 2012). Thirdly, we calculated the number of pixels and edges which are exclusively related to the RES.

2.4. Data processing and analysis

With R studio version 3.5.1 (R Core Team, 2018) we conducted descriptive and inferential statistics. As a first step, we tested the inter-item reliability for the two items of coherence, complexity, mystery and the three items of legibility (Appendix B.1). We consider a Cronbach alpha of 0.7 as an acceptable reliability (Palmer, 2000) and averaged the corresponding items with the median for each participant. We then checked the data for normal distribution (Shapiro test) and examined homogeneity of variances (Levene test). Secondly, we assessed the effect of the two independent variables RES_VISIBILITY and LANDSCAPE on each of the 20 visual features with split-plot ANOVAS. Split-plot ANOVAS were used because each LANDSCAPE contains two dependent levels of RES_VISIBILITY. In a third

step, we examined the effects of RES VISIBILITY and LANDSCAPE on the landscape quality ratings and the binary choice data. In order to select an appropriate model for the landscape quality ratings and the choice, we compared general linear models (glm) with mixed effect models (lmer) with Likelihood-ratio tests (Giampaoli and Singer, 2009). For the landscape quality ratings, Likelihood-ratio tests revealed lower AIC for mixed effect models with integrated random effects compared to general linear models. Thus, we calculated the lmer with the R-package 'Ime4' (Bates, Mächler, Bolker, & Walker, 2015). We considered the rated landscape qualities as dependent and the LANDSCAPE and RES_VISIBILITY as independent variables (Table 1). Mixed models have the advantage that they can account for variation in the data that is related to further aspects and not explicitly expected in the hypotheses (i.e., random factors). We used the subjects (SUBJ ID) as random factors to account for the fact that the ratings of landscape qualities are not fully independent since we have a repeated measurement design (each subject rated six times the landscape qualities). Further, mixed models are more suitable for imbalanced study designs (each participant assessed only 6 out of the total 14 scenarios). In addition to the main effects (RES VISIBILITY & LANDSCAPE), we tested the effect of socio demographics on the landscape qualities (Stamps, 1999; Strumse, 1996; Zheng et al., 2011). Specifically, we examined the effect of GENDER, AGE and the fact that a stimulus might represent the participants' home landscape (HOME_LT; Hunziker et al., 2008). In order to test the effect of age, we split our data into a group (AGE_GROUP) with people younger than 30 years and a group with people older than 29 years. For the preference date, Likelihood-ratio tests revealed lower AIC for generalized linear models (glm) with binomial logit regression compared to mixed effect models. The inclusion of random effect would lead to model overfitting. In consequence, we applied a glm to determine the effect of LANDSCAPE and RES VISI-BILITY on the binary choice data (Table 1). In order to calculate p and X^2 values for the main effects within the lmer and the glm models, we used additional Likelihood-ratio tests. We compared the null-model with the onefixed factor model to assess the effects of each individual fixed factor and the two-fixed factor model with the three-fixed factor model to determine the interaction effects of LANDSCAPE and RES VISIBILITY (Table 1). We then calculated the effect size (R² & f²) with the R-package 'Sjstats' (Lüdecke, 2020). For mixed models, we used marginal R^2 which represents the explained variance from the fixed, without the random effects (Nakagawa and Schielzeth, 2013). According to Cohen (1992), we consider $f^2 = 0.02$ as small effects, $f^2 = 0.15$ as medium effects and $f^2 > 0.3$ as strong effects. For the logit models of with the binary choice as dependent variable we calculated the coefficient of determination (D; Tjur, 2009).

With the median we aggregated the landscape quality ratings to the average quality rating for each scenario (N = 14). Similarly, we aggregated the binary choices to a preference score (PREF_SCORE) for each scenario by dividing the total number the scenario has been shown by the number the scenario has been preferred. Finally, we applied Kendall and spearman correlations with a Bonferroni corrected alpha level for multiple testing to analyze the relationship between the average landscape qualities, the visual features and PREF_SCORE. Non-parametric correlations were used because of a relatively small sample size (N = 14 scenarios) and not normally distributed data.

3. Results

A total of 135 participants (61 women and 72 men, mean age = 28 years, SD = 11.74, age range = 19–73 years) completed the experiment. Two participants were excluded because of software issues. The young AGE_GROUP contains 98 participants (47 woman, 51 men, mean age = 22.4 years, SD = 2.28, range = 19–29 years), the older AGE_GROUP 28 (11 woman, 17 men, mean age = 46.6 years, SD = 12.33, range = 32–73 years). Nine participants could not be included into an age group since they did not state their age.

The inter item reliability for each landscape quality was calculated for several data subsets and for the whole dataset (Table 2, Appendix C). Independent of the data subset, the two items of coherence and complexity as well as the three legibility items, revealed acceptable (>0.7)

Table 2

Inter-item reliability of the landscape qualities for the overall data set, the subsets	of landscape types (LANDSCAPE) and the subset of RES_VISIBILITY.
---	--

Landscape qualities	LANDSCAPE		RES_VISIBILITY		Overall					
	PLAT_URB	PLAT_AGRI	JURA	PRE_ALPS	ALP_URB	ALP_TOUR	ALP	LOW	HIGH	
Coherence	0.81	0.77	0.86	0.83	0.86	0.82	0.86	0.82	0.84	0.84
Complexity	0.71	0.78	0.82	0.75	0.71	0.7	0.85	0.78	0.79	0.78
Legibility	0.76	0.74	0.68	0.78	0.77	0.68	0.6	0.73	0.75	0.74
Mystery	0.54	0.73	0.7	0.67	0.46	0.72	0.72	0.66	0.58	0.62
N data subset	112	120	118	104	119	110	116	400	399	799

overall reliability. In consequence, we averaged the items with the median for each landscape quality. In contrast, the two items of mystery revealed too low overall reliability, hence the two items (MYST_A, MYST_B) will be treated as separate variables in the further analysis.

3.1. Rated qualities of RES altered landscapes

Since we have an unbalanced experimental design, the number of landscape quality ratings for each scenario varies between 48 and 70 (Appendix D.1). We calculated the median and Inter-Quartile Range (IQR) of the landscape quality ratings (Fig. 5) and reference to Appendix D.1 for exact values.

Main effects of RES_VISIBILITY and LANDSCAPE on landscape qualities Despite complexity, all landscape qualities revealed a significant effect on RES_VISIBILITY. However, coherence ratings show the strongest effect on RES_VISIBILITY ($f^2 = 0.2$, Table 3). Additionally, all landscape qualities varied significantly between the levels of LAND-SCAPE. A strong effect ($f^2 < 0.3$) of LANDSCAPE could be observed for Mystery_B, Legibility and Complexity (Table 3). Mystery_B revealed the only landscape quality with a significant and strong interaction effect of LANDSCAPE and RES_VISIBILITY. Contrarily, the effect of different LANDSCAPEs is stronger for complexity, legibility and Mystery_A.

Coherence is the only landscape quality which shows a stronger effect on the levels of RES_VISIBILITY compared to the levels of LAND-SCAPE. Therefore, a post hoc analysis between the levels of RES_VISIBILITY has been performed for each landscape type. In the near natural alpine landscape (ALP, df = 763, T = -3.591, p = .024, d = -0.6), the touristic alpine area (ALP_TOUR, df = 763, T = -3.502, p = .032, d = -0.58) and the urban alpine valley (ALP_URB, df = 763, T = -4.588, p = .0004, d = -0.78) post hoc analysis show significantly higher rated coherence in the low-, compared to the high visibility. In the urban and agricultural plateaus (PLAT_URB & PLAT_AGRI), the hilly landscapes of the JURA and the pre alps (PRE_ALPS) the degree of visibility of RES does not affect rated coherence significantly (Fig. 6).

Effects of socio demographics on landscape qualities

The effects of GENDER, AGE and HOME_LT on the ratings of the landscape qualities have been tested with mixed linear models and Likelihood-Ratio tests (Table 1). The test revealed a significant, medium effect of AGE_GROUP on coherence ($X^2(1) = 7.302$, p = .007, R² = 0.144, f² = 0.17). Additionally, a strong effect of GENDER and MYS-TERY_A evolved (X^2 (6) = 3.922, p = .048, R² = 0.243, f² = 0.32) (Appendix E). The effect that high visibility is rated with lower coherence is stronger for young people compared to older people (Table 4).

3.2. Relation between measured visual features and rated landscape qualities

We report the descriptive statistics of the 20 visual features in Appendix D.2.

Correlation between visual features and landscape qualities

To test the correlation between 20 visual features and five landscape qualities, we correct the alpha level to p = .002, in order to correct for possible alpha inflation. To find relations between rated visual landscape qualities and measured visual features for N = 14 scenarios, we applied spearman's rank and Kendall correlations (Fig. 7; see Appendix F, for exact numerical r and p values).

Only a few visual features correlate significantly (p < .002) with the landscape qualities. The contiguity index with rated complexity (*tau* = -0.582, *tau* p = .003, r = -0.758, p = .0016) and the amount of RES related edges with rated coherence (*tau* = -0.582, *tau* p = .003, r = -0.75, p = .002). Additionally, we report a strong correlation between the high spatial frequency (sf_high) and the individually treated item Mystery_B (*tau* = 0.736, *tau* p = 7.7e-5, r = 0.88, p = 2.5e-5).

Main effects of RES_VISIBILITY and LANDSCAPE on the visual features Split-plot ANOVAS showed that most of the visual features vary significantly between the landscape types (LANDSCAPE) but not between the LOW and the HIGH RES visibility. Nevertheless, subband entropy (*se*, *F*(1) = 6.069, p = .05), feature congestion (*fc*, *F*(1) = 28.78, p = .002), file size of the stimuli (*bytes*, *F*(1) = 16,15, p = .007) and color hue (*F*(1) = 39.6, p = .0008) show an additional significant effect on the levels of VISIBILITY (Appendix G). The two visual features, which relate exclusively on RES (PIX_RES, EDGE_RES), vary much stronger between the levels of RES VIS-IBILITY compared to the levels of LANDSCAPE (Appendix G). This result is a proof that the visual balancing between the landscape types was successful and the combined visibility of wind and photovoltaic infrastructures is comparable between the different landscape types.

3.3. Preference scores for RES altered landscapes

Fig. 8 indicates that people generally prefer the LOW compared to the HIGH visibility scenario. However, in the flat areas with mainly settlements the HIGH scenario is more preferred compared to the LOW. The highest preference scores, independent from the amount of RES can be observed in the hilly and less densely populated areas of the JURA and the pre alps (PRE_ALPS).Fig. A3.Fig. A4.

For the preference data, the comparison between the three-fixed-factor model and the two-fixed-factor model (Table 1) revealed a significant interaction between LANDSCAPE and VISIBILITY (X^2 (1) = 17.134, p = .008, D = 0.09). In addition, the comparisons between the one-fixed-factor models and the null model revealed significant effects for LANDSCAPE (X^2 (6) = 51.843, p < 2.6e-9, D = 0.06) and RES_VI-SIBILITY (X^2 (1) = 7.431, p = .006, D = 0.009) on participants' preferences. These effects suggest that participants preferred low visibility of RES (compared to high visibility) and that this effect varied across landscapes. Notably, this trend can be observed for each landscape except for PLAT_URB and PLAT_AGRI (Fig. 8).

Relation between visual features and preference scores

With a Pearson correlation we checked the relation between the 20 visual features and the preference score. Again, we consider an alpha level of 0.002 as a significance threshold. However, no correlation between the visual features and the preference score revealed significant (Appendix H.1)

Relation between landscape qualities and preference score

Finally, we applied Kendall and spearman rank correlation in order to find relations between the five aggregated ratings of landscape qualities and the preference score. We correct the alpha level to 0.01 since we tested only five possible correlations. Coherence (tau = 0.538, tau p = .006, p = .012, r = 0.13), legibility (tau = 0.494, tau p = .013, p =.003, r = 0.72) and mystery_B ratings (tau = 0.648, tau p = 7e-4, p = 7e-4, r = 0.79), correlated significantly with preference scores. Mystery_A (p =.48, r = 0.2) and complexity (p = .93, r = -0.02) did not show any significant relation with the preference score (Appendix H.2).



Fig. 5. Median and (IQR) of the landscape quality ratings.

4. Discussion

This section summarizes the results, discusses their implication for each goal separately and ends with a description of the study's limitations. Goal 1: Rated qualities of RES altered landscapes

First, we investigated the influence of RES infrastructure on people's landscape quality ratings. Rated coherence varies stronger between the two levels of RES visibility compared to mystery, legibility and

Table 3

Effects of VISIBILITY	, LANDSCAPE and	Interactions on	landscape of	jualities (N = 798).
-----------------------	-----------------	-----------------	--------------	-------------	-----------

Landscape qualities	Fixed effe	Fixed effects												
	RES VISIBILITY				LANDSCAL	LANDSCAPE				INTERACTION				
	$\chi^{2}(1)$	Marginal R ²	Cohens f ²	р	$\chi^2(1)$	Marginal R ²	Cohens f^2	р	$\chi^{2}(1)$	Marginal R ²	Cohens f ²	р		
Coherence	71.804	0.17	0.2	2.2e-16	58.478	. 153	0.18	9.16e-11	11.16	0.241	0.32	0.08		
Complexity	0.018	0.064	0.07	0.667	137.78	0.239	0.31	0	8.35	0.25	0.33	0.21		
Legibility	14.417	0.124	0.14	0.0001	120.28	0.257	0.35	0	8.8	0.272	0.37	0.18		
Mystery_A	6.746	0.049	0.05	0.009	93.499	0.162	0.19	0	7.55	0.179	0.22	0.27		
Mystery_B	15.877	0.027	0.03	6.76e-5	215.81	0.277	0.38	0	12.88	0.304	0.45	0.045		



Fig. 6. Rated coherence for each landscape-RES scenario.

Table 4	
Effects of RES_VISIBILITY, LANDSCAPE and Interactions on rated coherence, separated by AGE_G	ROUP.

		RES_VISIBILITY			LANDSCAPE				INTERACTION				
AGE GROUP	Ν	X ²	р	R ²	f^2	X ²	р	R ²	f^2	X ²	р	R^2	f^2
< 30	98	62.509	2.60E-15	0.1	0.11	45.64	1.40E-08	0.07		5.9514	0.428		
>30	28	7.412	0.006	0.03	0.038	12.662	0.048	0.061	0.065	7.6657	0.26		

complexity. Kaplan and Kaplan (1989) asserted that coherence is processed immediately. Consequently, or first result implies that people's perception of RES altered landscapes contains a substantial affective component. Therefore, we support the finding of Maehr, Watts, Hanratty, and Talmi (2015). The authors discovered that measured physiological arousal, which expresses an immediate affective reaction, is higher for landscapes containing wind turbines than other infrastructure. Further, we found that people generally rated a high RES visibility as less coherent over a low impact. We conclude that an increasing number of RES decreases the sense of coherence. This finding contrasts with the statement of Tveit et al. (2006) that enhancement of the visual redundancy should increase the sense of coherence in a scene. Although, RES add visual repetitions and uniformity to landscapes (Apostol et al., 2016), the sense of coherence might increase with only a low RES visibility up to a certain point. After reaching this point, which is specific for different landscape types, coherence decreases with more RES visibility. Further research should test the relationship between amount of RES and coherence more in detail. Aside from finding a generally lower rated



Fig. 7. Correlation matrix with averaged visual landscape qualities and visual features for N = 14 scenarios.

coherence of high RES visibility, we discovered this effect to be strongest in urbanized alpine areas, touristic alpine regions, and near natural alpine landscapes. Contrarily, in the densely populated flat plateau (PLAT URB), coherence is rated almost equally in high- and the low RES visibility scenarios. This finding indicates two things. Firstly, the visual coherence of the PLAT URB landscape is not affected by the amount of RES. Thus, from a visual perspective, this landscape seems to have potential for a stronger visibility of RES. Secondly, the degree of naturalness does not serve as a unique predictor for the coherence difference between low and high RES visibility scenarios. Most likely, the coherence differences between high- and low RES visibility is influenced by further landscape connotations. Therefore, we see coherence ratings as a potential indicator for the 'place-technology-fit' (Devine-Wright and Howes, 2010). However, further research needs to build up on this statement and test whether coherence mainly describes a visual part (e. g., scale; Stanton, 2016) or also a functional part of place-technology-fit.

In contrast to coherence, complexity varies strongly among the seven landscape types but not between the RES visibility scenarios. Complexity is an immediate exploration involving an assessment of how many visual elements are present in a landscape (Fry et al., 2009; Kaplan and Kaplan, 1989; Stamps, 2004). On the grounds of the landscape types selected for our experimental stimuli, we implicitly represented this visual richness to some extent. Our results show that landscapes containing many man-made elements and diverse topographies are rated as more complex, whereas uniform flat areas with few man-made elements are less complex. Mystery and legibility are not related exclusively to RES visibility or the landscape types. Both dimensions need further processing (Kaplan and Kaplan, 1989) and might consequently entail more cognitive considerations, which possibly involve scrutinizing a landscape, including an RES, as a whole.

Goal 2: Relationship between landscape qualities and visual features

Motivated by the critiques of Stamps (2004) regarding the reliability of the preference matrix, we inquired into the association between measured visual features derived from the stimuli and the rated visual qualities of RES altered landscapes. The analysis revealed that rated complexity is negatively correlated with measured contiguity. Specifically, the analysis shows that diversity in land use and topography leads to lower contiguity and high complexity values. This finding is in line with La Fuente Val et al. (2006) who demonstrated that contiguity is inversely related to patch richness and positively correlated with the Shannon evenness index. However, compared to Kuper (2017) and Stamps (2004), we could not find a significant correlation between the measured entropy and the rated



Fig. 8. The number of choices and no choices and the preference score for each of the 14 scenarios.

complexity of the scenes. This could be due to the slightly different formulation of the questions used to assess complexity in our Germanlanguage study. On the other hand, one has to consider that we did not calculate the entropy according to Shannon (1948), but the subband entropy according to Rosenholtz et al. (2007).

Rated coherence revealed as significantly correlated with counted edges, exclusively related to wind and solar infrastructure. The more RES edges are visible within a scenario, the less coherent the scenario, consistent with Kuper's (2017) work. Within a plant context, the author showed that a high number of planted regions in a scene drives down ratings of coherence. Although the measured feature congestion was not significantly correlated with rated coherence, we propose to further investigate this relationship. Feature congestion varied significantly between the RES visibility scenarios, indicates a degree of visual organization and redundancy, and might thus influence coherence (Bell, 2012; Tveit et al., 2006). However, most of the visual features commonly used to operationalize landscape qualities potentially fail to measure RES-prompted landscape changes.

Goal 3: Preference for RES-altered landscapes

We found that visual preference is significantly influenced by the landscape types and by the RES visibility level. However, the two variables landscape types and RES visibility could only explain a very small part of the preference. Although we specifically asked about people's visual preference for energy landscapes, this result implies that people think beyond visual aspects when evaluating such scenes. Overall, people prefer low RES visibility scenarios over high-visibility counterparts, in congruence with the findings of Betakova et al. (2015), who reported lower visual preferences for higher number of wind turbines. Independent of the number of RES, the hilly and sparsely populated pre alpine area is the most strongly favored landscape. By contrast, the inner alpine valley and the urban plateau are the least preferred. Interestingly, the urbanized plateau (PLAT URB), which is generally rated as of low visual quality, is the only landscape in which a high-visual-impact scenario is preferred over a low-impact equivalent. Thus, our results support those of Lothian (2008) and Betakova et al. (2015), who found that installing wind farms in high-visual-quality landscapes decreases visual quality, whereas adding wind turbines in low-visual-quality landscapes may increase visual quality.

We found no direct correlation between measured visual features and preferences, thus substantiating the finding of Fuente Val et al. (La Fuente Val et al., 2006). The study revealed only limited correlation of visual indices with scenic beauty estimations and conclude that might be attributed to the broad and very normative concept of landscape preference.

Rated complexity and the similar item of mystery (MYST_A: 'There is

much to discover in this landscape') are unrelated to preference. Contrarily, coherence, legibility and one item related mystery ('I would like to get to know this landscape better ') are correlated with visual preferences for RES altered landscapes. Hence, "making sense," "understanding," and "readability" are more important factors for preferences of RES altered landscapes compared to diversity aspects. Similarly, Johansson & Laike (2007) demonstrated that "how well landscape components fit and function together", are important aspects for the opposition against wind turbines.

Limitations

We worked under the premise that for successful energy transition, the right question is not "*if*" RES should be established but "*where*" and "*how many*". Correspondingly, we showed only landscape stimuli containing RES. This decision prevented us from identifying a connection between a landscape change from RES presence to no RES presence. Our study design nonetheless cleared the way for illuminating the effects on people's quality ratings and preferences with regard to different landscape types.

In addition, we showed only combined visual impact of wind and photovoltaic infrastructure. Future studies should show landscape stimuli separated by energy facilities in order to investigate differences in landscape quality ratings between wind turbines and solar panels. Although other researchers found connections between measured visual features and landscape preferences (Ibarra et al., 2017; Hunziker & Kienast, 1999; Kardan et al., 2015), we detected no direct relationship between visual features and rated preferences. This may be explained by the setting of our experiment, which involved only seven landscape types and therefore covered inadequate variance within single visual features. Further studies directed toward ascertaining this association may need a wider variety of landscape types and, thus, more visual stimuli. However, both mentioned issues lead to significantly more stimuli, which increases the risk that evaluations are influenced by fatigue effects.

Another shortcoming is that our study was conducted in German. The slightly different wording and formulation of the language could explain the differences in our results and those of research carried out in English. This issue might be of special interest in further *meta*-analysis, wherein the correct meanings of German and English terms are carefully considered.

Finally, we used the preference matrix (Kaplan and Kaplan, 1989) as only one of several methods for assessing people's perception of landscape qualities. Because the preference matrix distinguishes between immediate and inferred ratings of landscape quality, we used this assessment to relate landscape qualities to measured immediate, physiological arousal responses (Spielhofer et al., 2021). Combining physiological data and landscape quality ratings leads to a better

understanding of the individual response and assessment of landscape changes caused by renewable energy infrastructure.

5. Conclusion

The visual aesthetic effects of RES on landscapes are important factors for resistance against energy infrastructure and therefore hamper the cultivation of sustainable societies. Our results revealed that landscapes with the potential to facilitate the establishment of connections with energy facilities might be visually preferred by people. Possible connectors can be visual (repetitive shapes, colors, and textures) but also contextual/functional in nature, such as other man-made elements or similar land uses. Further study is needed to specifically look into these visual and contextual connectors. We cannot directly formulate recommendations for landscape planners and policy makers, but value for these stakeholders may come in the form of this work's unearthing of the potential to incorporate visual qualities into VIAs. Further our results can contribute to determine landscape specific visual thresholds, which incorporate visual qualities. Capitalizing on visual qualities might provide additional insights into what people think about landscape changes through RES instead of merely focusing on RES visibility. Finding suitable areas that incorporate people's visual perception can help mitigate resistance against RES. Rated coherence can serve as a starting point given its suggested effectiveness in the assessment of RES-induced visual changes in landscapes. Future research should investigate coherence, its connection to place-technology fit, and possible objective visual features in the context of RES landscapes. Specifically, such scholarship should account for the different connotations of various landscape types and how these are influenced by renewable energy systems.

CRediT authorship contribution statement

Reto Spielhofer: Conceptualization, Data curation, Formal analysis, Investigation, Software, Visualization, Writing - original draft. Marcel Hunziker: Conceptualization, Methodology. Felix Kienast: Methodology. Ulrike Wissen Hayek: Investigation, Methodology, Visualization. Adrienne Grêt-Regamey: Funding acquisition, Supervision, Writing original draft.

Acknowledgement

This research was funded by the Swiss National Science Foundation (SNSF), National Research Programme NRP 70 "Energy Turnaround", grant number 407040 173808/1 (ENERGYSCAPE). Further support was provided by the Swiss Federal Office of the Environment (FOEN), the Swiss Federal Office of Energy (SFOE), the Elektrizitätswerke des Kantons Zürich (EKZ), the Swissgrid AG, the Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), and the Sophie and Karl Binding Stiftung. The authors bear the entire responsibility for the content and for the conclusions that were drawn. Additionally, we thank Flurina Wartmann and Boris Salak for their helpful advice regarding the questionnaire and data analysis. Finally, we thank all of the participants for their time and effort during the experimental sessions.

Appendix A. Visual impact of RES

Scenario development with calculated visual impact values for photovoltaic panels (OAISPP) and wind turbines (VIWT). Parameters OAISPP

Table A1. Mean and standard deviation for calculated visual impact parameters of photovoltaic panels (OAISPP), in the LOW and the HIGH Renewable Energy Systems (RES) SCENARIO levels (Torres-Sibille et al., 2009)

	Feature	Visibility	Color	Fractality	Concurrence	Objective Aesthetic Impact of Solar Power Plants	Number of pixel covered with PV	Number of edge pixel relating to PV
	RES visual impact dimension	Area (pixel) covered with PV related to background pixel	Color contrast of panel vs. close surrounding	Artificiality of installations against natural background	More than one type of solar panel within a plant produce concurrence and influence on perception	Weighted overall visual impact of PV system		
	Scale	[0 -1]No PV – total view covered with PV	[0 – 1]No color contrast – high contrast between PV ad background	[0 – 2]Background has higher fractality as PV [If<1]PV has higher fractality than background [If>1]	[0 -1]Highest concurrence if two panel types are equally distributed	0 = No visual impact1 = highest visual impact		
	Calculation	Calculation steps J	proposed by Torres	-Sibille et. al., (2009) ir	nplemented in MATLAI	3 script (appendix)	Pixel subtraction (Scenario – No- RES scenario)	MATLAB Canny edge detection
LOW RES SCENARIO	mean	0.1	0.014	0.044	0.3	0.095	16264	3822
	Standard deviation	0.02	0.007	0.02	0.1	0.02	4606	1219
HIGH RES	mean	0.15	0.02	0.04	0.64	0.16	26738	7379
SCENARIO	Standard deviation	0.04	0.01	0.02	0.16	0.01	7432	4658

Parameters VIWT

Table A2. Mean and standard deviation for calculated visual impact parameters of wind turbines (VIWT) in the LOW and the HIGH Renewable Energy System (RES) SCENARIO levels (Brahms and Peters, 2012).

	Feature	Absoulut number of WT	Effect of additional WT	Pratial visibility	Distance effect	Total visual impact wind	Number of pixels covered with wind turbine	Number of edge pixels relating to wind turbines
	RES visual impact dimension	Total number of WT placed in scene	Reduction effect of additional WT	Effect of partial visible WT in landscapes	Effect of distance to closest WT	Total calculated visual impact of wind turbines		
	Scale		$\begin{array}{l} 1 = \text{only single} \\ \text{WT}{\sim}0.18 = \text{add} \\ 30 + \text{WT} \end{array}$	$\sum_{i=1}^{NUM_{-WT}} part_{vis_i}(0)$ = all WT total invisible or no WT)	$1 = 0 \text{ m} 0.03 \sim 10'000 \text{ m}$	=RED_NUM_WT * PART_VIS_WT*EFF_DIST_WT		
	Calculation		Calculation steps proposed by (Brahms und Peters., (2012) implemented in MATLAB script				Pixel subtraction (Scenario – No- RES scenario)	MATLAB Canny edge detection
LOW RES	mean	3	0.33	2.73	0.32	0.29	3341	1316
SCENARIO	Standard deviation	0	0	0.24	0.04	0.04	1845	1182
HIGH RES	mean	10	0.24	8.89	0.59	1.27	5085	3244
SCENARIO	Standard deviation	0	0	0.78	0.1	0.25	1982	920



Fig. A3



Fig. A3. VIWT for LOW and HIGH scenarios of all landscape types.





Fig. A4. OAISPP for LOW and HIGH scenarios of all landscape types.

Appendix B. Measures

Table B1. Nine items to assess the landscape qualities based on the preference matrix (Kaplan & Kaplan, 1989). Originally the questions were asked in German. All statements have been answered on a 5-point Likert scale ranging from 1 = totally disagree (Germ. "Trifft gar nicht zu") – 5 = totally agree (Germ. "Trifft sehr zu").

Item	Question	Landscape quality
COMPL_A	"There are many different things in this landscape."	Complexity
	German: "In dieser Landschaft gibt es viel Verschiedenes."	
COMPL_B	"This landscape is diverse."	Complexity
	German: "Diese Landschaft ist abwechslungsreich."	
COHER_A	"This landscape is coherent in itself."	Coherence
	German: "Diese Landschaft ist in sich stimmig."	
COHER_B	"The individual things or components of this landscape fit together."	Coherence
	German: "Die einzelnen Dinge oder Bestandteile dieser Landschaft passen gegenseitig zusammen."	
LEGI_A	"This landscape is clear."	Legibility
	German: "Diese Landschaft ist übersichtlich."	
LEGI_B	"There are landmarks in this landscape that help you find your way."	Legibility
	German: "In dieser Landschaft gibt es Orientierungspunkte, die einem helfen, sich zurechtzufinden."	
LEGI_C	"I find my way around this landscape very well."	Legibility
	German: "Ich finde mich in dieser Landschaft sehr gut zurecht."	
MYST_A	"There is much to discover in this landscape."	Mystery
	German: "In dieser Landschaft gibt es viel zu entdecken."	
MYST_B	"I would like to get to know this landscape better."	Mystery
	German: "Ich möchte diese Landschaft gerne besser kennen lernen."	

Table B2. Calculated visual features and their related concepts for visual perception of landscapes

Concept	Relevance for visual perception	Index (coding)	(Source) / calculation
Pixel based measures			
Color & contrast	Color tone	Hue (hue)	(Berman et al., 2014; Kardan et al., 2015) MATLAB image
	Color intensity	Saturation (sat)	processing toolbox
	Light and shadow	Brightness (bright)	
Visual information	Visual saliency / redundancy	Feature congestion	(Rosenholtz et al., 2007) Free available MATLAB script
density		(fc)	
		Subband entropy (se)	
	Visual complexity	JPEG file size (bytes)	(Tuch et al., 2009)File size on storage
Spatial frequency	Sharpness	Low spatial	(Valtchanov and Ellard, 2015) Gaussian filter and Fourier
		frequency (sf_low)	transformation. Python script provided by D. Valtchanov
		High spatial	
		frequency (sf_high)	
Landscape metrics			
Area & edges	Detection of edges \rightarrow perceived naturalness	Edge density (ed)	(Kardan et al., 2015) MATLAB image processing toolbox
Shape	Complexity or naturalness of shapes \rightarrow landscape preference	Shape index (shape)	(Ode et al., 2009)
		Fractality (frac)	(Forsythe et al., 2011; Hagerhall et al., 2008)
		Contiguity (contig)	(La Fuente Val, Atauri, & de Lucio , 2006) \rightarrow antagonist of
			evenness

(continued on next page)

R. Spielhofer et al.

(continued)

Concept	Relevance for visual perception	Index (coding)	(Source) / calculation
Core area	Area shape interaction	Core area (cai)	
Aggregation	Dispersion / interspersionHigh interspersion values if same patches	Dispersion (ai)	(Ode et al., 2009)
	are homogeneously distributed \rightarrow related to landscape preferences	Interspersion (iji)	(Hunziker and Kienast, 1999)
	Subdivision of patches \rightarrow landscape preference in agricultural	Number of patches	(Dramstad et al., 2006; Ode et al., 2009)
	landscape context	(np)	
	Diversity of patches landscape \rightarrow preference in agricultural	Shannon diversity	(Dramstad et al., 2006)
	landscape context \rightarrow complexity	index (shdi)	
	Evenness of patches, relation to landscape preferences	Shannon evenness	(Hunziker and Kienast, 1999; La Fuente Val et al., 2006)
		index (shei)	
Related exclusively on	RES		
Amount of edges	Aspect of visual contrast	(edges_res)	MATLAB canny edge detection
related to RES			
Amount of pixels		(pix_res)	Manual calculation with Adobe Photoshop
related to RES			
RGB Images decom	posed with 12 classes		
5	-		

Appendix C. Reliability analysis

Each participant rated the landscape qualities for six stimuli. Here we report Cronbach alpha separated for each of the six trails.

Landscape quality	RATING ORDER	NG ORDER							
	1st	2nd	3rd	4th	5th	6th			
Coherence	0.79	0.86	0.92	0.78	0.85	0.82			
Complexity	0.72	0.73	0.82	0.83	0.8	0.8			
Legibility	0.71	0.81	0.69	0.71	0.76	0.75			
Mystery	0.63	0.64	0.51	0.65	0.6	0.7			
N data subset	132	133	133	133	133	134			

Appendix D. Descriptive statistics

For the landscape qualities, we report the descriptive statistics separated by individual scenarios (Table D.1). Table D1. Descriptive statistics for the landscape qualities

Landscape qualities	SCEANRIOS													
	PLAT_U LOW	RB HIGH	PLAT_A LOW	GRI HIGH	JURA LOW	HIGH	PRE_AL LOW	PS HIGH	ALP_UR LOW	B HIGH	ALP_TO LOW	UR HIGH	ALP LOW	HIGH
COHERENCE [MED]	3.5	3.5	4	4	4.5	4	4.25	4	4	2.5	4	3	4	3.5
COHERENCE [SE]	0.14	0.14	0.1	0.15	0.13	0.15	0.14	0.18	0.16	0.13	0.16	0.14	0.13	0.18
COMPLEXITY [MED]	3.5	3.5	3	3	3.5	3	3.75	4	4	4	3	3	3.5	3.5
COMPLEXITY [SE]	0.14	0.13	0.09	0.14	0.1	0.12	0.1	0.12	0.13	0.09	0.13	0.11	0.13	0.15
LEGIBILITY [MED]	3.33	3	4	4	4.33	3.67	4	4	3.67	3.33	3.5	3.67	4	4
LEGIBILITY [SE]	0.12	0.12	0.09	0.12	0.09	0.1	0.12	0.11	0.13	0.11	0.09	0.1	0.09	0.12
MYSTERY_A [MED]	4	3	3	3	4	4	4	4	4	4	3	3	4	4
MYSTERY_A [SE]	0.13	0.15	0.12	0.16	0.13	0.12	0.1	0.13	0.16	0.12	0.17	0.14	0.14	0.17
MYSTERY_B [MED]	2	2	3	3	4	4	4	4	3	2	4	4	5	4
MYSTERY_B [SE]	0.15	0.15	0.14	0.17	0.12	0.15	0.12	0.15	0.16	0.15	0.17	0.15	0.11	0.16
Ν	57	55	68	52	65	53	56	48	49	70	46	64	59	57

	SCEANR	IOS												
	PLAT_UR	В	PLAT_AC	GRI	JURA		PRE_ALF	PS	ALP_URI	3	ALP_TO	UR .	ALP	
	LOW	HIGH	LOW	HIGH	LOW	HIGH	LOW	HIGH	LOW	HIGH	LOW	HIGH	LOW	HIGH
N CHOICE	17	24	33	26	41	32	38	32	19	16	28	27	44	24
N NOCHOICE	40	31	35	26	24	21	18	16	30	54	18	37	15	33
PREFERENCE	0.31	0.44	0.49	0.5	0.63	0.6	0.68	0.67	0.39	0.22	0.61	0.42	0.75	0.42

Table D2. Descriptive statistics for the visual features, measured on N = 14 scenarios

Image Statistics	Mean	SD	SE	Median
Pixel based measures				
hue	0.316	0.065	0.017	0.332
sat	0.352	0.044	0.012	0.369
bright	53.543	6.028	1.611	51.900
fc	2.466	0.360	0.096	2.488

(continued on next page)

(continued)

Image Statistics	Mean	SD	SE	Median
se	4.501	0.243	0.065	4.462
bytes	3201570.857	654711.232	174978.937	3058877.5
sf_low	7.756	0.178	0.048	7.763
sf_high	7.902	0.226	0.060	7.871
Landscape metrics				
ed	3447.157	1493.666	399.199	3041.314
shape	1.367	0.058	0.015	1.376
frac	1.259	0.031	0.008	1.26
ai	82.820	7.477	1.998	84.852
contig	0.287	0.049	0.013	0.270
cai	6.100	2.448	0.654	5.119
iji	58.024	5.238	1.400	59.245
np	126341.929	69948.262	18694.459	126,128
shdi	2.350	0.048	0.013	2.354
shei	0.946	0.019	0.005	0.948
Related exclusively on RES				
PIX_RES	20464.357	13465.360	3598.769	18277.000
EDGES_RES	6204.071	5539.226	1480.420	3864.500

Appendix E. Socio demographics

Fixed effects of age, gender and if the rated landscape was similar to the landscape where the participant live.

Landscape qualities	Fixed e	Fixed effects of socio demographics											
	$AGE \chi^2(1)$	Marginal R ²	Cohens f ²	р	HOME LA $\chi^2(1)$	ANDSCAPE Marginal R ²	Cohens f ²	р	GENDER $\chi^2(1)$	R Marginal R ²	Cohens f ²	р	
Coherence	7.009	0.144	0.17	0.008	0.655	0.131	0.15	0.418	0.068	0.13	0.15	0.794	
Complexity	0.114	0.142	0.17	0.736	0.109	0.147	0.17	0.741	1.127	0.151	0.18	0.288	
Legibility	0.101	0.129	0.15	0.749	36.923	0.131	0.15	1.229e-9	1.891	0.142	0.17	0.169	
Mystery_A	0.779	0.121	0.14	0.342	1.476	0.115	0.12	0.224	2.753	0.243	0.32	0.048	
Mystery_B	1.296	0.243	0.32	0.274	27.414	0.249	0.33	1.6e-7	0.773	0.247	0.33	0.379	

Appendix F. Correlation analysis

Correlation analysis with N=14 scenario and a corrected alpha level of 0.002. Correlation coefficient $\left(r\right)$

	COHERENCE	COMPLEXITY	LEGIBILITY	MYSTERY_A	MYSTERY_B
hue	0.275	-0.262	0.226	-0.525	-0.125
sat	-0.182	0.455	-0.429	0.495	0.121
bright	0.080	-0.549	0.177	-0.647	-0.372
fc	-0.578	0.169	-0.486	0.090	-0.451
se	-0.187	-0.547	-0.015	-0.147	0.020
bytes	-0.468	0.644	-0.534	0.604	-0.130
sf_low	-0.508	0.429	-0.521	0.099	-0.464
sf_high	0.288	0.081	0.648	0.376	0.886
ed	-0.415	0.424	-0.451	0.552	0.055
shape	-0.455	-0.204	-0.301	0.090	0.002
frac	-0.433	0.253	-0.508	0.380	-0.033
ai	0.415	-0.424	0.451	-0.552	-0.055
contig	0.174	-0.758	0.209	-0.631	-0.090
cai	0.073	-0.635	0.165	-0.613	-0.279
iji	-0.262	0.490	-0.358	0.244	-0.134
np	-0.240	0.587	-0.288	0.697	0.222
shdi	0.086	-0.090	0.059	-0.130	-0.046
shei	0.086	-0.090	0.059	-0.130	-0.046
PIX_RES	-0.446	0.055	-0.235	-0.284	-0.200
EDGES_RES	-0.749	-0.037	-0.314	-0.147	-0.262
Significance level ((p)				
	COHERENCE	COMPLEXITY	LEGIBILITY	MYSTERY_A	MYSTERY_B
hue	0.342	0.366	0.436	0.054	0.670
sat	0.533	0.102	0.126	0.072	0.681
bright	0.786	0.042	0.545	0.012	0.190
fc	0.030	0.563	0.078	0.759	0.106
se	0.523	0.043	0.958	0.615	0.946
bytes	0.091	0.013	0.049	0.022	0.659
sf_low	0.064	0.126	0.056	0.737	0.095
sf_high	0.318	0.782	0.012	0.185	0.000

(continued on next page)

R. Spielhofer et al.

(continued)

	COHERENCE	COMPLEXITY	LEGIBILITY	MYSTERY_A	MYSTERY_B
ed	0.140	0.131	0.106	0.041	0.852
shape	0.102	0.483	0.296	0.759	0.994
frac	0.122	0.383	0.064	0.180	0.911
ai	0.140	0.131	0.106	0.041	0.852
contig	0.553	0.002	0.474	0.016	0.759
cai	0.805	0.015	0.573	0.020	0.334
iji	0.366	0.075	0.208	0.401	0.648
np	0.409	0.027	0.318	0.006	0.446
shdi	0.771	0.759	0.840	0.659	0.876
shei	0.771	0.759	0.840	0.659	0.876
PIX_RES	0.110	0.852	0.418	0.326	0.493
EDGES_RES	0.002	0.899	0.274	0.615	0.366

Appendix G. Visual features

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
PIX_RES					
LANDSCAPE	6	4.49E + 08	7.48E + 07	4.42	0.047 *
RES_LEVEL	1	1.81E + 09	1.81E + 09	106.71	4.81e-05 ***
Residuals	6	1.02E + 08	1.69E + 07		
EDGES_RES					
LANDSCAPE	6	76,886,638	12,814,440	1.289	0.383
RES LEVEL	1	262,337,401	262,337,401	26.385	0.002 **
Residuals	6	59.655.324	9.942.554		
cai mn			- ,		
LANDSCAPE	6	76.52	12.753	55.322	5.45-05 ***
RES LEVEL	1	0.01	0.006	0.024	0.882
Residuals	6	1.38	0.231		
contig mn	0	100	01201		
LANDSCADE	6	0.03	0.005	43 156	0 0001 ***
RES I EVEL	1	>0.001	>0.0001	0 101	0.761
REO_EEVEE	6	0.0006	0.0001	0.101	0.701
	0	0.0000	0.0001		
	6	2 607	0.601	60.276	0.01 × 05 ***
LANDSCAPE	6	3.607	0.601	69.376	2.816-05
RES_LEVEL	I	0.007	0.007	0.862	0.389
Residuals	6	0.052	0.87		
frac_mn					
LANDSCAPE	6	0.012	0.002	17.521	0.001 **
RES_LEVEL	1	>0.0001	>0.0001	0.228	0.649
Residuals	6	0.0007	0.0001		
shape_mn					
LANDSCAPE	6	0.039	0.006	14.776	0.002 **
RES_LEVEL	1	0.0009	0.0008	1.896	0.217
Residuals	6	0.003	0.0004		
ai					
LANDSCAPE	6	718.1	119.68	84.846	1.55e-05 ***
RES_LEVEL	1	0.3	0.26	0.181	0.685
Residuals	6	8.5	1.41		
ed					
LANDSCAPE	6	28.656.295	4.776.049	85.031	1.54e-05 ***
RES LEVEL	1	10.194	10.194	0.181	0.685
Residuals	6	337 010	56 168	01101	0.000
iii	0	007,010	00,100		
I ANDSCADE	6	312.04	52.01	7.077	0.015 *
DECIEVEI	1	0.58	0.58	0.079	0.788
RES_LEVEL Residuels	1	0.58	7.25	0.079	0.788
nesiduais	0	44.09	7.33		
np LANDCCADE	<i>r</i>	C 16E + 10	1.005 + 10	20.072	0 0001 ***
LANDSCAPE	6	6.16E + 10	1.03E + 10	38.072	0.0001
RES_LEVEL	I	4.38E + 08	4.38E + 08	1.627	0.249
Residuals	6	1.62E + 09	2.69E + 08		
shdi					
LANDSCAPE	6	0.014	0.003	0.881	0.559
RES_LEVEL	1	0.0004	0.0004	0.169	0.695
Residuals	6	0.016	0.003		
shei					
LANDSCAPE	6	0.002	0.0004	0.881	0.559
RES_LEVEL	1	>0.0001	>0.0001	0.169	0.695
Residuals	6	0.003	0.0004		
sf_low					
LANDSCAPE	6	0.319	0.053	3.866	0.062 .
RES LEVEL	1	0.013	0.013	0.918	0.375
Residuals	-	0.082	0.014	5.710	2.07.0
sf high	v	0.002	0.011		
LANDSCAPE	6	0.655	0.109	72 847	2 430-05 ***
	0	0.000	0.109	/ 2.0 1/	2.100.00

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
RES_LEVEL	1	0.001	0.001	0.879	0.385
Residuals	6	0.009	0.002		
frac					
LANDSCAPE	6	0.078	0.013	1119.783	7.09e-09 ***
RES_LEVEL	1	>0.0001	>0.0001	3.523	0.11
Residuals	6	>0.0001	>0.0001		
bright					
LANDSCAPE	6	472.4	78.73	1.02e30	<2e16 ***
RES_LEVEL	1	0	0	1.18e - 01	0.643
Residuals	6	0	0		
bytes					
LANDSCAPE	6	5.57E + 12	9.29E + 11	18817.5	1.5e-12 ***
RES_LEVEL	1	7.97E + 08	7.97E + 08	16.15	0.007 **
Residuals	6	2.96E + 08	4.94E + 07		
ec					
LANDSCAPE	6	2.00E + 09	333,382,549	842.367	1.66e-08 ***
RES_LEVEL	1	6.98E + 05	697,991	1.764	0.232
Residuals	6	2.38E + 06	395,769		
fc					
LANDSCAPE	6	1.6839	0.281	8456.39	1.65e-11 ***
RES_LEVEL	1	0.001	0.001	28.78	0.002 **
Residuals	6	0.0002	>0.0001		
se					
LANDSCAPE	6	0.765	0.128	631.467	3.94e-08 ***
RES_LEVEL	1	0.001	0.001	6.069	0.049 *
Residuals	6	0.001	>0.0001		
hue					
LANDSCAPE	6	0.055.	0.009	17753.9	1.79e-12 ***
RES_LEVEL	1	>0.0001	>0.0001	39.6	0.0007 ***
Residuals	6	>0.0001	>0.0001		
sat					
LANDSCAPE	6	0.026	0.004	80.694	1.85e-10 ***
RES_LEVEL	1	>0.0001	>0.0001	2.557	0.161
Residuals	6	>0.0001	>0.0001		
Signif. co		:0 '***'	0.001 '**	' 0.01 '	*' 0.05 '.' 0.1 '

Landscape and Urban Planning 209 (2021) 104000

Appendix H. Preferences

R. Spielhofer et al.

H.1 Relation (R & p) between visual features and preference score

	R	р
hue	0.301	0.296
sat	0.011	0.970
bright	-0.151	0.607
fc	-0.692	0.006
se	-0.187	0.523
bytes	-0.437	0.118
sf_low	-0.556	0.039
sf_high	0.618	0.019
ed	-0.327	0.253
shape	-0.319	0.267
frac	-0.319	0.267
ai	0.327	0.253
contig	0.002	0.994
cai	-0.121	0.681
iji	-0.209	0.474
np	-0.108	0.714
shdi	0.046	0.876
shei	0.046	0.876
PIX_RES	-0.011	0.970
EDGES_RES	-0.305	0.288
es and preference scores		
	R	р
COHEBENCE	0.710	0.004

H.2 Relation between landscape qualities and preference score

	R	р
COHERENCE	0.710	0.004
COMPLEXITY	-0.024	0.935
LEGIBILITY	0.723	0.003
MYSTERY_A	0.204	0.483
MYSTERY_B	0.793	0.001

19

Appendix I. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.landurbplan.2020.104000. The following link to the papers data repository on mendeley data: https://dx.doi.org/10.17632/wvf56j6nps.1.

References

- Appleton, Jay (1996). The experience of landscape. Book. doi:9780471962359.
- Apostol, D., Palmer, J., Pasqualetti, M., Smardon, R., & Sullivan, R. (2016). The renewable energy landscape: preserving scenic values in our sustainable future. Florence: Taylor and Francis.
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software.*, 67(1). https://doi.org/10.18637/ jss.v067.i01.
- Bell, S. (2012). Landscape: pattern, perception and process (2nd ed.). Hoboken: Taylor and Francis.
- Berman, M. G., Hout, M. C., Kardan, O., Hunter, M. R., Yourganov, G., Henderson, J. M., et al. (2014). The perception of naturalness correlates with low-level visual features of environmental scenes. *PLoS ONE*, 9(12), e114572. https://doi.org/10.1371/ journal.pone.0114572.s002.
- Betakova, V., Vojar, J., & Sklenicka, P. (2015). Wind turbines location: how many and how far? Applied Energy, 151, 23–31. https://doi.org/10.1016/j. apenergy.2015.04.060.
- Bishop, I. D., & Miller, D. R. (2007). Visual assessment of off-shore wind turbines: the influence of distance, contrast, movement and social variables. *Renew. Energy*, 32(5), 814–831. https://doi.org/10.1016/j.renene.2006.03.009.
- Brahms, E., Peters, J. 2012. Germ: Landschaftsbild, Erholungsnutzung und Windenergieanlagen in der Planungsregion Magdeburg. (Visual landscape, recreation and wind turbines in the region of Magdeburg).
- Cattin, R., Schaffner, B., Humar-Mägli, T., Albrecht, S., Remund, J., Klauser, D., et al. (2012). Energiestrategie 2050 – Berechnung der Energiepotenziale für Wind- und Sonnenenergie. (Energy Strategy 2050 – Calculation of wind and solar energy potential for Switzerland), Bern.
- Chiabrando, R., Fabrizio, E., & Garnero, G. (2009). The territorial and landscape impacts of photovoltaic systems: Definition of impacts and assessment of the glare risk. *Renewable and Sustainable Energy Reviews*, 13(9), 2441–2451. https://doi.org/ 10.1016/j.rser.2009.06.008.
- Coeterier, J. F. (1996). Dominant attributes in the perception and evaluation of the Dutch landscape. Landscape and Urban Planning, 34(1), 27–44. https://doi.org/10.1016/ 0169-2046(95)00204-9.
- Cohen, J. (1992). Statistical power analysis. Current Directions in Psychological Science, 1 (3), 98–101. https://doi.org/10.1111/1467-8721.ep10768783.
- Council of Europe. (2000). European Landscape Convention: ECL.
- Courcoux, P., & Semenou, M. (1997). Preference data analysis using a paired comparison mode. Food Quality and Performance, 8(5), 353–358.
- Cushman, S. A., McGarigal, K., & Neel, M. C. (2008). Parsimony in landscape metrics: strength, universality, and consistency. *Ecological Indicators*, 8(5), 691–703. https:// doi.org/10.1016/j.ecolind.2007.12.002.
- Devine-Wright, P., & Howes, Y. (2010). Disruption to place attachment and the protection of restorative environments: a wind energy case study. *Journal of Environmental Psychology*, 30(3), 271–280. https://doi.org/10.1016/j. jenvp.2010.01.008.
- Dramstad, W. E., Tveit, M. S., Fjellstad, W. J., & Fry, G. L. A. (2006). Relationships between visual landscape preferences and map-based indicators of landscape structure. *Landscape and Urban Planning*, 78(4), 465–474. https://doi.org/10.1016/j. landurbplan.2005.12.006.
- A. Forsythe M. Nadal N. Sheehy C.J. Cela-Conde M. Sawey 2011 Predicting beauty: fractal dimension and visual complexity in art British Journal of Psychology (London, England 1953) 102 1 49–70 10.1348/000712610X498958.
- Frank, S., Fürst, C., Koschke, L., Witt, A., & Makeschin, F. (2013). Assessment of landscape aesthetics—validation of a landscape metrics-based assessment by visual estimation of the scenic beauty. *Ecological Indicators*, 32, 222–231. https://doi.org/ 10.1016/j.ecolind.2013.03.026.
- Fry, G., Tveit, M. S., Ode, Å., & Velarde, M. D. (2009). The ecology of visual landscapes: exploring the conceptual common ground of visual and ecological landscape indicators. *Ecological Indicators*, 9(5), 933–947. https://doi.org/10.1016/j. ecolind.2008.11.008.
- Giampaoli, V., & Singer, J. M. (2009). Likelihood ratio tests for variance components in linear mixed models. *Journal of Statistical Planning and Inference*, 139(4), 1435–1448. https://doi.org/10.1016/j.jspi.2008.06.016.
- Gobster, P. H. (1999). An ecological aesthetic for forest landscape management. Landscape Journal, 18(1), 54–64. https://doi.org/10.3368/lj.18.1.54.
- Gonseth, Y., Wohlgemuth, T., Sansonnens, B., & Buttler, A. (2001). Die biogeographischen Regionen der Schweiz. Bern: Erläuterungen und Einteilungsstandard.
- Grübel, J., Weibel, R., Jiang, M. H., Hölscher, C., Hackman, D. A., & Schinazi, V. R. (2017). EVE: a framework for experiments in virtual environments. In T. Barkowsky, H. Burte, C. Hölscher, & H. Schultheis (Eds.), Spatial cognition X: 13th Biennial Conference, Kogwis 2016, Bremen, Germany, September 26–30, 2016, and 10th International Conference, Spatial Cognition 2016, Philadelphia (pp. 159–176). New York Inc, Cham: Springer-Verlag.

- Hagerhall, C. M., Laike, T., Taylor, R. P., Küller, M., Küller, R., & Martin, T. P. (2008). Investigations of human EEG response to viewing fractal patterns. *Perception*, 37(10), 1488–1494. https://doi.org/10.1068/p5918.
- Hägerhäll, C. M., Ode Sang, Å., Englund, J.-E., Ahlner, F., Rybka, K., Huber, J., & Burenhult, N. (2018). Do humans really prefer semi-open natural landscapes? a cross-cultural reappraisal. *Frontiers in Psychology*, 9, 822. https://doi.org/10.3389/ fpsyg.2018.00822.
- Hastik, R., Basso, S., Geitner, C., Haida, C., Poljanec, A., Portaccio, A., et al. (2015). Renewable energies and ecosystem service impacts. *Renewable and Sustainable Energy Reviews*, 48, 608–623. https://doi.org/10.1016/j.rser.2015.04.004.
- Hesselbarth, M. H. K., Sciaini, M., With, K. A., Wiegand, K., & Nowosad, J. (2019). Landscapemetrics: an open-source R tool to calculate landscape metrics. *Ecography*, 42(10), 1648–1657. https://doi.org/10.1111/ecog.04617.
- Hunziker, M., Felber, P., Gehring, K., Buchecker, M., Bauer, N., & Kienast, F. (2008). Evaluation of landscape change by different social groups: results of two empirical studies in Switzerland. *Mountain Research and Development*, 28(2), 140–147. https:// doi.org/10.1659/mrd.0952.
- Hunziker, M., & Kienast, F. (1999). Potential impacts of changing agricultural activities on scenic beauty – a prototypical technique for automated rapid assessment. *Landscape Ecology*, 14(2), 161–176. https://doi.org/10.1023/A:1008079715913.
- Ibarra, F. F., Kardan, O., Hunter, M. R., Kotabe, H. P., Meyer, F. A. C., & Berman, M. G. (2017). Image feature types and their predictions of aesthetic preference and naturalness. *Frontiers in Psychology*, 8, 632. https://doi.org/10.3389/ fpsyg.2017.00632.
- Johansson, M., & Laike, T. (2007). Intention to respond to local wind turbines: the role of attitudes and visual perception. Wind Energy, 10(5), 435–451. https://doi.org/ 10.1002/we.232.
- Kaplan, R., & Kaplan, S. (1989). The experience of nature a psychological perspective. Cambridge: Cambridge University Press.
- Kaplan, S. (1987). Aesthetics, affect, and cognition: environmental preference from an evolutionary perspective. *Environment and Behavior*, 19(1), 3–32. https://doi.org/ 10.1177/0013916587191001.
- Kardan, O., Demiralp, E., Hout, M. C., Hunter, M. R., Karimi, H., Hanayik, T., et al. (2015). Berman Is the preference of natural versus man-made scenes driven by bottom-up processing of the visual features of nature? *Frontiers in Psychology*, *6*, 471. https://doi.org/10.3389/fpsyg.2015.00471.
- Kienast, F., Frick, J., van Strien, M. J., & Hunziker, M. (2015). The Swiss landscape monitoring program – a comprehensive indicator set to measure landscape change. *Ecological Modelling*, 295, 136–150. https://doi.org/10.1016/j. ecolmodel.2014.08.008.
- Kienast, F., Huber, N., Hergert, R., Bolliger, J., Moran, L. S., & Hersperger, A. M. (2017). Conflicts between decentralized renewable electricity production and landscape services – a spatially-explicit quantitative assessment for Switzerland. *Renewable and Sustainable Energy Reviews*, 67, 397–407. https://doi.org/10.1016/j. rser.2016.09.045.
- Kuiper, J. (1998). Landscape quality based upon diversity, coherence and continuity. Landscape and Urban Planning, 43(1-3), 91–104. https://doi.org/10.1016/S0169-2046(98)00075-9.
- Kuper, R. (2017). Evaluations of landscape preference, complexity, and coherence for designed digital landscape models. *Landscape and Urban Planning*, 157, 407–421. https://doi.org/10.1016/j.landurbplan.2016.09.002.
- La Fuente Val, G.de, Atauri, J. A., & de Lucio, J. V. (2006). Relationship between landscape visual attributes and spatial pattern indices: a test study in Mediterraneanclimate landscapes. *Landscape and Urban Planning*, 77(4), 393–407. https://doi.org/ 10.1016/j.landurbplan.2005.05.003.
- Lothian, A. (2008). Scenic perceptions of the visual effects of wind farms on south Australian landscapes. *Geographical Research*, 46(2), 196–207. https://doi.org/ 10.1111/j.1745-5871.2008.00510.x.
- Lüdecke, D. (2020). Sjstats: Collection of convenient functions for common statistical computations.
- Manyoky, M., Wissen Hayek, U., Pieren, R., Heutschi, K., Grêt-Regamey, A., et al. (2016). Evaluating a visual-acoustic simulation for wind park assessment. *Landscape and Urban Planning*. https://doi.org/10.1016/j.landurbplan.2016.03.013.
- Maehr, A. M., Watts, G. R., Hanratty, J., & Talmi, D. (2015). Emotional response to images of wind turbines: a psychophysiological study of their visual impact on the landscape. *Landscape and Urban Planning*, 142, 71–79. https://doi.org/10.1016/j. landurbplan.2015.05.011.

Mandelbrot, B. B. (1983). The fractal geometry of nature, Updated and augmented. New York, NY: Freeman.

- Mathiesen, B. V., Lund, H., & Karlsson, K. (2011). 100% Renewable energy systems, climate mitigation and economic growth. *Applied Energy*, 88(2), 488–501. https:// doi.org/10.1016/j.apenergy.2010.03.001.
- McGarigal, K., Cushman, S. A., & Ene, E. (2012). FRAGSTATS: Spatial Pattern Analysis Program for Categorical and Continuous Maps. Computer software program produced by the authors at the University of Massachusetts, Amherst.
- Molnarova, K., Sklenicka, P., Stiborek, J., Svobodova, K., Salek, M., & Brabec, E. (2012). Visual preferences for wind turbines: location, numbers and respondent

R. Spielhofer et al.

characteristics. Applied Energy, 92, 269–278. https://doi.org/10.1016/j. apenergy.2011.11.001.

Nadaï, A., & van der Horst, D. (2010). Introduction: landscapes of energies. Landscape Research, 35(2), 143–155. https://doi.org/10.1080/01426390903557543.

- Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4(2), 133–142. https://doi.org/10.1111/j.2041-210x.2012.00261.x.
- Ode, Å., Fry, G., Tveit, M. S., Messager, P., & Miller, D. (2009). Indicators of perceived naturalness as drivers of landscape preference. *Journal of Environment Management*, 90(1), 375–383. https://doi.org/10.1016/j.jenvman.2007.10.013.
- O'Neill, R. V., Krummel, J. R., Gardner, R. H., Sugihara, G., Jackson, B., DeAngelis, D. L., et al. (1988). Indices of landscape pattern. *Landscape Ecology*, 1(3), 153–162. https://doi.org/10.1007/BF00162741.

Orians, G. H. (1980). General theory and applications to human behavior. In J. S. Lockard (Ed.), *The Evolution of human social behavior* (pp. 49–63). New York: Elsevier.

- Palmer, J. F. (2000). Reliability of rating visible landscape qualities. Landscape Journal, 19(1–2), 166–178. https://doi.org/10.3368/lj.19.1-2.166.
- Palmer, J. F. (2004). Using spatial metrics to predict scenic perception in a changing landscape: Dennis, Massachusetts. *Landscape and Urban Planning*, 69(2–3), 201–218. https://doi.org/10.1016/j.landurbplan.2003.08.010.
- Pasqualetti, M. J. (2000). Morality, space, and the power of wind-energy landscapes. Geographical Review, 90(3), 381–394.
- Plieninger, T., Draux, H., Fagerholm, N., Bieling, C., Bürgi, M., Kizos, T., et al. (2016). The driving forces of landscape change in Europe: a systematic review of the evidence. *Land Use Policy*, 57, 204–214. https://doi.org/10.1016/j. landusepol.2016.04.040.

R Core Team. (2018). R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.

- Rosenholtz, R., Li, Y., & Nakano, L. (2007). Measuring visual clutter. Journal of Vision, 7 (2), 17. https://doi.org/10.1167/7.2.17.
- Santangeli, A., Di Minin, E., Toivonen, T., Pogson, M., Hastings, A., Smith, P., et al. (2016). Synergies and trade-offs between renewable energy expansion and biodiversity conservation – a cross-national multifactor analysis. *GCB Bioenergy*, 8 (6), 1191–1200. https://doi.org/10.1111/gcbb.12337.
- Schirpke, U., Tappeiner, G., Tasser, E., & Tappeiner, U. (2019). Using conjoint analysis to gain deeper insights into aesthetic landscape preferences. *Ecological Indicators*, 96, 202–212. https://doi.org/10.1016/j.ecolind.2018.09.001.
- Shannon, C. E. (1948). A mathematical theory of communication. Bell System Technical Journal, 27(3), 379–423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x.
- Singh, S. N., Todd Donavan, D., Mishra, S., & Little, T. D. (2008). The latent structure of landscape perception: a mean and covariance structure modeling approach. *Journal* of Environmental Psychology, 28(4), 339–352. https://doi.org/10.1016/j. ienvp.2008.03.004.
- Stamps, A. E., III (1999). Sex, complexity, and preferences for residential facades. Perceptual and Motor Skills, 88(3_suppl), 1301–1312. https://doi.org/10.2466/ pms.1999.88.3c.1301.
- Stamps, A. E., III (2004). Mystery, complexity, legibility and coherence: a meta-analysis. Journal of Environmental Psychology, 24(1), 1–16. https://doi.org/10.1016/S0272-4944(03)00023-9.
- Spielhofer, Reto, Thrash, T., Wissen Hayek, U., Grêt-Regamey, A., Salak, B., Grübel, J., & Schinazi, V. R. (2021). Physiological and behavioral reactions to renewable energy systems in various landscape types. *Renewable and Sustainable Energy Reviews*, 135. https://doi.org/10.1016/j.rser.2020.110410.
- Stanton, Caroline (2016). Perception of scale and scale effects in the landscape, with specific reference to wind turbines in Scotland. The University of Edindburgh. http://hdl.ha ndle.net/1842/22941.
- Steg, L. (2013). Environmental psychology: an introduction (first ed.). Chichester: Wiley-Blackwell.
- Strumse, E. (1996). Demographic differences in the visual preferences for agrarian landscapes in western Norway. *Journal of Environmental Psychology*, 16(1), 17–31. https://doi.org/10.1006/jevp.1996.0002.
- Swiss Federal Office of Energy. (2012). Die Energieperspektiven für die Schweiz bis 2050 (Energy perspectives for Switzerland up to the year 2050), Bern, Basel.

Swiss Federal Office of Energy. (2019). Overall energy statistics.

- Tjur, T. (2009). Coefficients of determination in logistic regression models—a new proposal: the coefficient of discrimination. *The American Statistician*, 63(4), 366–372. https://doi.org/10.1198/tast.2009.08210.
- Torres Sibille, A. D. C., Cloquell-Ballester, V.-A., Cloquell-Ballester, V.-A., & Darton, R. (2009). Development and validation of a multicriteria indicator for the assessment of objective aesthetic impact of wind farms. *Renewable and Sustainable Energy Reviews*, 13(1), 40–66. https://doi.org/10.1016/j.rser.2007.05.002.
- Torres-Sibille, A. D. C., Cloquell-Ballester, V.-A., Cloquell-Ballester, V.-A., & Artacho Ramírez, M.Á. (2009). Aesthetic impact assessment of solar power plants: an objective and a subjective approach. *Renewable and Sustainable Energy Reviews*, 13 (5), 986–999. https://doi.org/10.1016/j.rser.2008.03.012.
- Tuch, A. N., Bargas-Avila, J. A., Opwis, K., & Wilhelm, F. H. (2009). Visual complexity of websites: Effects on users' experience, physiology, performance, and memory. *International Journal of Human-Computer Studies*, 67(9), 703–715. https://doi.org/ 10.1016/j.ijhcs.2009.04.002.
- Tveit, M., Ode, Å., & Fry, G. (2006). Key concepts in a framework for analysing visual landscape character. Landscape Research, 31(3), 229–255. https://doi.org/10.1080/ 01426390600783269.
- Twigger-Ross, C. L., & Uzzell, D. L. (1996). Place and identity processes. Journal of Environmental Psychology, 16(3), 205–220. https://doi.org/10.1006/ jevp.1996.0017.
- Ulrich, R. S. (1986). Human responses to vegetation and landscapes. Landscape and Urban Planning, 13, 29–44. https://doi.org/10.1016/0169-2046(86)90005-8.
- Ulrich, R. S., Simons, R. F., Losito, B. D., Fiorito, E., Miles, M. A., & Zelson, M. (1991). Stress recovery during exposure to natural and urban environments. *Journal of Environmental Psychology*, 11(3), 201–230. https://doi.org/10.1016/S0272-4944 (05)80184-7.
- Valtchanov, D., & Ellard, C. G. (2015). Cognitive and affective responses to natural scenes: effects of low level visual properties on preference, cognitive load and eyemovements. *Journal of Environmental Psychology*, 43, 184–195. https://doi.org/ 10.1016/j.jenvp.2015.07.001.
- Van den Berg, A. E., & Koole, S. L. (2006). New wilderness in the Netherlands: an investigation of visual preferences for nature development landscapes. *Landscape and Urban Planning*, 78(4), 362–372. https://doi.org/10.1016/j. landurplan.2005.11.006.
- van der Jagf, A. P. N., Craig, T., Anable, J., Brewer, M. J., & Pearson, D. G. (2014). Unearthing the picturesque: the validity of the preference matrix as a measure of landscape aesthetics. *Landscape and Urban Planning*, 124, 1–13. https://doi.org/ 10.1016/i.landurbolan.2013.12.006.
- Ware, C. (2011). Visual thinking for design. Morgan Kaufmann series in interactive technologies. Amsterdam: Elsevier/Morgan Kaufmann.
- Warren, C. R., Lumsden, C., O'Dowd, S., & Birnie, R. V. (2005). 'Green On Green': public perceptions of wind power in Scotland and Ireland. *Journal of Environmental Planning* and Management, 48(6), 853–875. https://doi.org/10.1080/09640560500294376.
- Wolsink, M. (2007). Wind power implementation: The nature of public attitudes: equity and fairness instead of 'backyard motives'. *Renewable and Sustainable Energy Reviews*, 11(6), 1188–1207. https://doi.org/10.1016/j.rser.2005.10.005.
- Wolsink, M. (2018). Co-production in distributed generation: renewable energy and creating space for fitting infrastructure within landscapes. *Landscape Research*, 43(4), 542–561. https://doi.org/10.1080/01426397.2017.1358360.
- World Energy Council. (2019). World Energy Scenarios 2019: European regional perspective, London.
- Wüstenhagen, R., Wolsink, M., & Bürer, M. J. (2007). Social acceptance of renewable energy innovation: an introduction to the concept. *Energy Policy*, 35(5), 2683–2691. https://doi.org/10.1016/j.enpol.2006.12.001.
- Zheng, B., Zhang, Y., & Chen, J. (2011). Preference to home landscape: wildness or neatness? *Landscape and Urban Planning*, 99(1), 1–8. https://doi.org/10.1016/j. landurbplan.2010.08.006.
- Zoellner, J., Schweizer-Ries, P., & Wemheuer, C. (2008). Public acceptance of renewable energies: results from case studies in Germany. *Energy Policy*, 36(11), 4136–4141. https://doi.org/10.1016/j.enpol.2008.06.026.