

Machine learning reveals climate forcing from aerosols is dominated by increased cloud cover

Journal Article

Author(s):

Chen, Ying; Haywood, Jim; Wang, Yu; Malavelle, Florent; Jordan, George; Partridge, Daniel; Fieldsend, Jonathan; De Leeuw, Johannes; Schmidt, Anja; Cho, Nayeong; Oreopoulos, Lazaros; Platnick, Steven; Grosvenor, Daniel; Field, Paul; Lohmann, Ulrike



Publication date:

2022-08

Permanent link:

<https://doi.org/10.3929/ethz-b-000561341>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Originally published in:

Nature Geoscience 15, <https://doi.org/10.1038/s41561-022-00991-6>

Supplementary Information:

Machine-learning reveals climate forcing from aerosols is dominated by increased cloud cover

Ying Chen^{1*,#}, Jim Haywood^{1,2}, Yu Wang³, Florent Malavelle⁴, George Jordan², Daniel Partridge¹, Jonathan Fieldsend¹, Johannes De Leeuw⁵, Anja Schmidt^{5,6,†}, Nayeong Cho⁷, Lazaros Oreopoulos⁷, Steven Platnick⁷, Daniel Grosvenor⁸, Paul Field^{4,9}, Ulrike Lohmann³

¹College of Engineering, Mathematics, and Physical Sciences, University of Exeter, UK

²Met Office Hadley Centre, Exeter, UK

³Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland

⁴Met Office, Exeter, UK

⁵Centre for Atmospheric Science, Yusuf Hamied Department of Chemistry, University of Cambridge, UK

⁶Department of Geography, University of Cambridge, UK

⁷Earth Sciences Division, NASA GSFC, Greenbelt, Maryland, USA

⁸National Centre for Atmospheric Sciences, University of Leeds, Leeds, UK

⁹School of Earth and Environment, University of Leeds, Leeds, UK

*Correspondence to: Ying Chen (y.chen6@exeter.ac.uk; ying.chen@psi.ch)

#Now at Laboratory of Atmospheric Chemistry, Paul Scherrer Institut, Villigen, Switzerland

†Now at Institute of Atmospheric Physics (IPA), German Aerospace Center (DLR), Oberpfaffenhofen, Germany and Meteorological Institute, Ludwig Maximilian University of Munich, Munich, Germany

This PDF file includes:

Discussion:

Section S1– Discussion of results in September 2014;

Section S2– Discussion on the influence of the sea-surface temperature anomaly in 2014 in disentangling the ACI-induced CF increase;

Tables:

Table S1 – The explanatory variables for machine-learning.

32 **Supplementary Discussion**

33 **Section S1: Discussion of results in September 2014**

34 The Holuhraun effusive eruption also resulted in a massive aerosol plume in the lower
35 troposphere in September 2014. Unlike October 2014, the unusual easterly wind in September
36 2014 brought European outflow with anthropogenic aerosol to the southeast part of the
37 geographical region (latitude $< 63^{\circ}\text{N}$, longitude $> 30^{\circ}\text{W}$)¹⁵ in addition to the volcanic
38 plume^{15,63}.

39 The predictions of cloud properties using the machine-learning surrogate MODIS (ML-
40 MODIS) also validate well with the MODIS observations in non-eruption Septembers during
41 2001-2020 (left panels in Extended Data Fig. S7). In line with October, a clear difference
42 between ML-MODIS prediction and MODIS observation is observed in September 2014 due
43 to the volcanic eruption (right panels in Extended Data Fig. S7). The volcanic aerosol-
44 perturbation led to a clear increase in N_d and a decrease in r_{eff} , as expected, especially over the
45 northeast quarter of the geographical region (Extended Data Fig. S9a and S9b), which is
46 dominated by volcanic aerosol plume¹⁵. The enhanced N_d and the resultant Twomey r_{eff} effect
47 are clearly discerned over all latitude bands, and lead to a higher cloud fraction (Extended
48 Data Fig. S9c). The spatial patterns of volcanic aerosol-perturbation induced changes in cloud
49 properties are similar to the spatial patterns of climatological anomalies (right panels in
50 Extended Data Fig. S9). However, the unusual easterly European outflow increases aerosol
51 loading in the southeast part of the region and leads to stronger cloud responses than that
52 induced solely by Holuhraun plume. This noise cannot be ruled out using climatological
53 anomaly analysis¹⁵. Our machine-learning approach is able to account for meteorology
54 variability, and rules out the noise driven by the unusual easterly flow. We therefore quantify

55 significantly weaker responses in N_d , r_{eff} and CF using the machine-learning approach over
56 the southeast part of the region (compared left and right panels in Fig. S9). This again further
57 demonstrates the viability of our machine-learning approach in identifying changes in cloud
58 created by volcanic aerosols above and beyond the expected meteorological variability.

59 Our Monte Carlo analysis also shows statistically significant changes in N_d , r_{eff} and CF due to
60 the volcanic eruption. The variability of N_d and r_{eff} response signals lie outside the uncertainty
61 range (Extended Data Fig. S8a). Most parts of CF response signal lies outside the uncertainty
62 range, with a significant shift to higher cloud fraction (Extended Data Fig. S8a). The N_d is
63 increased by 22% on average, leading to a 6% decrease in r_{eff} and a 5% relative increase in CF
64 on median (and average) over the domain. We observe a weak LWP increase of 2% (1.01
65 minus 0.99) on average in September 2014, but it is not significant because the signal
66 variability lies in the uncertainty range (Extended Data Fig. S8a). The susceptibilities of r_{eff} ,
67 LWP and CF to N_d are estimated as 0.31 [CI90: 0.15 ~ 0.59], 0.10 [CI90: -0.11 ~ 0.42] and
68 0.25 [CI90: -0.10 ~ 0.55], respectively, where CI90 stands for 90% confidence interval.
69 Therefore, according to Eq. (3), the relative contributions to ACI-induced radiative forcing are
70 $46 \pm 29\%$ (CF adjustment), $42 \pm 16\%$ (Twomey r_{eff} effect) and $12 \pm 20\%$ (LWP adjustment),
71 respectively. It is worth noting that these values in September are potentially less
72 representative of the global distribution of cloud regimes than in October, because of the more
73 limited areal extent of the plume¹⁵. However, all of these responses are in line with the findings
74 in October, providing further evidence of that our findings are robust.

75

76 **Section S2: Discussion on the influence of the sea-surface temperature (SST) anomaly in**
77 **2014 in disentangling the ACI-induced CF increase**

78 In October 2014, a cold SST anomaly developed to the south ($45^{\circ}\text{N} \sim 60^{\circ}\text{N}$, $20^{\circ}\text{W} \sim 45^{\circ}\text{W}$) of
79 the study domain, owing to factors that appear to be independent from the Holuhraun
80 eruption⁴². These colder SSTs could favor a higher low-level liquid CF even without volcanic
81 aerosol perturbation, due to enhanced static stability and thinner boundary layers⁶⁴⁻⁶⁷. While
82 such a confounding factor induced by SST anomaly is not accounted for in the climatological
83 analysis using only MODIS observations, our machine-learning (ML) approach however
84 accounts for it, because CF results from ML-MODIS predictions experience the same SST
85 conditions as the MODIS observations.

86 The cold SST anomaly does not undermine the ML representation of SST variability. The SST
87 in 2014 lies entirely in the variability range of the ML training dataset (see Extended Data Fig.
88 S10a). The prediction of CF over the anomaly region is based on the ML trained by the large
89 dataset over the entire study domain spanning the years 2001-2020, which consists of 64,713
90 pairs of training data. The cold SST anomaly in 2014 actually shifts the SST probability
91 distribution towards the center of the SST distribution of the training dataset, instead of
92 shifting it outside the range of variability (blue bars in Extended Data Fig. S10a). However,
93 there remains the possibility that the co-variation of meteorological variables associated with
94 the cold SST anomaly may result in multi-variate conditions that are not well captured by the
95 range found in the training conditions. We therefore perform a new Monte Carlo ML analysis
96 which excludes the regions where for the October 2014 SSTs lie outside of the climatological
97 range for the same place. This extreme cold SST anomalies occurred in fewer than 5% of the
98 pixels in the domain. We find a negligible difference between this new ML analysis (Extended

99 Data Fig. S8b) and the initial ML analysis (Fig. 3), indicating that the ACI signals derived
100 using our ML approach are not significantly impacted or contaminated by the SST anomaly.

101 The strong cold SST anomaly is limited to a region south of approximately 60°N (Extended
102 Data Fig. S5a), while the impact of Twomey effect (a well-documented indicator of ACI^{8,9,15,18})
103 is clearly seen Atlantic-wide (Fig. 2b), and coincide with an Atlantic-wide CF response (Fig.
104 2c). Compared with climatological analysis (Extended Data Fig. S3c), one of the impacts of
105 the ML approach is to reduce the CF response in the south where SSTs are below average
106 (most clearly seen in the zonal mean at ~52°N, Fig. 2c .vs. Extended Data Fig. S3c). These
107 indicate that ML is able to distinguish the extra CF increase due to aerosol on top of the likely
108 CF increase due to SST-covariant factors.

109 To further demonstrate the fidelity of the ML approach in disentangling ACI signals, the total
110 impact of the SST anomaly on CF should be indicated by anomalous low-level cloud cover
111 (LCC) in ERA5 reanalysis, where the volcanic aerosol is not included. Despite the higher LCC
112 in October 2014 to the south due to cold SST anomaly, this cloud fraction increase in the
113 ERA5 anomaly analysis (which accounts for meteorology only) is significantly less than the
114 perturbation that is derived from either ML-MODIS (which accounts for aerosols) or MODIS
115 alone (which accounts for aerosols and meteorology). In addition, many regions in the ERA5
116 anomaly analysis show reduction in cloud fraction (Extended Data Fig. S5c), while ML-
117 MODIS (Fig. 2c) and MODIS (Extended Data Fig. S3c) show Atlantic-wide increases in cloud
118 fraction. We recognize that the CF from MODIS and LCC from ERA5 are derived from
119 satellite and model reanalysis, respectively, and are not entirely equivalent or directly
120 comparable; but again, this result suggests that aerosols are the primary driver of the increase
121 in CF over the study domain. Negligible LCC anomaly is found in September 2014 in ERA5

122 (not shown here), because SST anomaly is much weaker in September; however, a consistent
123 increase of CF in all latitude zonal means with an average of 0.02 is found in our ML-MODIS
124 results, which should be solely due to ACI (see Extended Data Fig. S9c).

125

126

127

128

Table S1 | The explanatory variables in random forest based machine-learning training*.

1	Temperature at 1000 hpa	Temperature at 950 hpa	Temperature at 900 hpa	Temperature at 850 hpa
2	Temperature at 800 hpa	Temperature at 750 hpa	Temperature at 700 hpa	Temperature at 650 hpa
3	Temperature at 600 hpa	Temperature at 550 hpa	Relative Humidity at 1000 hpa	Relative Humidity at 950 hpa
4	Relative Humidity at 900 hpa	Relative Humidity at 850 hpa	Relative Humidity at 800 hpa	Relative Humidity at 750 hpa
5	Relative Humidity at 700 hpa	Relative Humidity at 650 hpa	Relative Humidity at 600 hpa	Relative Humidity at 550 hpa
6	Potential Vorticity at 1000 hpa	Potential Vorticity at 950 hpa	Potential Vorticity at 900 hpa	Potential Vorticity at 850 hpa
7	Potential Vorticity at 800 hpa	Potential Vorticity at 750 hpa	Potential Vorticity at 700 hpa	Potential Vorticity at 650 hpa
8	Potential Vorticity at 600 hpa	Potential Vorticity at 550 hpa	Wind-U at 1000 hpa	Wind-U at 950 hpa
9	Wind-U at 900 hpa	Wind-U at 850 hpa	Wind-U at 800 hpa	Wind-U at 750 hpa
10	Wind-U at 700 hpa	Wind-U at 650 hpa	Wind-U at 600 hpa	Wind-U at 550 hpa
11	Wind-V at 1000 hpa	Wind-V at 950 hpa	Wind-V at 900 hpa	Wind-V at 850 hpa
12	Wind-V at 800 hpa	Wind-V at 750 hpa	Wind-V at 700 hpa	Wind-V at 650 hpa
13	Wind-V at 600 hpa	Wind-V at 550 hpa	Wind: updraft at 1000 hpa	Wind: updraft at 950 hpa
14	Wind: updraft at 900 hpa	Wind: updraft at 850 hpa	Wind: updraft at 800 hpa	Wind: updraft at 750 hpa
15	Wind: updraft at 700 hpa	Wind: updraft at 650 hpa	Wind: updraft at 600 hpa	Wind: updraft at 550 hpa
16	Vorticity at 1000 hpa	Vorticity at 950 hpa	Vorticity at 900 hpa	Vorticity at 850 hpa

17	Vorticity at 800 hpa	Vorticity at 750 hpa	Vorticity at 700 hpa	Vorticity at 650 hpa
18	Vorticity at 600 hpa	Vorticity at 550 hpa	Specific Humidity at 1000 hpa	Specific Humidity at 950 hpa
19	Specific Humidity at 900 hpa	Specific Humidity at 850 hpa	Specific Humidity at 800 hpa	Specific Humidity at 750 hpa
20	Specific Humidity at 700 hpa	Specific Humidity at 650 hpa	Specific Humidity at 600 hpa	Specific Humidity at 550 hpa
21	Geopotential at 1000 hpa	Geopotential at 950 hpa	Geopotential at 900 hpa	Geopotential at 850 hpa
22	Geopotential at 800 hpa	Geopotential at 750 hpa	Geopotential at 700 hpa	Geopotential at 650 hpa
23	Geopotential at 600 hpa	Geopotential at 550 hpa	Longitude	Latitude
24	Dew point at 2 meter	K-index	Wind gust at 10 meter	Instant moisture flux
25	Large-scale precipitation fraction	Large-scale precipitation	Precipitation type	Friction velocity
26	Wind speed at 10 meter	Wind-U at 100 meter	Wind-V at 100 meter	Sea-ice area fraction
27	Skin temperature	Total column water vapor	Convective available potential energy	Sea surface temperature
28	Mean sea level pressure	Large-scale rain rate	Total column rain water	Total precipitation
29	Boundary layer height	Trapping layer base height		

130 *All datasets are available from ECMWF ERA5 reanalysis.

131

132