


Partial convergence of the human vaginal and rectal maternal microbiota in late gestation and early post-partum

Journal Article

Author(s):

Shin, Hakdong; Martinez, Keith A.; Henderson, Nora; Jay, Melanie; Schweizer, William; Bogaert, Debby; Park, Gwoncheol; [Bokulich, Nicholas](#) ; Blaser, Martin J.; Dominguez-Bello, Maria Gloria

Publication date:

2023-06-13

Permanent link:

<https://doi.org/10.3929/ethz-b-000618149>

Rights / license:

[Creative Commons Attribution 4.0 International](#)

Originally published in:

npj Biofilms and Microbiomes 9(1), <https://doi.org/10.1038/s41522-023-00404-5>

ARTICLE OPEN



Partial convergence of the human vaginal and rectal maternal microbiota in late gestation and early post-partum

Hakdong Shin¹, Keith A. Martinez^{1,2,3}, Nora Henderson², Melanie Jay^{2,4}, William Schweizer⁵, Debby Bogaert⁶, Gwoncheol Park¹, Nicholas A. Bokulich⁷, Martin J. Blaser⁸ and Maria Gloria Dominguez-Bello^{3,9}✉

The human vaginal and fecal microbiota change during pregnancy. Because of the proximity of these perineal sites and the evolutionarily conserved maternal-to-neonatal transmission of the microbiota, we hypothesized that the microbiota of these two sites (rectal and vaginal) converge during the last gestational trimester as part of the preparation for parturition. To test this hypothesis, we analyzed 16S *rRNA* sequences from vaginal introitus and rectal samples in 41 women at gestational ages 6 and 8 months, and at 2 months post-partum. The results show that the human vaginal and rectal bacterial microbiota converged during the last gestational trimester and into the 2nd month after birth, with a significant decrease in *Lactobacillus* species in both sites, as alpha diversity progressively increased in the vagina and decreased in the rectum. The microbiota convergence of the maternal vaginal-anal sites perinatally might hold significance for the inter-generational transmission of the maternal microbiota.

npj Biofilms and Microbiomes (2023)9:37; <https://doi.org/10.1038/s41522-023-00404-5>

INTRODUCTION

The maternal vaginal microbiome is a primordial source of microbes for the developing newborn^{1–3}, carrying microbiotas from feces and other body sites⁴, providing seeding for multiple body sites in the baby, and protection against pathogens after birth⁵. For the mother, it remains important for health during and after birth⁶. Women undergo important physiological changes during gestation that also affect their microbiomes^{7,8}. Vaginal *Lactobacillus* increase relative abundances during pregnancy and this is observed across American⁹, European¹⁰, African¹¹, and Asian¹² populations. The fecal microbiota also changes with pregnancy, with reduced richness, and increased abundances of Actinobacteria (including Bifidobacteriales) and Proteobacteria⁷. Third trimester fecal microbiota induce increased pro-inflammatory responses in germ-free mice compared with those from the first trimester⁷. The vaginal microbiota may have increased diversity post-partum, driven in part by increases in species of anaerobes (*Peptoniphilus*, *Prevotella*, and *Anaerococcus*), and a reduction in *Lactobacillus* species⁹.

Since in addition to the vagina, babies are exposed at birth to the mother's perineum, which is a potential connection between the microbiomes of the intestine and the vagina, both of which undergo gestational changes. We hypothesized that the rectal and vaginal microbiotas converge during the last gestational trimester. This would be consistent with the conserved transformation of the maternal microbiota for transmission to the next generation. In this work, we determined the ante- and post-partum vaginal and rectal bacterial community structure in 21 mothers who delivered vaginally and in 20 mothers who delivered by Cesarean section (C-section), whose samples were obtained as part of an earlier study^{13,14}.

RESULTS

Comparison of vaginal and rectal communities during pregnancy and postpartum

We compared vaginal and rectal communities early (month 6) and late (month 8) in the third gestational trimester, and at 2 months post-partum, in 41 mothers (Supplementary Table 1 and Supplementary Table 2). A total of ~3 million V4 16S *rRNA* gene sequences with an average of ~20,000 sequences per sample were obtained (Supplementary Table 3). Bacterial alpha diversity in the rectum gradually declined over the last gestational trimester and continued to decrease into the second month postpartum. In contrast, there was a postpartum increase in phylogenetic diversity of vaginal microbiota (Fig. 1A), with increasing tendencies in either richness or evenness (Supplementary Fig. 1).

More bacterial taxa differed significantly in relative abundance between vagina and rectum during the prenatal period (52 taxa) than in the postpartum period (45 taxa; Fig. 2A and Supplementary Fig. 2), consistent with convergence of community structure. Similarly, community distances between the rectal and vaginal microbiotas became progressively reduced from the last gestational trimester to post-partum month 2 (Figs. 1B, C). Paired bootstrapping analysis indicated that postpartum convergence was more pronounced in the vagina than in the rectum (Figs. 1D, 2, and Supplementary Fig. 3).

Temporal changes from prenatal to post-natal (Fig. 2) included reductions of *Lactobacillus* species relative abundances in both rectal and vaginal sites (Fig. 1D and Supplementary Fig. 4), and increases in Bacteroidetes including *Prevotella*, and in Firmicutes including *Streptococcus*, *Anaerococcus*, *Peptoniphilus*, and *Dialister* in the vaginal site (Fig. 2 and Supplementary Fig. 3). The predicted species for *Lactobacillus* features (DADA2-based) identified in this

¹Department of Food Science & Biotechnology, and Carbohydrate Bioproduct Research Center, College of Life Science, Sejong University, Seoul, South Korea. ²Department of Medicine, New York University Langone Medical Center, New York, NY, USA. ³Department of Biochemistry and Microbiology, Rutgers University, New Brunswick, NJ, USA. ⁴Department of Population Health, New York University Langone Medical Center, New York, NY, USA. ⁵Department of Obstetrics and Gynecology, New York University Langone Medical Center, New York, NY, USA. ⁶MRC Centre for Inflammation Research, University of Edinburgh, Edinburgh, Scotland. ⁷Laboratory of Food Systems Biotechnology, Institute of Food, Nutrition and Health, ETH Zürich, Zürich, Switzerland. ⁸Center for Advanced Biotechnology and Medicine, Rutgers University, New Brunswick, NJ, USA. ⁹Department of Anthropology, Rutgers University, New Brunswick, NJ, USA. ✉email: mg.dominguez-bello@rutgers.edu




Fig. 1 Diversity in maternal pre- and post-partum rectal and vaginal microbiomes in analyses of 16 S rRNA sequences. **A** Phylogenetic diversity in each site at three time-points. Labeled means without a common letter differ significantly, p -value < 0.05 . Center line, median; quartile lines, upper and lower quartiles; violin plot, 1.5x interquartile range. **B** UniFrac distances between rectal and vaginal microbiomes at three perinatal time-points. Error bars represent mean \pm SEM. **C** PCoA of unweighted UniFrac distances between rectal and vaginal microbiomes at each time-point (PERMANOVA p -value < 0.001). **D** Taxa plots of maternal rectal (left panel) and vaginal (right panel) microbiomes at three peripartum time-points; *indicates LDA score > 3.0 between $-1M$ and $+2M$.




Fig. 2 Differences in the bacterial taxa of maternal pre- and post-partum rectal and vaginal sites, based on 16 S rRNA high throughput sequencing. A LefSe analysis of rectal versus vaginal sites late in gestation (left panel) or post-partum (right panel). **B** LefSe analysis of prenatal versus post-partum bacterial taxa in rectal (left panel) or vaginal (right panel) sites.

dataset were compared between vaginal and rectal microbiota. *Lactobacillus* features related to *L. iners*, *L. crispatus*, and *L. acidophilus* showed major components in shared microbial composition communities between the rectal and vaginal microbiotas. These temporal changes were more pronounced in vagina than in rectum, as evidenced by showing less error rates in supervised classification of vaginal samples by time-points, in relation to rectal samples (Supplementary Fig. 5 and Supplementary Fig. 6). The pre- or peripartum antibiotics exposure showed an equivalent tendency on vaginal/rectal microbial convergence (Supplementary Fig. 7).

DISCUSSION

Mammals evolved separate canals for reproduction, urination, and defecation but the rectal and vaginal orifices are typically proximal, while the urethra is not¹⁵. Because of their anatomical proximity and their perineal connectivity, we asked whether the two sites share bacteria during pregnancy. Our results show a clear convergence of the rectal and vaginal microbiota in the third trimester that extends post-partum. These results are consistent with independent reports of gestational changes in the vaginal^{10–12,16} and gut^{7,9} microbiota that were assessed separately, in different women. Although the vaginal microbiome varies with ethnicity, the gestational reduction in diversity abrogates these differences^{17,18}. In the current study, we compared vaginal and rectal gestational changes longitudinally, in the same mothers, with consistent and robust changes, regardless of delivery mode. Microbial convergence found in this study reflects the increase in shared microbial composition communities, leading to higher community similarity, as evidenced by the reduced community distances between the rectal and vaginal microbiotas. Indeed, the convergence continued after birth, to post-partum month 2 (Fig. 1B–D) as shown by the results. Since rectal samples did not change in diversity, this convergence is not an artifact due to simply shallower sampling. The vaginal ecosystem converged towards the rectal one. We speculate that these higher similarities in vaginal/rectal microbial composition communities could lead to selection in favor of (i) a *Lactobacillus*-enriched vaginal microbiota that maximally protects the gravid uterus from invasive pathogens¹⁹, and (ii) mechanisms that will expose the neonate to the widest diversity of intestinal microbes to be transferred from the mother². These biological roles of vaginal/rectal microbial convergence should be examined in further research. Limitations of this study included a small number of samples and a lack of control for the use of antibiotics during labor. Further studies should follow the maternal microbiota beyond 2 months post-partum and when it normalizes to the non-pregnant typical state.

Prior studies have shown that there is a large increase in beta-diversity in the maternal fecal microbiota in the third trimester, as the fecal microbial populations become more host-specific⁷. The post-labor vaginal microbiota differs from the baseline non-pregnant state, and the gestational changes have been shown to persist for up to 1 year after birth⁹.

Studies of mother-to-infant strain transmission show that the early infant gut microbiome contains maternal fecal bacteria²⁰, further supporting our findings of vaginal/rectal microbiota convergence. Indeed, proof of concept of gut microbiota restoration in infants born by C-section have been shown using both vaginal²¹ and fecal²² maternal sources. In any case, elective C-sections also have the confounding effect of antibiotics for the procedure, and will impair the post-partum observations. Although different body sites of the baby during birth are exposed to the same maternal microbiome, baby site selection effect is observable at day 2 after birth, in the gut, skin and oral microbiota²¹, as the developmental succession continues after the infant's birth¹⁴. Gestational changes in the microbiota might also be relevant to pregnancy outcomes⁹, since the fetus receives

products of the metabolism of the maternal microbiota^{23,24}, although the direction of causality is not well-understood.

Although the pre- and post-partum changes in these for infant health remains unknown. We used DADA2-based methods to predict *Lactobacillus* species related to shared microbial composition communities between the rectal and vaginal microbiotas, and found that the major species were *L. iners*, *L. acidophilus*, and *L. crispatus*. *L. iners* has been associated with vaginosis²⁵, and *L. crispatus* and *L. acidophilus* are associated with the high *Lactobacillus* dominance vaginal profile in healthy women²⁶. If these changes optimize the exposure of infants to the beneficial maternal microbiota, they are of adaptive value. More research is needed to understand both the dynamics and functional significance of the maternal microbiota during the critical peripartum period of maternal recovery and child development and across wider racial demographics.

METHODS

Sample and sequence information

In this study, we prospectively collected samples from the vaginal introitus and rectum of 41 women during pregnancy and at 2 months postpartum, as reported¹³. Vaginal -introitus- swabs were collected by the obstetrician and/or through self-collection by the mothers using sterile cotton-tipped swabs, a method shown to be reliable to quantify BV-associated bacteria²⁷ and to identify vaginal bacterial structures²⁸. For this analysis, we used rectal ($n = 73$) or vaginal ($n = 75$) samples from pregnant mothers in the early (weeks 23–35) or late (weeks 36–delivery) third gestational trimester and 2 months after delivery (Supplementary Table 1 and Supplementary Fig. 7). DNA had been extracted and the 16S *rRNA* V4 gene region had been sequenced using the Illumina MiSeq platform, as reported¹³ (European Nucleotide Archive: PRJEB14529). The samples analyzed in this study consist of a subset of samples collected with informed consent according to a New York University Institutional Review Board-approved study that took place in New York City from 2011 to 2014. Participation was voluntary and included written informed consent.

Data analysis

The 16S *rRNA* gene sequencing dataset, bacterial taxonomic compositions, and analyses of alpha-diversity (phylogenetic diversity²⁹, observed ASVs, and ASV evenness) and beta-diversity (Unweighted/Weighted UniFrac distance³⁰) were performed using QIIME 2 version 2020.6³¹ and its associated plugins. The q2-demux plugin was used for the demultiplexing and quality filtering of raw sequencing reads. Qualified reads were trimmed and denoised with DADA2³². All amplicon sequence variants (ASVs) were aligned using MAFFT³³ and used to generate a rooted phylogenetic tree with FastTree 2³⁴. The q2-feature-classifier plugin³⁵ was used to trim the 99% SILVA 16S *rRNA* gene database³⁶ to the 515F–806R (V4) region, train a naïve Bayes taxonomy classifier on these sequences, and use it to taxonomically classify each ASV. For comparisons of bacterial diversity, all communities were rarefied to 4347 reads (lowest number of reads) per sample, to include all samples the dataset. To determine significant differences in diversity, the Kruskal–Wallis test was used as a non-parametric test. Linear discriminant analysis (LDA) effect size (LEfSe) was used to detect statistically significant differences in the representation of bacterial taxa in comparisons (LDA score >3.0)³⁷. Also, ALDEx2 tools³⁸, which uses clr-transformed data generated from 128 Monte Carlo instances, used to confirm the LEfSe results. The Welch's *t*-test was used followed by Benjamini–Hochberg false discovery rate (FDR) correction, and effect size was calculated. To evaluate community separation in beta diversity, PERMANOVA

(with 999 random permutations)³⁰ was used. Supervised classification of vaginal and rectal samples by time-points based on the ASVs table collapsed to genus level was performed using a q2-sample-classifier³⁹ based on Random Forest classifier⁴⁰ and nested stratified 5-fold cross-validation.

Reporting summary

Further information on research design is available in the Nature Research Reporting Summary linked to this article.

DATA AVAILABILITY

The sequence data have been deposited in the European Nucleotide Archive under accession number ERP016173.

Received: 2 May 2021; Accepted: 2 June 2023;

Published online: 13 June 2023

REFERENCES

- Palmer, C., Bik, E. M., DiGiulio, D. B., Relman, D. A. & Brown, P. O. Development of the human infant intestinal microbiota. *PLoS Biol.* **5**, e177 (2007).
- Dominguez-Bello, M. G. et al. Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc. Natl Acad. Sci. USA* **107**, 11971–11975 (2010).
- Wampach, L. et al. Birth mode is associated with earliest strain-conferred gut microbiome functions and immunostimulatory potential. *Nat. Commun.* **9**, 5091 (2018).
- Song, S. J. et al. Naturalization of the microbiota developmental trajectory of Cesarean-born neonates after vaginal seeding. *Med* **2**, 951–964.e955 (2021).
- Shao, Y. et al. Stunted microbiota and opportunistic pathogen colonization in caesarean-section birth. *Nature* **574**, 117–121 (2019).
- Gajer, P. et al. Temporal dynamics of the human vaginal microbiota. *Sci. Transl. Med* **4**, 132ra152 (2012).
- Koren, O. et al. Host remodeling of the gut microbiome and metabolic changes during pregnancy. *Cell* **150**, 470–480 (2012).
- Romero, R. et al. The composition and stability of the vaginal microbiota of normal pregnant women is different from that of non-pregnant women. *Microbiome* **2**, 4 (2014).
- DiGiulio, D. B. et al. Temporal and spatial variation of the human microbiota during pregnancy. *Proc. Natl Acad. Sci. USA* **112**, 11060–11065 (2015).
- MacIntyre, D. A. et al. The vaginal microbiome during pregnancy and the postpartum period in a European population. *Sci. Rep.* **5**, 8988 (2015).
- Bisanz, J. E. et al. Microbiota at multiple body sites during pregnancy in a rural Tanzanian population and effects of Moringa-supplemented probiotic yogurt. *Appl Environ. Microbiol* **81**, 4965–4975 (2015).
- Huang, Y. E. et al. Homogeneity of the vaginal microbiome at the cervix, posterior fornix, and vaginal canal in pregnant Chinese women. *Micro. Ecol.* **69**, 407–414 (2015).
- Bokulich, N. A. et al. Antibiotics, birth mode, and diet shape microbiome maturation during early life. *Sci. Transl. Med* **8**, 343ra382 (2016).
- Koenig, J. E. et al. Succession of microbial consortia in the developing infant gut microbiome. *Proc. Natl Acad. Sci. USA* **108**, 4578–4585 (2011).
- Standring, S. *Gray's Anatomy: the Anatomical Basis of Clinical Practice*. 1261–1266 (Elsevier, 2016).
- Romero, R. et al. The vaginal microbiota of pregnant women who subsequently have spontaneous preterm labor and delivery and those with a normal delivery at term. *Microbiome* **2**, 18 (2014).
- Serrano, M. G. et al. Racioethnic diversity in the dynamics of the vaginal microbiome during pregnancy. *Nat. Med* **25**, 1001–1011 (2019).
- Dominguez-Bello, M. G. Gestational shaping of the maternal vaginal microbiome. *Nat. Med* **25**, 882–883 (2019).
- Borgdorff, H. et al. Unique insights in the cervicovaginal *Lactobacillus iners* and *L. crispatus* proteomes and their associations with microbiota dysbiosis. *PLoS ONE* **11**, e0150767 (2016).
- Ferretti, P. et al. Mother-to-infant microbial transmission from different body sites shapes the developing infant gut microbiome. *Cell Host Microbe* **24**, 133–145.e135 (2018).
- Dominguez-Bello, M. G. et al. Partial restoration of the microbiota of cesarean-born infants via vaginal microbial transfer. *Nat. Med.* **22**, 250–253 (2016).

- Korpela, K. et al. Maternal fecal microbiota transplantation in cesarean-born infants rapidly restores normal gut microbial development: a proof-of-concept study. *Cell*, <https://doi.org/10.1016/j.cell.2020.08.047> (2020).
- Romano-Keeler, J. & Weitkamp, J. H. Maternal influences on fetal microbial colonization and immune development. *Pediatr. Res* **77**, 189–195 (2015).
- Hemberg, E. et al. Occurrence of bacteria and polymorphonuclear leukocytes in fetal compartments at parturition; relationships with foal and mare health in the peripartum period. *Theriogenology* **84**, 163–169 (2015).
- Verstraelen, H. et al. Longitudinal analysis of the vaginal microflora in pregnancy suggests that *L. crispatus* promotes the stability of the normal vaginal microflora and that *L. gasseri* and/or *L. iners* are more conducive to the occurrence of abnormal vaginal microflora. *BMC Microbiol.* **9**, 116 (2009).
- Zheng, N., Guo, R., Wang, J., Zhou, W. & Ling, Z. Contribution of *Lactobacillus iners* to vaginal health and diseases: a systematic review. *Front. Cell. Infect. Microbiol.* **11**, 792787 (2021).
- Nelson, D. B., Bellamy, S., Gray, T. S. & Nachamkin, I. Self-collected versus provider-collected vaginal swabs for the diagnosis of bacterial vaginosis: an assessment of validity and reliability. *J. Clin. Epidemiol.* **56**, 862–866 (2003).
- Forney, L. J. et al. Comparison of self-collected and physician-collected vaginal swabs for microbiome analysis. *J. Clin. Microbiol* **48**, 1741–1748 (2010).
- Faith, D. P. Conservation evaluation and phylogenetic diversity. *Biol. Conserv.* **61**, 1–10 (1992).
- Anderson, M. A new method for non-parametric multivariate analysis of variance. *Aust. Ecol.* **26**, 32–46 (2001).
- Bolyen, E. et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).
- Callahan, B. J. et al. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* **13**, 581–583 (2016).
- Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066 (2002).
- Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2-approximately maximum-likelihood trees for large alignments. *PLoS ONE* **5**, e9490 (2010).
- Bokulich, N. A. et al. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* **6**, 90 (2018).
- Yilmaz, P. et al. The SILVA and “All-species Living Tree Project (LTP)” taxonomic frameworks. *Nucleic Acids Res.* **42**, D643–D648 (2014).
- NonSegata, N. et al. Metagenomic biomarker discovery and explanation. *Genome Biol.* **12**, R60 (2011).
- Fernandes, A. D., Macklaim, J. M., Linn, T. G., Reid, G. & Gloor, G. B. ANOVA-like differential expression (ALDEx) analysis for mixed population RNA-Seq. *PLoS One* **8**, e67019 (2013).
- Bokulich, N. A. et al. q2-sample-classifier: machine-learning tools for microbiome classification and regression. *J. Open Res. Softw.* **3**, <https://doi.org/10.21105/joss.00934> (2018).
- Breiman, L. Random forests. *Mach. Learn.* **45**, 5–32 (2001).

ACKNOWLEDGEMENTS

We acknowledge the financial support from EMCH Fund, the C&D Fund (to MGDB), T32 AI007180 and R01 DK090989 from the National Institutes of Health, and from the C & D and Zlinkoff Funds, and the TransAtlantic program of the Fondation Leducq (to M.J.B.). The NYULMC Genome Technology Core was partially supported by the Cancer Center Support Grant, P30CA016087. H.S. was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2021R1A2C1095215 and NRF-2022R1A6A1A03055869). We acknowledge the collaboration of Arnon D. Leiber, Fritz Francois, Sukhleen Bedi, and Guillermo I. Perez-Perez. MGDB is a fellow of the Canadian Institute for Advanced Research (CIFAR).

AUTHOR CONTRIBUTIONS

M.G.D.B. and H.S. designed the study. N.H., M.J., and W.S. completed sample collection and NH performed sample processing. H.S., N.A.B., G.P., K.A.M. II, M.J.B., and M.G.D.B. analyzed the data. All authors reviewed the data and manuscript drafts, and H.S., K.A.M. II, D.B., M.J.B., and M.G.D.B. interpreted results and wrote the final manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41522-023-00404-5>.

Correspondence and requests for materials should be addressed to Maria Gloria Dominguez-Bello.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023