

A special issue preface: Radiocarbon in the Anthropocene

Other Journal Item**Author(s):**

Eglinton, Timothy I.; Graven, Heather D.; Raymond, Peter A.; Trumbore, Susan E.

Publication date:

2023-11-27

Permanent link:

<https://doi.org/10.3929/ethz-b-000638607>

Rights / license:

[Creative Commons Attribution 4.0 International](#)

Originally published in:

Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences 381(2261), <https://doi.org/10.1098/rsta.2022.0209>

Funding acknowledgement:

193770 - Radiocarbon Inventories of Switzerland (RICH): An integrated approach to understand the changing carbon cycle (SNF)

Preface



Cite this article: Eglinton TI, Graven HD, Raymond PA, Trumbore SE. 2023 A special issue preface: Radiocarbon in the Anthropocene.

Phil. Trans. R. Soc. A **381**: 20220209.

<https://doi.org/10.1098/rsta.2022.0209>

Received: 25 August 2023

Accepted: 25 August 2023

One contribution of 10 to a Theo Murphy meeting issue 'Radiocarbon in the Anthropocene'.

Subject Areas:

atmospheric chemistry, environmental chemistry, atmospheric science, climatology, geochemistry

Keywords:

radiocarbon (^{14}C), carbon cycle, climate change, Anthropocene, fossil fuels, bomb ^{14}C

Author for correspondence:

Timothy I. Eglinton

e-mail: timothy.eglinton@erdw.ethz.ch

A special issue preface: Radiocarbon in the Anthropocene

Timothy I. Eglinton¹, Heather D. Graven², Peter A. Raymond³ and Susan E. Trumbore⁴

¹Department of Earth Sciences, ETH Zurich, Zurich 8092, Switzerland

²Department of Physics, Imperial College London, London, UK

³School of the Environment, Yale University, New Haven, CT, USA

⁴Department of Biogeochemical Processes, Max Planck Institute for Biogeochemistry, Jena 07745, Germany

TIE, 0000-0001-5060-2155

The Anthropocene is defined by marked acceleration in human-induced perturbations to the Earth system. Anthropogenic emissions of CO_2 and other greenhouse gases to the atmosphere and attendant changes to the global carbon cycle are among the most profound and pervasive of these perturbations. Determining the magnitude, nature and pace of these carbon cycle changes is crucial for understanding the future climate that ecosystems and humanity will experience and need to respond to. This special issue illustrates the value of radiocarbon as a tool to shed important light on the nature, magnitude and pace of carbon cycle change.

This article is part of the Theo Murphy meeting issue 'Radiocarbon in the Anthropocene'.

1. Introduction

Radiocarbon, the radioactive isotope of carbon, serves as a powerful tool to examine carbon cycle processes over a range of timescales by virtue of two phenomena: (i) its quasi-constant production in the atmosphere and decay with an approximately 5700-year half-life and (ii) the abrupt spike of 'bomb' ^{14}C introduced into the atmosphere in the mid twentieth century

© 2023 The Authors. Published by the Royal Society under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0/>, which permits unrestricted use, provided the original author and source are credited.

as a consequence of the initiation and cessation of above-ground nuclear weapons testing. Together, these phenomena allow tracking, apportionment and assessment of carbon exchange and turnover times between and within Earth's carbon reservoirs. Presently, the radiocarbon content of the atmosphere is returning to pre-industrial levels as a consequence of the redistribution of bomb ^{14}C between atmospheric, oceanic and terrestrial reservoirs, as well as on-going emissions of ^{14}C -depleted carbon from fossil fuel burning and land-use change. Future variations of ^{14}C within and between different carbon reservoirs will depend on how trajectories of Anthropocene change evolve in the context of greenhouse gas emission scenarios and the efficacy of carbon reduction strategies.

With a backdrop of on-going environmental change, together with an appreciation of the unique value of ^{14}C as a tool to understand the changing carbon cycle, a Royal Society-sponsored meeting on *Radiocarbon in the Anthropocene* was convened on 16 and 17 May 2022 at Whittlebury Park, UK, in which carbon cycle scientists and radiocarbon experts discussed current research related to the application of ^{14}C to understand natural carbon cycle processes and to assess past and future changes. The theme issue includes papers stemming from presentations at this meeting, augmented by others that reflect current frontiers in radiocarbon research.

2. Contents of the theme issue

The first paper of the theme issue, an Opinion Piece by Eglinton *et al.* [1], seeks to highlight the value of ^{14}C for understanding Anthropocene carbon cycle change, and to propose an intense phase of coordinated and sustained research activity that harnesses the full potential of radiocarbon.

Three papers describe applications of radiocarbon to assess soil carbon dynamics, and the factors controlling soil carbon turnover. The paper by Stoner *et al.* [2] describes the application of serial or ramped thermal oxidation of subsoils developed on different bedrock lithologies in order to examine the influence of mineralogy on organic matter stabilization and carbon quality. The products released were examined for molecular and isotopic (radiocarbon) composition and in the context of mineral composition, particularly pedogenic oxides and clays. Their findings provide strong evidence for mineralogical control on the age distribution, chemical diversity and persistence of soil organic matter, with implications for modelling soil C turnover, and for predictions of soil carbon vulnerability in the face of changing environmental conditions. Sierra *et al.* [3] present a modelling study that seeks to describe the transit time of carbon through the terrestrial biosphere, from photosynthetic carbon fixation to respiration. Variables that are explored include the impact of increasing productivity and respiration. Model results indicate that temperature-driven increases in respiration are fuelled by the degradation of older carbon pools, resulting in an increase in transit time, whereas the opposite holds for increased primary productivity. Simulations using different terrestrial carbon models indicate overall shorter transit times prevail due to the dominance of temperature-driven increases in productivity over the twentieth century. However, increased asymmetry in transit time distributions reflects the differing carbon dynamics of low- versus high-latitude ecosystems, with decreasing transit times in the former and increasing transit times in the latter due to the respiration of long-stored carbon. In an observational study that relates to the findings of Sierra *et al.*, Schuur *et al.* [4] describe results from almost 20 years of time-series CO_2 flux and radiocarbon measurements at a permafrost site in Alaska where carbon has been stored in frozen and waterlogged soils that shed light on long-term ecosystem and carbon cycle perturbations in a region of the world that is experiencing the accelerated climate change. The detection of respired CO_2 with depleted ^{14}C values relative to atmospheric CO_2 provides evidence of old permafrost soil C degradation, and a steeper trajectory in respired $^{14}\text{CO}_2$ decrease relative to the atmosphere over the time series indicates that this degradation is enhanced with regional climate change, with temperature and moisture being the key variables. These findings suggest that changing environmental conditions are favouring release of old, previously stabilized carbon, representing a positive feedback with respect to on-going climate change.

Two manuscripts focus on the use of radiocarbon to examine controls on carbon dynamics during mobilization and transport along the aquatic continuum. Rhyner *et al.* [5] examined radiocarbon signatures of dissolved and particulate carbon phases in a suite of 21 Swiss rivers sampled during an interval of elevated water and sediment discharge that span a broad range of watershed characteristics. They found that mean river basin elevation was most strongly linked to ^{14}C values, suggesting that the dynamics of carbon mobilized and exported by rivers is subject to significant ecoregional control. Gies *et al.* [6] use radiocarbon measurements of different groups of source-specific ‘biomarker’ compounds isolated from river sediments to examine relationships with watershed properties and carbon dynamics within river basins. They find coherence between ^{14}C characteristics of terrestrial plant biomarkers and markers of soil microbial residues, suggesting that biospheric carbon export from watersheds is modulated by soil storage.

Two papers address ^{14}C measurements of atmospheric CO_2 in the context of constraining local to regional fossil fuel CO_2 sources and emissions. Maier *et al.* [7] discuss the complexities in constraining fossil fuel CO_2 surplus (ΔffCO_2) emissions at a regional scale through the conventional approach of comparing local $^{14}\text{CO}_2$ measurements to measurements at a background site. Specific challenges, biases and uncertainties are discussed in the context of the Integrated Carbon Observation System (ICOS) atmospheric network for Europe and the background station at Mace Head, Ireland. They highlight the significant biases that can be incurred in ΔffCO_2 estimates as a consequence of the small signals as well as the impacts of nuclear contamination and heterotrophic respiration. Young *et al.* [8] describe the use of ^{14}C as part of urban flask measurements from Auckland New Zealand that form part of a programme to assess local ffCO_2 as well as excess carbon monoxide emissions relative to background. They determine that excess CO largely results from traffic emissions, and this information is used to develop a revised inventory for excess CO .

The final paper in the theme issue by Bard *et al.* [9] reports annual-resolution ^{14}C measurements on tree rings from subfossil Scots Pines from the Southern French Alps in the context of past atmospheric ^{14}C variability spanning a 700-year interval during the Late Glacial period. The record reveals two prominent events that are attributed to solar activity, providing further linkages with Greenland ice core records and temporal context for interpretation of past climate variability.

Data accessibility. This article has no additional data.

Declaration of AI use. We have not used AI-assisted technologies in creating this article.

Authors’ contributions. T.I.E.: writing—original draft, writing—review and editing; H.D.G.: writing—original draft, writing—review and editing; S.E.T.: writing—original draft, writing—review and editing; P.A.R.: writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. This theme issue was put together by the Guest Editor team under supervision from the journal’s Editorial staff, following the Royal Society’s ethical codes and best-practice guidelines. The Guest Editor team invited contributions and handled the review process. Individual Guest Editors were not involved in assessing papers where they had a personal, professional, or financial conflict of interest with the authors, or the research described. Independent reviewers assessed all papers.

Funding. No financial support was received for this article.

Acknowledgements. The Guest Editors wish to thank all the contributing authors for their hard work in preparing their manuscripts, and in meeting deadlines, and their patience while this theme issue has been prepared. We also thank the referees for their constructive feedback on the articles. We are indebted to the staff of the *Phil. Trans. R. Soc. A* Editorial Office, and wish to extend our special thanks to Alice Power, the Commissioning Editor, for her efficiency and patience in guiding us through the editorial process.

References

1. Eglinton TI *et al.* 2023 Making the case for an International Decade of Radiocarbon. *Phil. Trans. R. Soc. A* **381**, 20230081. (doi:10.1098/rsta.2023.0081)

2. Stoner S, Trumbore SE, González-Pérez JA, Schrumpp M, Sierra CA, Hoyt AM, Chadwick O, Doetterl S 2023 Relating mineral–organic matter stabilization mechanisms to carbon quality and age distributions using ramped thermal analysis. *Phil. Trans. R. Soc. A* **381**, 20230139. (doi:10.1098/rsta.2023.0139)
3. Sierra CA, Quetin GR, Metzler H, Müller M. 2023 A decrease in the age of respired carbon from the terrestrial biosphere and increase in the asymmetry of its distribution. *Phil. Trans. R. Soc. A* **381**, 20220200. (doi:10.1098/rsta.2022.0200)
4. Schuur EAG, Hicks Pries C, Mauritz M, Pegoraro E, Rodenhizer H, See C, Ebert C. 2023 Ecosystem and soil respiration radiocarbon detects old carbon release as a fingerprint of warming and permafrost destabilization with climate change. *Phil. Trans. R. Soc. A* **381**, 20220201. (doi:10.1098/rsta.2022.0201)
5. Rhyner TMY *et al.* 2023 Radiocarbon signatures of carbon phases exported by Swiss rivers in the Anthropocene. *Phil. Trans. R. Soc. A* **381**, 20220326. (doi:10.1098/rsta.2022.0326)
6. Gies H, Lupker M, Galy V, Hemingway J, Boehman B, Schwab M, Haghypour N, Eglinton TI. 2023 Multi-molecular ^{14}C evidence for mineral control on terrestrial carbon storage and export. *Phil. Trans. R. Soc. A* **381**, 20220328. (doi:10.1098/rsta.2022.0328)
7. Maier F, Levin I, Gachkivskyi M, Rödenbeck C, Hammer S. 2023 Estimating regional fossil fuel CO_2 concentrations from $^{14}\text{CO}_2$ observations: challenges and uncertainties. *Phil. Trans. R. Soc. A* **381**, 20220203. (doi:10.1098/rsta.2022.0203)
8. Young HA *et al.* 2023 Urban flask measurements of CO_2^{ff} and CO to identify emission sources at different site types in Auckland, New Zealand. *Phil. Trans. R. Soc. A* **381**, 20220204. (doi:10.1098/rsta.2022.0204)
9. Bard E, Miramont C, Capano M, Guibal F, Marschal C, Rostek F, Tuna T, Fagault Y, Heaton TJ. 2023 A radiocarbon spike at 14 300 cal yr BP in subfossil trees provides the impulse response function of the global carbon cycle during the Late Glacial. *Phil. Trans. R. Soc. A* **381**, 20220206. (doi:10.1098/rsta.2022.0206)