



Other Journal Item

Correction: Testing the potential of a virtual reality neurorehabilitation system during performance of observation, imagery and imitation of motor actions recorded by wireless functional near-infrared spectroscopy (fNIRS)

Author(s):

Holper, Lisa; Mühlemann, Thomas; Scholkmann, Felix; Eng, Kynan; Kiper, Daniel; Wolf, Martin

Publication Date:

2013-02

Permanent Link:

<https://doi.org/10.3929/ethz-b-000064553> →

Originally published in:

Journal of NeuroEngineering and Rehabilitation 10, <http://doi.org/10.1186/1743-0003-10-16> →

Rights / License:

[Creative Commons Attribution 2.0 Generic](#) →

This page was generated automatically upon download from the [ETH Zurich Research Collection](#). For more information please consult the [Terms of use](#).

CORRECTION

Open Access

Correction: Testing the potential of a virtual reality neurorehabilitation system during performance of observation, imagery and imitation of motor actions recorded by wireless functional near-infrared spectroscopy (fNIRS)

Lisa Holper^{1,2*}, Thomas Muehleemann^{1,3}, Felix Scholkmann¹, Kynan Eng², Daniel Kiper² and Martin Wolf¹

Correction

Following publication of our article [1], we realised that some of the statistical tests used were not appropriate. We have now conducted the appropriate statistical tests, and updated the relevant tables, figures five and six (Figures 1 and 2 here, respectively) and conclusions accordingly.

For both the unilateral and the bilateral groups, analyses were recalculated.

Intra-condition differences:

- original publication [1]: paired *t*-test using means per trial for unilateral and bilateral group
- update: one-way repeated measures ANOVA using means per subject for unilateral and bilateral group

Inter-condition differences:

- original publication [1]: one-way ANOVA using means per trial for unilateral and bilateral group

- update: one-way repeated measures ANOVA using means per subject for unilateral and bilateral group

Conclusions

For the unilateral group (Table 1 and Figure 1), no changes in significance levels were found in $\Delta[O_2Hb]$ signals. For the bilateral group (Table 2 and Figure 2), the main differences compared to the original publication are that the intra-condition differences ($[O_2Hb]_{rest}$ versus $[O_2Hb]_{stim}$) for the two conditions 'Observation Right' (O_R, $p = 0.077$) and 'Observation Left' (O_L, $p = 0.080$) recorded over the ipsilateral hemisphere do not reach significant level any more. Hence, the paragraphs discussing the intra-condition significances in those two conditions (sections *Observation, imagery and imitation* and *Bilateral oxygenation* of the *Discussion*) are only applicable for the contralateral hemisphere. Further, in both groups changes in significance levels were found for $\Delta[HHb]$. However, since the *Discussion* and *Conclusion* of the originally published article only focuses on the concentration changes found in $\Delta[O_2Hb]$, this aspect does not change these sections.

* Correspondence: holper@ini.phys.ethz.ch

¹Biomedical Optics Research Laboratory (BORL), Division of Neonatology, Department of Obstetrics and Gynecology, University Hospital Zurich, Frauenklinikstrasse 10, 8091, Zurich, Switzerland

²Institute of Neuroinformatics (INI), University of Zurich and ETH Zurich, Winterthurerstrasse 190, 8057, Zurich, Switzerland

Full list of author information is available at the end of the article




Figure 1 Unilateral group recorded over contralateral hemisphere: shown are the $\Delta[\text{O}_2\text{Hb}]$ amplitude changes with standard error of the mean (SEM) and statistical significances of repeated measures ANOVA.




Figure 2 Bilateral group recorded over contra- (dark gray) and ipsilateral (light gray) hemisphere: shown are the $\Delta[\text{O}_2\text{Hb}]$ amplitude changes with standard error of the mean (SEM) and statistical significances of repeated measures ANOVA.

Table 1 Unilateral group

Unilateral group [N = 15]	Observation	Motor imagery	Observation & motor imagery	Imitation
Left hemisphere (contralateral) ($\mu\text{mol/l} \pm \text{SD}$)				
Mean Δ [O_2Hb]	0.06953 \pm 0.1800	0.0833 \pm 0.1404	0.0460 \pm 0.2218	0.2309 \pm 0.3212
Mean Δ [HHb]	-0.0051 \pm 0.03855	0.0356 \pm 0.0771	-0.0089 \pm 0.0963	0.0079 \pm 0.0832
Intra-condition, ANOVA, repeated measures				
[O_2Hb] rest-stim	p = 0.157	p = 0.037*	p = 0.435	p = 0.015*
[HHb] rest-stim	p = 0.612	p = 0.097	p = 0.727	p = 0.717
Inter-condition ANOVA, repeated measures, post-hoc-tests, Bonferroni 0.05				
		Δ [HHb]	Δ [O_2Hb]	
	O – MI	p = 0.347	p = 1.000	
	O – O&MI	p = 1.000	p = 1.000	
	O – IM	p = 1.000	p = 0.286	
	MI – O&MI	p = 0.132	p = 1.000	
	MI – IM	p = 1.000	p = 0.622	
	O&MI – IM	p = 1.000	p = 0.321	
Main effect on condition		p = 0.253	p = 0.062	

(Top) Mean signal amplitudes ($\mu\text{mol/l} \pm \text{SD}$) of channels with significant concentration changes. Separate calculations for increases in [O_2Hb], decreases in [HHb] in response to the four conditions for each group. Numbers were rounded to four decimal places. (Middle) Intra-condition statistical significance of the mean changes between [O_2Hb]_{rest} and [O_2Hb]_{stim} and [HHb]_{rest} and [HHb]_{stim} repeated measures ANOVA. (Bottom) Inter-condition statistical significance of mean changes of Δ [O_2Hb] and Δ [HHb] between the four conditions using repeated measures ANOVA. Shown are post-hoc tests (with Bonferroni correction); significant values ($p \leq 0.05$) are highlighted by * (observation = O, motor imagery = MI, observation & motor imagery = O & MI, imitation = IM).

Table 2 Bilateral group

Bilateral group [N = 8]	Observation right	Observation left	Imitation right	Imitation left
Left hemisphere (contralateral) ($\mu\text{mol/l} \pm \text{SD}$)				
Mean Δ [O ₂ Hb]	0.1231 \pm 0.1506	0.1231 \pm 0.1507	0.3941 \pm 0.4598	0.3715 \pm 0.4289
Mean Δ [HHb]	-0.0056 \pm 0.0676	-0.0408 \pm 0.0915	0.0371 \pm 0.1131	0.0474 \pm 0.0665
Intra-condition, ANOVA, repeated measures				
[O ₂ Hb] rest-stim	p = 0.044*	p = 0.047*	p = 0.025*	p = 0.038*
[HHb] rest-stim	p = 0.821	p = 0.247	p = 0.384	p = 0.084
Inter-condition, ANOVA, repeated measures, post-hoc-tests, Bonferroni 0.05				
		Δ [HHb]	Δ [O ₂ Hb]	
	O_R - O_L	p = 1.000	p = 1.000	
	O_R - IM_R	p = 1.000	p = 0.519	
	O_R - IM_L	p = 1.000	p = 0.862	
	OL_ - IM_R	p = 0.227	p = 0.486	
	O_L - IM_L	p = 0.223	p = 0.777	
	IM_R - IM_L	p = 1.000	p = 1.000	
Main effect on condition				
		p = 0.072	p = 0.119	
Right hemisphere (ipsilateral) ($\mu\text{mol/l} \pm \text{SD}$)				
Mean Δ [O ₂ Hb]	0.1541 \pm 0.0735	0.1957 \pm 0.1957	0.4036 \pm 0.2097	1.3728 \pm 1.6143
Mean Δ [HHb]	-0.0113 \pm 0.0334	0.0068 \pm 0.0274	0.0235 \pm 0.0402	0.7016 \pm 1.9167
Intra-condition, ANOVA, repeated measures				
[O ₂ Hb] rest-stim	p = 0.077	p = 0.080	p = 0.002*	p = 0.032*
[HHb] rest-stim	p = 0.367	p = 0.502	p = 0.142	p = 0.335
Inter-condition, ANOVA, repeated measures, post-hoc-tests, Bonferroni 0.05				
		Δ [HHb]	Δ [O ₂ Hb]	
	O_R - O_L	p = 1.000	p = 1.000	
	O_R - IM_R	p = 0.445	p = 0.014*	
	O_R - IM_L	p = 1.000	p = 0.324	
	OL_ - IM_R	p = 1.000	p = 0.015*	
	O_L - IM_L	p = 1.000	p = 0.231	
	IM_R - IM_L	p = 1.000	p = 0.710	
Main effect on condition				
		p = 0.384	p = 0.008*	

(Top) Mean signal amplitudes ($\mu\text{mol/l} \pm \text{SD}$) of channels with significant concentration changes. Separate calculations for increases in [O₂Hb], decreases in [HHb] in response to the four conditions for each group. Numbers were rounded to four decimal places. (Middle) Intra-condition statistical significance of the mean change between [O₂Hb]_{rest} and [O₂Hb]_{stim} and [HHb]_{rest} and [HHb]_{stim} using repeated measures ANOVA. (Bottom) Inter-condition statistical significance of mean changes of Δ [O₂Hb] and Δ [HHb] between the four conditions using repeated measures ANOVA. Shown are post-hoc tests (with Bonferroni correction); significant values ($p \leq 0.05$) are highlighted by * (observation left = O_L, observation right = O_R, imitation left = IM_L, imitation right = IM_R).

Received: 1 February 2013 Accepted: 1 February 2013
 Published: 8 February 2013

Reference

- Holper L, et al: Testing the potential of a virtual reality neurorehabilitation system during performance of observation, imagery and imitation of motor actions recorded by wireless functional near-infrared spectroscopy (fNIRS). *J Neuroeng Rehabil* 2012, **7**:57.

doi:10.1186/1743-0003-10-16

Cite this article as: Holper et al.: Correction: Testing the potential of a virtual reality neurorehabilitation system during performance of observation, imagery and imitation of motor actions recorded by wireless functional near-infrared spectroscopy (fNIRS). *Journal of NeuroEngineering and Rehabilitation* 2013 **10**:16.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

