



Report

Sparse Quadrature Approach to Bayesian Inverse Problems

Author(s):

Schwab, Christoph; Schillings, Claudia

Publication Date:

2013-08

Permanent Link:

<https://doi.org/10.3929/ethz-a-010394235> →

Rights / License:

[In Copyright - Non-Commercial Use Permitted](#) →

This page was generated automatically upon download from the [ETH Zurich Research Collection](#). For more information please consult the [Terms of use](#).

Sparse Quadrature Approach to Bayesian Inverse Problems

Ch. Schwab and C. Schillings

Research Report No. 2013-27
August 2013

Seminar für Angewandte Mathematik
Eidgenössische Technische Hochschule
CH-8092 Zürich
Switzerland

Sparse Quadrature Approach to Bayesian Inverse Problems

CHRISTOPH SCHWAB

(joint work with Claudia Schillings)

We consider the parametric deterministic formulation of Bayesian inverse problems with distributed parameter uncertainty from infinite dimensional, separable Banach spaces X , with uniform prior probability measure on space X of all uncertainties. Under the assumption of given observation data δ subject to additive observation noise $\eta \sim N(0, \Gamma)$ with positive covariance Γ , an infinite-dimensional version of Bayes' formula has been shown in [14].

For problems with uncertain, distributed parameters $u \in X$ (which could be a diffusion coefficient, elastic moduli in solid mechanics, shape of the domain D of definition of the physical problem [1], kinetic parameters in stoichiometric models of reaction-systems in biological systems [4, 7], permeability in porous media or optimal control of uncertain systems [9]), we develop a practical, adaptive computational algorithm for the efficient approximation of the infinite-dimensional integrals with respect to the Bayesian posterior (conditional on given data δ) μ^δ which arise in Bayes' formula in [14].

The Bayesian posterior μ^δ is shown to admit a representation in terms of a (generalized) polynomial chaos expansion in the (countably many) coordinates y_j which parametrize the uncertainty. We prove that if the uncertain datum $u \in X$ admits the (norm-convergent in X) expansion

$$u = \langle u \rangle + \sum_{j \geq 1} y_j \psi_j(x)$$

with $|y_j| \leq 1$ and with $(\|\psi_j\|_X)_{j \geq 1} \in \ell^p(\mathbb{N})$ for some $0 < p < 1$, then the solution $q(u) = (A(u))^{-1}f$ of the parametric operator equation will depend holomorphically on the parameters y_j , with precise control of the domain of holomorphy [3, 8, 5, 1, 9]. In two-scale limits of homogenization theory, these domains are independent of physical scale parameters [8].

We prove, generalizing [13], that the holomorphic dependence on the parameters y_j of the forward solution of the above problems implies p -sparsity of the polynomial chaos expansion for the parametric forward solution $q(u) = (A(u))^{-1}f$ and also for the density function of derivative of the Bayesian posterior μ^δ with respect to the prior μ_0 , conditional on given data δ .

The proof of the p -sparsity in [12, 10] is based on verification of holomorphic dependence of the polynomial chaos representation for the density of the Bayesian posterior with respect to the prior, following the proofs in the linear, elliptic diffusion problems in [3, 13].

Based on this sparsity result, dimension independent convergence rates of best N -term approximations of the parametric forward map as well as of the parametric density of the Bayesian posterior with respect to uniform prior follows from Stechkin's lemma.

We propose a deterministic, adaptive algorithm inspired by [6] and analogous to the adaptive interpolation methods in [2] and the references there. The proposed algorithm determines iteratively, and depending on the observation data δ and the observation noise variance Γ a sequence of quadrature dimensions and quadrature orders.

Convergence rates for the adaptive Smolyak quadrature approximation are shown, computationally, to coincide with the best N term approximation rates of the Bayesian posterior density which, in turn [12, 10] depend only on the sparsity class (characterized in turn by the summability exponent $p \in (0, 1)$) of the uncertain distributed parameter $u \in X$.

Convergence rates are obtained in [12, 10] via monotone L^∞ N -term approximations of the posterior density from [2]. The resulting rates $1/p - 1$ are larger than the rate $1/2$ afforded by Monte-Carlo methods and their variants (notably MCMC) for $p < 2/3$ when stated in terms of the number N of (numerical) solutions of the forward problems which are necessary in the quadrature algorithm.

Applications with verified holomorphic dependence include high-dimensional parametric initial value problems [4], semilinear elliptic equations [5, 1] with uncertain differential operators, parabolic evolution problems with uncertain operators [1], elliptic multiscale problems with uncertain coefficients [8] and from biological systems sciences [7], as well as optimal control of uncertain systems [9], and problems with uncertain shape [1].

Numerical examples are presented for diffusion problems with uncertain diffusion coefficient from [12], [10], and for large, parametric systems of initial value problems from stoichiometric models for biological systems with mass-action kinetics from [4, 7]. Computational savings with respect to adaptivity in the forward solver are indicated. Here, we present numerical experiments based on the parametric, parabolic initial boundary-value problem

$$\begin{aligned} \partial_t q(t, x) - \operatorname{div}(u(x) \nabla q(t, x)) &= f(t, x) \quad (t, x) \in T \times D, \\ q(0, x) &= 0 \quad x \in D, \\ q(t, 0) = q(t, 1) &= 0 \quad t \in T, \end{aligned}$$

with $f(t, x) = 100 \cdot tx$, $D = (0, 1)$ and $T = (0, 1)$. The uncertain coefficient u is parametrized as $u(x, y) = \bar{a} + \sum_{j=1}^{128} y_j \psi_j$, where $\bar{a} = 1$ and $\psi_j = \alpha_j \chi_{D_j}$ with $D_j = [(j-1)\frac{1}{128}, j\frac{1}{128}]$, $\mathbf{y} = (y_j)_{j=1, \dots, 128}$ and $\alpha_j = \frac{0.6}{j^\zeta}$, $\zeta = 3$. Figure 1 shows the convergence behavior of the adaptive Smolyak algorithm for the approximation of the normalization constant considering a variation of the number of observation points as well as of the observational noise.




FIGURE 1. Comparison of the estimated error and actual error of the normalization constant $Z = \mathbb{E}^{\mu^\delta}[1]$ with respect to the cardinality $\#\Lambda$ of the index sets Λ_N (Clenshaw-Curtis quadrature) with $K = 1, 3, 9$ (number of observation points), $\eta \sim \mathcal{N}(0, 1)$ (l.), $\eta \sim \mathcal{N}(0, 0.5^2)$ (m.), $\eta \sim \mathcal{N}(0, 0.1^2)$ (r.).

A detailed discussion of the numerical experiments for the parametric, parabolic evolution problem with random coefficients can be found in [10]. The numerical experiments indicate that, as the observation noise with variance $\Gamma \rightarrow 0$, growth of the constants in the Smolyak quadrature error estimates. In [11], we present an asymptotic analysis and prove $C \sim \exp(b/\Gamma)$ for some constants $b, C > 0$. We also show in [11] that the Bayesian estimate admits a finite limit in the case $\Gamma \rightarrow 0$, and propose regularization of the integrand functions arising in the computation of the conditional expectation.

Acknowledgement: This work was supported by the European Research Council (ERC) under AdG 247277 STAHPDE.

REFERENCES

- [1] Albert Cohen and Abdellah Chkifa and Christoph Schwab, Breaking the curse of dimensionality in sparse polynomial approximation of parametric PDEs, Report 2013-25, Seminar for Applied Mathematics, ETH Zürich, 2013.
- [2] Abdellah Chkifa and Albert Cohen and Christoph Schwab, High-dimensional adaptive sparse polynomial interpolation and applications to parametric PDEs, *Journ. Found. Comp. Math.*, <http://dx.doi.org/10.1007/s10208-013-9154-z>, 2013.
- [3] Albert Cohen and Ronald A. DeVore and Christoph Schwab, Analytic regularity and polynomial approximation of parametric and stochastic elliptic PDEs, *Analysis and Applications* **9**(1) 11–47, <http://dx.doi.org/10.1142/S0219530511001728>, 2010.
- [4] Markus Hansen and Christoph Schwab, Sparse Adaptive Approximation of High Dimensional Parametric Initial Value Problems, *Vietnam Journal of Mathematics***41** (2) 181–215, <http://dx.doi.org/10.1007/s10013-013-0011-9>, 2013.
- [5] Markus Hansen and Christoph Schwab, Analytic regularity and nonlinear approximation of a class of parametric semilinear elliptic PDEs, *Mathematische Nachrichten*, <http://dx.doi.org/10.1002/mana201100131>, 2013.
- [6] Thomas Gerstner and Michael Griebel, Dimension-adaptive tensor-product quadrature *Computing* **71** 65-87, 2003.
- [7] Markus Hansen, Claudia Schillings and Christoph Schwab, Sparse Approximation Algorithms for High Dimensional Parametric Initial Value Problems, Report 2013-10, Seminar for Applied Mathematics, ETH Zürich (2013) (to appear in Proc. HPSC5, Hanoi, 2013 in Springer LNCSE).
- [8] Viet H. Hoang and Christoph Schwab, Analytic regularity and polynomial approximation of stochastic, parametric elliptic multiscale PDEs, *Analysis and Applications (Singapore)* **11** (01), <http://dx.doi.org/10.1142/S0219530513500012>, 2013.
- [9] Angela Kunoth and Christoph Schwab, Analytic regularity and GPC approximation for control problems constrained by linear parametric elliptic and parabolic PDEs, *SIAM Journ. Control and Optimization* **51**(3) pp. 2442 – 2471, 2013.
- [10] Claudia Schillings and Christoph Schwab, Sparsity in Bayesian Inversion of Parametric Operator Equations, Report 2013-17, Seminar for Applied Mathematics, ETH Zürich, 2013.
- [11] Claudia Schillings and Christoph Schwab, Scaling Limits in Computational Bayesian Inversion of Parametric and Stochastic Operator Equations (in preparation) , 2013.
- [12] Claudia Schillings and Christoph Schwab, Sparse, adaptive Smolyak quadratures for Bayesian inverse problems, *Inverse Problems*, **29** (6), <http://dx.doi.org/10.1088/0266-5611/29/6/065011>, 2012.
- [13] Christoph Schwab and Andrew M. Stuart, Sparse deterministic approximation of Bayesian inverse problems, *Inverse Problems* **28**(4) <http://dx.doi.org/10.1088/0266-5611/28/4/045003>, 2012.
- [14] Andrew M. Stuart, Inverse problems: a Bayesian perspective, *Acta Numerica*, **19** (2010) pp. 451–559, <http://dx.doi.org/10.1017/S0962492910000061>.

Recent Research Reports

Nr.	Authors/Title
2013-17	Cl. Schillings and Ch. Schwab Sparsity in Bayesian Inversion of Parametric Operator Equations
2013-18	V. Kazeev and Ch. Schwab Tensor approximation of stationary distributions of chemical reaction networks
2013-19	K. Schmidt and R. Hiptmair Asymptotic Boundary Element Methods for Thin Conducting Sheets
2013-20	R. Kruse Consistency and Stability of a Milstein-Galerkin Finite Element Scheme for Semilinear SPDE
2013-21	J. Sukys Adaptive load balancing for massively parallel multi-level Monte Carlo solvers
2013-22	R. Andreev and A. Lang Kolmogorov-Chentsov theorem and differentiability of random fields on manifolds
2013-23	P. Grohs and M. Sprecher Projection-based Quasiinterpolation in Manifolds
2013-24	P. Grohs and S. Keiper and G. Kutyniok and M. Schaefer α -Molecules: Curvelets, Shearlets, Ridgelets, and Beyond
2013-25	A. Cohen and A. Chkifa and Ch. Schwab Breaking the curse of dimensionality in sparse polynomial approximation of parametric PDEs
2013-26	A. Lang Isotropic Gaussian random fields on the sphere