



Conference Paper

Successes and challenges of the APP Coronagraph

Author(s):

Kenworthy, Matthew A.; Quanz, Sascha Patrick; Otten, Gilles; Meshkat, Tiffany; Codona, Johanan; Snik, Frans; Meyer, Michael E.; Kasper, Markus; Girard, Julien H.

Publication Date:

2013-06

Permanent Link:

<https://doi.org/10.3929/ethz-b-000086489> →

Originally published in:

IAU Symposium 8(299), <http://doi.org/10.1017/S1743921313007771> →

Rights / License:

[In Copyright - Non-Commercial Use Permitted](#) →

This page was generated automatically upon download from the [ETH Zurich Research Collection](#). For more information please consult the [Terms of use](#).

Successes and challenges of the APP Coronagraph

Matthew A. Kenworthy¹, Sascha Quanz², Gilles Otten¹, Tiffany Meshkat¹, Johanan Codona³, Frans Snik¹, Michael E. Meyer², Markus Kasper⁴ and Julien Girard⁵

¹Leiden Observatory, Leiden University, P.O. Box 9513, 2300 RA Leiden, The Netherlands
email: kenworthy@strw.leidenuniv.nl

²Institute for Astronomy, ETH Zurich, CH-8093 Zurich, Switzerland

³Steward Observatory, 933 N. Cherry Avenue, Tucson, AZ 85721

⁴European Southern Observatory, Garching bei Munchen, Germany

⁵European Southern Observatory, Vitacura, Cassilla 19001, Santiago, Chile

Abstract.

The Apodizing Phase Plate (APP) coronagraph has been used to image the exoplanet β Pictoris b and the protoplanet candidate around HD 100546, and is currently in use in surveys with NaCo at the VLT. Its success is due to its tolerance to tip-tilt pointing errors in current AO systems, which degrade the performance of nearly all other coronagraphs. Currently the sensitivity of the APP is limited by non-common path errors in the science camera systems and by its chromatic behaviour. We present the achromatized Vector APP coronagraph and address how we will measure and minimise non-common path errors with Focal Plane Wavefront Sensing algorithms.

Keywords. instrumentation: high angular resolution, stars: planetary systems, stars: individual (β Pic, HD 100546)

1. Introduction

Coronagraphs are used to suppress the diffraction halo around bright stars, revealing the presence of faint companions (brown dwarfs and planets) next to their primary star. There are many coronagraph designs that work at small inner working angles (Guyon *et al.* 2006) and continuing technological developments are ongoing (for a recent review see Mawet *et al.* 2012).

The Apodizing Phase Plate (APP) Coronagraph is a pupil-plane optic that is a realization of Phase Apodization Coronagraphy (Codona & Angel 2004; Codona *et al.* 2006). In common with other pupil plane reformatters (Carlotti *et al.* 2012; Carlotti 2013), it is therefore relatively insensitive to residual tip-tilt motions in the focal plane of the science camera from residual atmospheric correction and from vibrations in the telescope superstructure. A first light demonstrator on the MMT0 6.5m telescope at $\lambda = 4.85\mu\text{m}$ (Kenworthy *et al.* 2007), was followed by an L' band APP on the Very Large Telescope (Kenworthy *et al.* 2010a,c; Girard *et al.* 2010) and most recently at L' band on the Large Binocular Telescope in LMIRCam (Wilson *et al.* 2008; Kenworthy *et al.* 2010b; Leisenring *et al.* 2012). Characterizations of planets include β Pictoris b (Quanz *et al.* 2010) and the discovery of a low mass companion to HD 100546 (Quanz *et al.* 2013), along with its inclusion in several VLT surveys for extrasolar planets around nearby stars with debris disks (Meshkat *et al.* 2013) and A/F stars (Quanz *et al.* 2011).

2. Current Performance and Future Directions

APP observations have placed strong constraints on low mass objects at Solar System scales around Fomalhaut (Kenworthy *et al.* 2013), typically with contrasts of 12 mag at 0.4 arcseconds. The refinement of Principal Component Analysis (PCA) techniques (Amara & Quanz 2012; Soummer *et al.* 2012) lead to an additional ~ 0.5 mag of sensitivity (see Meshkat *et al.* these proceedings; Meshkat *et al.* (2013) submitted).

The performance of the APP coronagraph is limited by (i) the non-common path aberrations (NCPA) between the AO camera and the wavefront sensor path in the science instrument, and (ii) its chromatic design. With the APP we will use Phase Sorting Interferometry (Codona & Kenworthy 2013) which uses phase diversity from the AO-corrected atmosphere to sense NCPA. Another development is the Vector APP (VAPP), an implementation of the APP that uses multiple stacked layers of polarization phase plates to provide a broadband wavelength solution (Snik *et al.* 2012) that also provides two simultaneous and complementary PSFs with the dark D region on opposing sides around the star. New versions of this optic are currently being characterized at Leiden Observatory, and the APP is being considered as the baseline coronagraph for ERIS, the new thermal infrared imager for the VLT, and for METIS, the thermal infrared imager and spectrograph for the European ELT.

References

- Amara A. & Quanz S. P., 2012, MNRAS, 427, 948
 Carlotti A., 2013, A&A, 551, A10
 Carlotti A., Kasdin N. J., Vanderbei R. J., & Delorme J.-R., 2012, SPIE 8442.
 Codona J. L. & Angel R., 2004, ApJ, 604, L117
 Codona J. L. & Kenworthy M., 2013, ApJ, 767, 100
 Codona J. L., Kenworthy M. A., Hinz P. M., Angel J. R. P., & Woolf N. J., 2006, SPIE 6269, pp. 62691N
 Girard J. H. V., *et al.*, 2010, SPIE 7736
 Guyon O., Pluzhnik E. A., Kuchner M. J., Collins B., & Ridgway S. T., 2006, ApJS, 167, 81
 Kenworthy M., Quanz S., Meyer M., Kasper M., Girard J., Lenzen R., Codona J., & Hinz P., 2010a, The Messenger, 141, 2
 Kenworthy M. A., Codona J. L., Hinz P. M., Angel J. R. P., Heinze A., & Sivanandam S., 2007, ApJ, 660, 762
 Kenworthy M. A., Hinz P. M., Codona J. L., Wilson J. C., Skrutskie M. F., & Solheid E., 2010b, SPIE 7734
 Kenworthy M. A., Meshkat T., Quanz S. P., Girard J. H., Meyer M. R., & Kasper M., 2013, ApJ, 764, 7
 Kenworthy M. A., Quanz S. P., Meyer M. R., Kasper M. E., Lenzen R., Codona J. L., Girard J. H., & Hinz P. M., 2010c, SPIE 7735
 Leisenring J. M., *et al.*, 2012, SPIE 8446
 Mawet D., *et al.*, 2012, SPIE 8442
 Meshkat T., Kenworthy M. A., Su K. Y., & Bailey V. P., 2013, AAS 221, 149.28
 Quanz S. P., Amara A., Meyer M. R., Kenworthy M. A., Kasper M., & Girard J. H., 2013, ApJ, 766, L1
 Quanz S. P., Kenworthy M. A., Meyer M. R., Girard J. H. V., & Kasper M., 2011, ApJ, 736, L32
 Quanz S. P., Meyer M. R., Kenworthy M. A., Girard J. H. V., Kasper M., Lagrange A.-M., Apai D., Boccaletti A., Bonnefoy M., Chauvin G., Hinz P. M., & Lenzen R., 2010, ApJ, 722, L49
 Snik F., Otten G., Kenworthy M., Miskiewicz M., Escuti M., Packham C., Codona J., 2012, SPIE 8450
 Soummer R., Pueyo L., & Larkin J., 2012, ApJ, 755, L28
 Wilson J. C., Hinz P. M., Skrutskie M. F., Jones T., Solheid E., Leisenring J., Garnavich P., Kenworthy M., Nelson M. J., & Woodward C. E. 2008, SPIE 7013