

Combinatorial Depth Measures for Hyperplane Arrangements

Conference Paper**Author(s):**

Schnider, Patrick; Soberón, Pablo

Publication date:

2023-06

Permanent link:

<https://doi.org/https://doi.org/10.3929/ethz-b-000621951>

Rights / license:

[Creative Commons Attribution 4.0 International](#)

Originally published in:

258, <https://doi.org/10.4230/LIPIcs.SoCG.2023.55>

Combinatorial Depth Measures for Hyperplane Arrangements

Patrick Schneider ✉ 🏠 

Department of Computer Science, ETH Zürich, Switzerland

Pablo Soberón ✉ 🏠 

Department of Mathematics, Baruch College, City University of New York, NY, USA

Department of Mathematics, The Graudate Center, City University of New York, NY, USA

Abstract

Regression depth, introduced by Rousseeuw and Hubert in 1999, is a notion that measures how good of a regression hyperplane a given query hyperplane is with respect to a set of data points. Under projective duality, this can be interpreted as a depth measure for query points with respect to an arrangement of data hyperplanes. The study of depth measures for query points with respect to a set of data points has a long history, and many such depth measures have natural counterparts in the setting of hyperplane arrangements. For example, regression depth is the counterpart of Tukey depth. Motivated by this, we study general families of depth measures for hyperplane arrangements and show that all of them must have a deep point. Along the way we prove a Tverberg-type theorem for hyperplane arrangements, giving a positive answer to a conjecture by Rousseeuw and Hubert from 1999. We also get three new proofs of the centerpoint theorem for regression depth, all of which are either stronger or more general than the original proof by Amenta, Bern, Eppstein, and Teng. Finally, we prove a version of the center transversal theorem for regression depth.

2012 ACM Subject Classification Theory of computation → Computational geometry; Mathematics of computing → Combinatorics

Keywords and phrases Depth measures, Hyperplane arrangements, Regression depth, Tverberg theorem

Digital Object Identifier 10.4230/LIPIcs.SoCG.2023.55

Related Version *Full Version*: <https://arxiv.org/abs/2302.07768>

Funding *Pablo Soberón*: Supported by NSF DMS grant 2054419.

1 Introduction

A central topic in combinatorial geometry and computational geometry is the study of structural properties of finite families of points in Euclidean spaces. Studying which sets can be separated from others by hyperplanes is a natural question, which leads us to study combinatorial properties of convex sets. Classic results, such as Tverberg's theorem [26] and Rado's centerpoint theorem [17] follow from this line of thought.

In some cases, instead of being provided our data as a finite set of points in \mathbb{R}^d , we might receive it as a set of hyperplanes. Understanding which results for families of points transfer to families of hyperplanes is a natural question.

Given a hyperplane arrangement A in \mathbb{R}^d and a point q , we first consider the depth of q with respect to A as follows.

► **Definition 1.** *The regression depth of a query point q with respect to hyperplane arrangement A , denoted by $RD(A, q)$, is the minimum number of hyperplanes in A intersected by or parallel to any ray emanating from q .*

Note that if q lies on a hyperplane H , then any ray emanating from q intersects H . Regression depth has been widely studied [1, 7, 21, 22, 28]. In this manuscript we provide



© Patrick Schneider and Pablo Soberón;


licensed under Creative Commons License CC-BY 4.0

39th International Symposium on Computational Geometry (SoCG 2023).

Editors: Erin W. Chambers and Joachim Gudmundsson; Article No. 55; pp. 55:1–55:14

Leibniz International Proceedings in Informatics

LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



new structural results for regression depth, related to Tverberg's theorem and *enclosing depth*. In particular, given a finite arrangement A of hyperplanes in \mathbb{R}^d , we might measure the depth of a point q in A in several different ways, so we study general properties of depth measures with respect to arrangements of hyperplanes. This follows a similar approach recently taken for depth measures with respect to finite families of points [23].

Given an arrangement A of n hyperplanes, the existence of points with regression depth at least $n/(d+1)$ has been established by Amenta, Bern, Eppstein, and Teng [1], and later by Mizera [15] as well as Karasev [11]. This can be considered a hyperplane version of Rado's centerpoint theorem [17]. We give three new proofs of the existence of points with large regression depth. First, we prove a Tverberg-type theorem for hyperplanes, confirming a conjecture of Rousseeuw and Hubert [21].

► **Theorem 2.** *Let r, d be positive integers and A be an arrangement of at least $(r-1)(d+1)+1$ hyperplanes in \mathbb{R}^d . Then, there exists a point q in \mathbb{R}^d and a partition of A into r parts such that q has positive regression depth with respect to each of the r parts.*

This was previously known when $d = 2$ [21] or when r is a prime power [12, 13]. The version for prime powers by Karasev holds with a slightly more restrictive version of regression depth. Based on this result, we define the *hyperplane Tverberg depth* of a point.

► **Definition 3.** *The hyperplane Tverberg depth of a query point q with respect to hyperplane arrangement A , denoted by $HTvD(A, q)$, is the maximum r such that there is a partition of A into r parts such that q has positive regression depth with respect to each part.*

Our other two proofs are topological, and each also has stronger consequences. One proof based on a topological version of Helly's theorem shows the existence of points of high *open regression depth*, which is a slightly weaker measure of depth introduced in Section 5. The last proof, based on properties of vector bundles, works for regression depth in families of weighted arrangements.

Another way to measure the depth of a point with respect to a hyperplane arrangement is via k -enclosures. We say that an arrangement A k -encloses a query point q if A can be partitioned into $d+1$ pairwise disjoint subsets A_1, \dots, A_{d+1} , each of size k , such that for every choice $h_1 \in A_1, \dots, h_{d+1} \in A_{d+1}$ we have that $RD(\{h_1, \dots, h_{d+1}\}, q) \geq 1$.

► **Definition 4.** *The hyperplane enclosing depth of a query point q with respect to a hyperplane arrangement A , denoted by $HED(A, q)$, is the maximum k such that there is a sub-arrangement of A which k -encloses q .*

Given a finite hyperplane arrangement A , we prove the existence of points with high hyperplane enclosing depth with respect to A . In particular, our lower bound is linear in $|A|$. The existence of points with large enclosing depth for families of points has been established by Pach [16] and by Fabila-Monroy and Huemer [6] (see [23] for improved constants).

One striking generalization of Rado's centerpoint theorem is the central transversal theorem, proven independently by Dolnikov and by Živaljević and Vrećica [5, 29]. In Section 7 we prove an analogue for hyperplane arrangements. Given a hyperplane arrangement A in \mathbb{R}^d and a linear subspace L in \mathbb{R}^d , we denote by $A \cap L$ the restriction of A to L . In Theorem 28, we show that given $d-k+1$ different arrangements of hyperplanes in \mathbb{R}^d , there exists a k -dimensional linear subspace L such that the restrictions of each arrangement to L share a point with high regression depth.

In particular, just as the central transversal theorem generalizes the ham sandwich theorem, Theorem 28 has the following corollary.

► **Corollary 5.** *Let A_1, \dots, A_d be d hyperplane arrangements in \mathbb{R}^d . There exists a line ℓ through the origin in \mathbb{R}^d and a point $q \in \ell$ such that each of the two rays in ℓ starting from q intersects at least $|A_i|/2$ hyperplanes of A_i , for each $i = 1, \dots, d$.*

The corollary above is similar to mass partition results for families of hyperplanes with segments [3, 18], and to projective versions of the central transversal theorem [13].

2 Correspondence to depth measures for point sets

For an arrangement A and a query point q , we define the *dual of A at q* , denoted by A_q^* , as follows. For each hyperplane $h \in A$, let $p(h)$ be the unique point on h that is closest to q . We define A_q^* as the set formed by all these points, that is, $A_q^* := \{p(h) \mid h \in A\}$. Note that if q lies on k hyperplanes, then those k dual points coincide with q in A_q^* .

Using this duality, for every depth measure ρ on point sets we can define a corresponding depth measure ρ^* on hyperplane arrangements and vice versa, by setting $\rho^*(A, q) = \rho(A_q^*, q)$. We have the following observation.

- **Observation 6.**
1. *a ray r emanating from q intersects a hyperplane h if and only if the half-space r^\perp defined by the hyperplane through q orthogonal to r , oriented such that it contains r , contains $p(h)$;*
 2. *the point q has positive regression depth with respect to h_1, \dots, h_n if and only if it is in the convex hull of $p(h_1), \dots, p(h_n)$.*
 3. *the point q lies in the simplex defined by h_1, \dots, h_{d+1} if and only if it is in the interior of the convex hull of $p(h_1), \dots, p(h_{d+1})$.*

The three depth measures for hyperplane arrangements defined in Section 1 all have natural corresponding depth measures for point sets that follow immediately from Observation 6. For regression depth, the corresponding depth measure is *Tukey depth* (TD), which is defined as the minimum number of data points contained in any closed half-space containing the query point q [25]. For hyperplane Tverberg depth we get Tverberg depth (TvD), which is defined as the maximum r for which there exists an r -partition of the data points containing the query point q in their intersection. Finally, for hyperplane enclosing depth, we get *enclosing depth* (ED), which is defined as the maximum k for which there exists a subset of the data points that k -encloses the query point q [23].


► **Corollary 7.** *Let A be an arrangement of hyperplanes in general position in \mathbb{R}^d and let q be a query point. Then*

1. $RD(A, q) = TD(A_q^*, q)$;
2. $HTvD(A, q) = TvD(A_q^*, q)$;
3. $HED(A, q) = ED(A_q^*, q)$.

3 Axioms for hyperplane depth

Let $A^{\mathbb{R}^d}$ denote the family of all finite arrangements of hyperplanes in \mathbb{R}^d . A depth measure for hyperplanes is a function $\rho : (A^{\mathbb{R}^d}, \mathbb{R}^d) \rightarrow \mathbb{R}_{\geq 0}$ which assigns to each pair (A, q) consisting of a hyperplane arrangement A and a query point q a value, which describes how deep the query point q lies within the arrangement A . A depth measure is called *combinatorial* if it is the same for all points in a face of A . Similar to [23], we introduce some axioms, that reasonable depth measures for hyperplane arrangements should satisfy.

We say that a combinatorial depth measure for hyperplanes is *super-additive* if it satisfies the following four conditions.



■ **Figure 1** Hyperplane enclosing depth does not satisfy condition (iv): the point q has hyperplane enclosing depth 1 with respect to both the blue and the red lines, but its hyperplane enclosing depth with respect to the union of the two sets is still 1.

- (i) for all $A \in A^{\mathbb{R}^d}$ and $q \in \mathbb{R}^d$ and any hyperplane h we have $|\rho(A, q) - \rho(A \cup \{h\}, q)| \leq 1$,
- (ii) for all $A \in A^{\mathbb{R}^d}$ we have $\rho(A, q) = 0$ if q is in an unbounded cell of A ,
- (iii) for all $A \in A^{\mathbb{R}^d}$ we have $\rho(A, q) \geq 1$ if q is in a bounded cell or if q lies on a hyperplane of A ,
- (iv) for any disjoint subsets $A_1, A_2 \subseteq A$ and $q \in \mathbb{R}^d$ we have $\rho(A, q) \geq \rho(A_1, q) + \rho(A_2, q)$.

► **Observation 8.** *Regression depth and hyperplane Tverberg depth are super-additive, but hyperplane enclosing depth is not.*

For hyperplane enclosing depth, an example with $\text{HED}(A_1, q) = \text{HED}(A_2, q) = \text{HED}(A, q) = 1$ can be found in Figure 1.

► **Lemma 9.** *Let ρ be any combinatorial depth measure that satisfies conditions (i) and (ii). Then for all $A \in A^{\mathbb{R}^d}$ and $q \in \mathbb{R}^d$ we have $\rho(A, q) \leq \text{RD}(A, q)$.*

Proof. Let $\text{RD}(A, q) = k$. This means that there is a ray r which intersects or is parallel to some k hyperplanes of A . Removing these k hyperplanes, we get a new arrangement A' and we have $\text{RD}(A', q) = 0$. In particular, q is in an unbounded cell of A' and thus also $\rho(A', q) = 0$ by condition (ii). By condition (i) we have $\rho(A, q) \leq \rho(A', q) + k = k$. ◀

► **Lemma 10.** *Let ρ be any combinatorial depth measure that satisfies conditions (iii) and (iv). Then for all $A \in A^{\mathbb{R}^d}$ and $q \in \mathbb{R}^d$ we have $\rho(A, q) \geq \text{HTvD}(A, q)$.*

Proof. Let $\text{HTvD}(A, q) = k$. This means that there is a k -partition A_1, \dots, A_k such that q has regression depth ≥ 1 with respect to each part. By condition (iii) we have $\rho(A_i, q) \geq 1$ for each A_i . By condition (iv) we get $\rho(A, q) \geq \rho(A_1, q) + \dots + \rho(A_k, q) \geq k$. ◀

► **Lemma 11.** *For all $A \in A^{\mathbb{R}^d}$ and $q \in \mathbb{R}^d$ we have $\text{HTvD}(A, q) \geq \frac{1}{d} \text{RD}(A, q)$.*

Proof. By Corollary 7 we have $\text{HTvD}(A, q) = \text{TvD}(A_q^*, q)$ and $\text{RD}(A, q) = \text{TD}(A_q^*, q)$. It is well known that for any point set S in \mathbb{R}^d and any query point q we have $\text{TvD}(S, q) \geq \frac{1}{d} \text{TD}(S, q)$, see e.g. [1, 8, 19]. ◀

Combining all of the above, we get

► **Theorem 12.** *Let ρ be a super-additive depth measure for hyperplanes. Then for all $A \in A^{\mathbb{R}^d}$ and $q \in \mathbb{R}^d$ we have $RD(A, q) \geq \rho(A, q) \geq HTvD(A, q) \geq \frac{1}{d}RD(A, q)$.*

As we have seen above, not all depth measures are super-additive: hyperplane enclosing depth is an example of a measure that is not. To include more general depth measures, we define a second family of measures, defined by a weaker set of axioms. We call a combinatorial depth measure for hyperplanes *enclosable* if it satisfies the following conditions.

- (i) for all $A \in A^{\mathbb{R}^d}$ and $q \in \mathbb{R}^d$ and any hyperplane h we have $|\rho(A, q) - \rho(A \cup \{h\}, q)| \leq 1$,
- (ii) for all $A \in A^{\mathbb{R}^d}$ we have $\rho(A, q) = 0$ if q is in an unbounded cell of A ,
- (iii') for all $A \in A^{\mathbb{R}^d}$ we have $\rho(A, q) \geq k$ if A k -encloses q ,
- (iv') for all $A \in A^{\mathbb{R}^d}$ and $q \in \mathbb{R}^d$ and any hyperplane h we have $\rho(A \cup \{h\}, q) \geq \rho(A, q)$.

► **Observation 13.** *Regression depth, hyperplane Tverberg depth and hyperplane enclosing depth are all enclosable.*

By Lemma 9, any enclosable depth measure is bounded from above by regression depth. On the other hand, it follows immediately from conditions (iii') and (iv') that any enclosable depth measure is bounded from below by hyperplane enclosing depth. We finish this section by showing a lower bound for hyperplane enclosing depth. In Theorem 17 in [23] it was shown that there is a constant $c(d)$ such that for any point set S in \mathbb{R}^d and any query point q we have $ED(S, q) \geq c \cdot TD(S, q)$. Let now q be a point of largest regression depth for a hyperplane arrangement A . We will see in Theorem 24 that q has regression depth at least $\frac{|A|}{d+1}$. By Observation 6, this means $TD(A_q^*, q) \geq \frac{|A|}{d+1}$. By Theorem 17 in [23], it follows that $ED(A_q^*, q) \geq \frac{c|A|}{d+1}$. Using Observation 6 again, we deduce the following:

► **Theorem 14.** *Let A be an arrangement of hyperplanes in \mathbb{R}^d . There is a constant $c = c(d)$ such that there is a query point q with hyperplane enclosing depth $HED(A, q) \geq \frac{c|A|}{d+1}$.*

Combining all of the above, we get an analogue to Theorem 12.

► **Theorem 15.** *Let ρ be an enclosable depth measure for hyperplanes. Then for all $A \in A^{\mathbb{R}^d}$ and $q \in \mathbb{R}^d$ we have $RD(A, q) \geq \rho(A, q) \geq HED(A, q) \geq c \cdot RD(A, q)$.*

In particular, all combinatorial depth measures for hyperplanes that we consider in this paper are constant factor approximations of regression depth. In the next three sections, we give three lower bounds for the depth of a deepest point. In Section 4 we give a lower bound for hyperplane Tverberg depth, in Section 5 a slightly stronger bound for regression depth, and in Section 6 we give a lower bound for super-additive depth measures with contractible depth regions in the more general setting of weighted arrangements.

4 A first lower bound: Hyperplane Tverberg Depth

In this section we prove an analogue of Tverberg's theorem for hyperplane arrangements, resolving a conjecture by Rousseeuw and Hubert from 1999 [21]. Our proof is inspired by the proof of Tverberg's theorem by Roudneff [20], see also [2].

► **Theorem 2.** *Let r, d be positive integers and A be an arrangement of at least $(r-1)(d+1)+1$ hyperplanes in \mathbb{R}^d . Then, there exists a point q in \mathbb{R}^d and a partition of A into r parts such that q has positive regression depth with respect to each of the r parts.*

Proof. Let π be a partition of A into r parts, each of size at most $d + 1$. Note that π can have at most d parts of size $\leq d$. Define the following function $f_\pi : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$: for each point $q \in \mathbb{R}^d$, consider the point set A_q^* . The partition π induces a partition of this point set into parts $X_1(q), \dots, X_r(q)$. Let $B(q)$ be the smallest ball centered at q which for every part intersects the convex hull, and define $f_\pi(q)$ as the radius of this ball. As the map which for a hyperplane h assigns to a point in \mathbb{R}^d the closest point on h is continuous as a function of q , the function f_π is also continuous. Further, the function goes to infinity along any ray, so it attains a minimum. Denote by $C(q)$ the set of parts whose convex hulls $B(q)$ is tangent to. By general position, we may assume that $|C(q)| \leq d + 1$.

Let now π be a partition which minimizes $\min f_\pi$ and let p be a point where f_π attains its minimum. If $f_\pi(p) = 0$, then by Observation 6, p is the desired point. So, assume that $f_\pi(p) > 0$. For each $X_j(q)$ let $y_j(q)$ denote the unique point in $\text{conv}X_j(q)$ that minimizes the distance to q , i.e., $d(q, \text{conv}X_j) = \|q - y_j(q)\|$, and define $Y_j(q) \subset X_j(q)$ as the unique subset for which $y_j(q)$ lies in the relative interior of $\text{conv}(Y_j(q))$. In particular we can write $f_\pi(q) = \frac{1}{|C(q)|} \sum_{X_j \in C(q)} \|q - y_j(q)\|$, and its gradient as $\nabla f_\pi(q) = \frac{1}{|C(q)|} \sum_{X_j \in C(q)} (q - y_j(q))$. As f_π is minimized at p , we have $\nabla f_\pi(p) = 0$.


We claim that $C(p)$ consists of exactly $d + 1$ parts and that no d of the corresponding vectors $(p - y_j(p))$ lie in a common hyperplane with p . Assume for the sake of contradiction that the latter is not the case, that is, that there is a hyperplane h containing all except possibly one of the vectors $(p - y_j(p))$. Let ℓ be a line through p that is orthogonal to h . Note that all except possibly one of the affine hulls $\text{aff}Y_j(p)$ for $X_j(p) \in C(p)$ are parallel to ℓ . If there is a single vector not in h , then this vector induces a direction on ℓ . Move p a distance ε in the opposite direction. If all vectors are in h , then move p along ℓ in any direction. Call the resulting point p' . We can choose ε small enough that $C(p') = C(p)$. Let h' be the hyperplane through p' that is parallel to h and let h^+ be its side containing p . Consider now the point $y_j(p')$ for some $X_j(p) \in C(p)$. This point is in the relative interior of the points in $Y_j(p')$. Let $a(p') \in Y_j(p')$ and let $a(p)$ be the corresponding point in $Y_j(p)$. If $a(p)$ is on the same side of h as p' , then $d(a(p'), p') < d(a(p), p)$ and if $a(p)$ is on the other side then $d(a(p'), p') > d(a(p), p)$, see Figure 2. In particular, The affine subspace $Y_j(p')$ is not parallel to ℓ and the vector $(p' - y_j(p'))$ points into h^+ . As this holds for any $X_j \in C(p)$, then also the gradient $\nabla f_\pi(p') = \frac{1}{|C(p)|} \sum_{X_j \in C(p)} (p' - y_j(p'))$ points into h^+ , and as this is the side that contains p , this means that p cannot be a local minimum, which is a contradiction to the choice of p . It follows that any d of the vectors $(p - y_j(p))$ are linearly independent, and thus we need at least $d + 1$ of them to have $\nabla f_\pi(p) = 0$.

Thus, the ball $B(p)$ is tangent to exactly $d + 1$ convex hulls, and the $d + 1$ tangent hyperplanes form a simplex containing p in its interior. As there are at most d parts of size $\leq d$, there must be a point v in some X_j such that $B(p)$ still intersects the convex hull of $X_j \setminus \{v\}$. This point must lie on the same side as p of some other tangent hyperplane, say of X_i . Then adding v to X_i gives a new partition π' in which $B(p)$ intersects the interior of the convex hull of X_i . In particular, due to the arguments above, p is not a minimum of $f_{\pi'}$, and thus $\min f_{\pi'} < \min f_\pi$. This is a contradiction to the choice of π , showing that $\min f_\pi = 0$. ◀

From Theorem 2, for any super-additive depth measure the existence of a point with depth at least $\frac{|A|}{d+1}$ follows using Theorem 12.

The existence of a Tverberg theorem for regression depth naturally leads us to ask about a colorful version of such a result.

► **Conjecture 16.** *Let r, d be positive integers and A_1, \dots, A_{d+1} be sets of r hyperplanes each*



■ **Figure 2** Moving p to p' the affine hulls of Y_j are not parallel to ℓ anymore.

in \mathbb{R}^d . There is a partition of their union into r sets B_1, \dots, B_r such that $|A_i \cap B_j| = 1$ for every $i \in [d + 1], j \in [r]$ and a point q such that q has positive regression depth for each B_j .

In the plane, Karasev conjectured, provided the hyperplanes are in general position, such a partition could be found so that q was in the simplex determined by each B_j , since his Tverberg-type results for hyperplanes hold in that setting [11]. However, his conjecture and its natural extensions to \mathbb{R}^d have been disproved [4, 14]. Yet, those counterexamples do not disprove the regression depth version, in which the q can be in the simplex determined by each B_j or the union of the hyperplanes making B_j .

5 A second lower bound: topological Helly theorem

In this section, we give a proof for the centerpoint theorem for regression depth based on one of the first topological versions of Helly’s theorem, which states that given a finite family F of subsets of \mathbb{R}^d with the property that for any $d + 1$ or fewer of them their intersection is non-empty and contractible, there is a point in the intersection of all families [9]. In fact, this method proves a stronger statement: we will show that for an arrangement A in general position, there is always as point in a cell of A of regression depth $\lceil \frac{|A|-d}{d+1} \rceil$. As we will see, this implies that there is always a point of regression depth $\lfloor \frac{|A|}{d+1} \rfloor + 1$.

The basic idea is the following: given an arrangement A of hyperplanes, consider some direction ℓ , and for every point q in \mathbb{R}^d compute how many hyperplanes of A the open ray with direction ℓ emanating from q intersects. Denote this number by $\ell(q)$. Define $R_A(k, \ell)$ as the set of points where $\ell(q) \geq k$. As A is finite, there are only finitely many different such regions. If we can show that for $k = \lceil \frac{|A|-d}{d+1} \rceil$ the intersection of any $d + 1$ or fewer such regions is contractible, then the existence of a deep point as claimed above follows from the topological Helly theorem. In fact, our arguments will show that any non-empty depth region is contractible.

There is however one technical issue: the depth regions of regression depth are in general not contractible. Consider three lines in the plane that form a triangle. The regression depth is 1 on any line or in the interior of the triangle, but it is 2 on the three corners, where two

of the lines intersect. So, the region of depth 2 consists of three isolated points and is thus not contractible.

If we however look only at the 2-dimensional cells of a planar line arrangement, then it is easy to show that the closure of the union of cells of depth at least k is contractible: no cell can be completely surrounded by cells of larger depth, as any ray witnessing depth k , that is, intersecting exactly k lines, also witnesses that the other cells it intersects all have depth smaller than k .

To overcome this issue, we define a new measure, which we call *open regression depth*, denoted by RD' as follows: let A be an arrangement of hyperplanes in \mathbb{R}^d . We first slightly perturb A to get an arrangement A' in general position. In particular, in any k -dimensional affine subspace at most $d - k$ of the hyperplanes intersect. Reversing this perturbation induces a surjective map π of the faces of A' to the faces of A . For any face F of A , we call $\pi^{-1}(F)$ the faces *perturbed* from F . Note that if A was already in general position, then π is a bijection.

Consider now the perturbed arrangement A' . For any point $q \in \mathbb{R}^d$, define the open regression depth with respect to the perturbed arrangement as the minimum number of hyperplanes of A that any ray emanating from q crosses or is parallel to, where a ray crosses a hyperplane if there is a point in the relative interior of the ray that is also on the hyperplane. In other words, the open regression depth for perturbed arrangement is just the regression depth, where we do not count the hyperplanes that q lies on. The depth regions of open regression depth in a perturbed arrangement are the unions of cells with large enough depth, with lower-dimensional faces added whenever they are incident to only deep enough cells.

In order to extend the definition to the original arrangement, we define the open regression depth of a query point q in some face F_q of the arrangement A as $\text{RD}'(A, q) := \max_{q' \in F \in F_q} \{\text{RD}'(A', q')\}$, that is, as the maximum open regression depth of any point in one of the faces perturbed from F_q . Note that we can perturb the arrangement in a deterministic way, ensuring that the open regression depth is well defined. The following lemma follows immediately from the definition:

► **Lemma 17.** *For any arrangement of hyperplanes A and any query point q , we have $\text{RD}'(A, q) \leq \text{RD}(A, q)$.*

In particular, proving the existence of deep points for open regression depth implies the existence of deep points for regression depth. Note, however, that open regression depth is not super-additive: it does not satisfy condition (iii). We will now prove the existence of deep points for open regression depth using the approach sketched above. We show that we have the necessary ingredients to apply the topological Helly theorem, starting with the contractability of the relevant regions. Recall that we defined the regions $R_A(k, \ell)$ as the set of points where $\ell(q) \geq k$ for a hyperplane arrangement A and a direction ℓ , where we considered the relevant ray to be open, that is, not containing q . Also recall that as A is finite, it is sufficient to restrict our attention to finitely many directions, and we may assume that these directions are d -wise linearly independent, that is, any d of them span a d -dimensional cone.

Our proof of contractability requires some algebraic topology and is independent of the rest of the manuscript, so we defer its proof to the full version [24]. For our purposes, a *homology cell* is the same as a contractible space.

► **Lemma 18.** *Let A_1, \dots, A_m be open subsets of \mathbb{R}^d . Assume that each set is a homology cell and that the union of any d of them is a homology cell. Then $\bigcap A := \bigcap_{i=1}^m A_i$ is either empty or a homology cell.*

We can apply this result to our setting.

► **Lemma 19.** *Let ℓ_1, \dots, ℓ_m be directions in d -wise general position in \mathbb{R}^d , let A be a hyperplane arrangement and let $R(k) := \bigcap_{i=1}^m R_A(k, \ell_i)$. If $R(k) \neq \emptyset$ then $R(k)$ is contractible.*

In particular, the depth regions, that is, the intersections of $R_A(k, \ell)$ over all considered directions ℓ is contractible.

Proof. As in our setting homology cells are contractible, by Lemma 18 it suffices to show that the union of any d regions $R_A(k, \ell_1), \dots, R_A(k, \ell_d)$ is contractible. Denote this union by $U(k)$ and let $n = |A|$. Let $C \subsetneq \mathbb{R}^d$ be the cone spanned by the d directions and let ℓ_0 be a direction in $-C$. In particular, moving from any point in $U(k)$ in direction ℓ_0 we never leave $U(k)$. Thus, $U(k)$ is contractible. ◀

The final property that we need in order to apply the topological Helly theorem is that the intersection of any $d + 1$ or fewer regions is non-empty.

► **Lemma 20.** *Let ℓ_1, \dots, ℓ_m be directions in \mathbb{R}^d , $m \leq d+1$, let A be a hyperplane arrangement and let $R := \bigcap_{i=1}^m R_A(k, \ell_i)$ for $k \leq \lceil \frac{|A|-d}{d+1} \rceil$. Then $R \neq \emptyset$.*

Proof. For every direction ℓ_i let h_i be a hyperplane orthogonal to ℓ_i which bounds a half-space h_i^+ that contains $R_A(k, \ell_i)$. In particular, for any point p in h_i^+ , moving p in direction ℓ_i , we eventually enter $R_A(k, \ell_i)$ and never leave it again. Thus, if all these half-spaces have a common intersection, then this intersection can be translated to lie in R , showing $R \neq \emptyset$. So, assume that these half-spaces have an empty intersection. As we assumed that any d of our directions are linearly independent, this can only happen for $m = d + 1$. In this case, we find a point $q \in \mathbb{R}^d$ such that the $d + 1$ (closed) rays emanating from q with directions $-\ell_i$ all intersect strictly more than $|A| - k$ hyperplanes of A . Each hyperplane that does not contain q can intersect at most d of the rays, and by the general position assumption, at most d hyperplanes contain q . Thus, if x denotes the number of intersections between rays and hyperplanes, we have

$$(|A| - k)(d + 1) < x \leq (|A| - d)d + d(d + 1) = (|A| + 1)d.$$

Rearranging this and using that all numbers are integers gives $k > \lceil \frac{|A|-d}{d+1} \rceil$, which is a contradiction to the assumption, showing that $R \neq \emptyset$. ◀

Now we have all the ingredients that are necessary for the topological Helly theorem, and we deduce the following

► **Corollary 21.** *For every hyperplane arrangement A in \mathbb{R}^d there is a point $q \in \mathbb{R}^d$ for which the open regression depth is $RD^+(A, q) \geq \lceil \frac{|A|-d}{d+1} \rceil$.*

In particular, by the definition of open regression depth, if the arrangement A is in general position, such a point can be chosen in a cell of A . It remains to show that we can get even deeper points for regression depth.

► **Lemma 22.** *For every hyperplane arrangement A in \mathbb{R}^d there is a point $q \in \mathbb{R}^d$ for which the regression depth is $RD(A, q) \geq \lfloor \frac{|A|}{d+1} \rfloor + 1$.*

Proof. Consider a point q in a cell C of maximum open regression depth k , and let ∂C be the boundary of the cell C . If there is a point on ∂C with open regression depth k , then this point has regression depth $k + 1$, and the claim follows. So assume that the open regression depth is strictly smaller everywhere on ∂C . Then we again find $d + 1$ directions such that

the rays emanating from q with these directions intersect exactly k hyperplanes. Looking at the opposite directions, the rays thus intersect exactly $|A| - k$ hyperplanes, and as q lies in the interior of a cell every hyperplane intersects at most d rays. Analogous to the proof of Lemma 20 we thus get $k > \lceil \frac{|A|}{d+1} \rceil$. This proves the claim for all cases where $d + 1$ does not divide $|A|$. If $d + 1$ divides $|A|$, note that as soon as one of the hyperplanes only intersects $d - 1$ of the considered rays, then we get $k > \lceil \frac{|A|+1}{d+1} \rceil$, and the claim follows again. So, assume that each hyperplane intersects exactly d rays. This gives a partition of the set of hyperplanes into $d + 1$ parts, each of size $\frac{|A|}{d+1}$ defined by the ray they do not intersect. The boundary ∂C inherits this partition, and each of the parts is contractible. In particular, ∂C contains a vertex q that is the intersection of d hyperplanes of d different parts. Now every ray emanating from q must intersect all hyperplanes of some part, but also lies on at least $d - 1$ other hyperplanes, showing that the regression depth of q is at least $k + d - 1$, which is a contradiction to the assumption that the open regression depth is strictly smaller everywhere on ∂C . ◀

Using the above insights, we can also conclude the contractability of many regions of regression depth.

► **Lemma 23.** *Let $k \leq \lceil \frac{|A|}{d+1} \rceil$. Then the region R of points p whose regression depth is $RD(A, q) \geq k$ is contractible.*

Proof. If there is a point with open regression depth k , then R is just the closure of the region of points with open regression depth at least k , which is contractible by Lemma 19. Otherwise, by the proof of Lemma 22, R is the union of faces incident only to cells of maximum open regression depth. As no cell is completely surrounded by deeper faces there is a contraction from a cell of maximum open regression depth to the deeper faces incident to it. Thus, as the region of maximum open regression depth is contractible, so is R . ◀

6 A third lower bound: weighted arrangements

In this section we give yet another proof for the existence of points with large regression depth. The proof we give here works for (and actually requires) the more general case of *weighted arrangements* of hyperplanes. A weighted arrangement of hyperplanes is a tuple (A, w) consisting of a finite arrangement A of hyperplanes and a weight function $w : A \rightarrow \mathbb{R}_{\geq 0}$ which assigns to each hyperplane a weight. By a slight abuse of notation we will often just write A for a weighted arrangement. For a subarrangement $A' \subseteq A$ we have $w'(h) \leq w(h)$, where w' is the weight function on A' , and we write $w'(A') := \sum_{h \in A'} w'(h)$. We say that $A' \subsetneq A$ is a strict subset of A if the underlying hyperplane arrangement of A' is a strict subset of that of A . The definition of regression depth extends to weighted arrangements: for any ray r emanating from a query point q , let $A(r)$ be the hyperplanes intersected by r . Then, the regression depth $RD(A, q)$ of q is the minimum of $w(A(r))$ taken over all rays emanating from q . This definition is similar to, but more restrictive than a measure-theoretic generalization of regression depth considered by Mizera [15].

Our proof also works for more general families of depth measures on weighted hyperplane arrangements. We extend the definition of super-additive depth measures above to weighted hyperplane arrangements as follows:

- (i) for all $A \in A^{\mathbb{R}^d}$ and $q \in \mathbb{R}^d$ and any hyperplane h we have $|\rho(A, q) - \rho(A \cup \{h\}, q)| \leq w(h)$,
- (ii) for all $A \in A^{\mathbb{R}^d}$ we have $\rho(A, q) = 0$ if q is in an unbounded cell of A ,

- (iii) for all $A \in \mathbb{R}^d$ we have $\rho(A, q) \geq \min\{w(h) \mid h \in A\}$ if q is in a bounded cell or if q lies on a hyperplane of A ,
- (iv) for any disjoint subsets $A_1, A_2 \subseteq A$ and $q \in \mathbb{R}^d$ we have $\rho(A, q) \geq \rho(A_1, q) + \rho(A_2, q)$.

Note that any hyperplane arrangement can be considered as a weighted hyperplane arrangement by assigning weight 1 to each hyperplane. On the other hand, each depth measure for hyperplane arrangement can be extended to a depth measure on weighted hyperplanes: using the fact that \mathbb{Q} is dense in \mathbb{R} , we can place multiple hyperplanes in the same position and then normalize to get a weighted arrangement.

For a weighted arrangement of hyperplanes A and a depth measure ρ denote by $R_\rho^A(\alpha) := \{q \in \mathbb{R}^d \mid \rho(A, q) \geq \alpha\}$ the α -depth region. The median region, which is the deepest non-empty depth region, is denoted by M_ρ^A .

► **Theorem 24.** *Let A be a weighted arrangement of hyperplanes in \mathbb{R}^d and let ρ be a super-additive depth measure on weighted hyperplanes whose depth regions are compact and contractible. Then there exists a point $q \in \mathbb{R}^d$ for which $\rho(A, q) \geq \frac{w(A)}{d+1}$.*

Before we prove Theorem 24, we give some lemmata that we will need in the proof. The first lemma concerns a generalization of a section in a vector bundle. Let $\pi : E \rightarrow B$ be a real vector bundle over a compact manifold B . Following [29] we say that $\phi : B \rightarrow E$ is a *multisection* if for every $x \in B$ we have that $\phi(x) \subseteq F_x := \pi^{-1}(x)$. We further say that ϕ is *contractible* if it is contractible in each fiber, that is, for every $x \in B$ the set $\phi(x)$ is contractible. Finally, we say that ϕ is compact if $\Gamma(\phi) := \{(x, v) \mid v \in \phi(x)\} \subseteq B \times E$ is compact. For any multisection ϕ , denote by $Z(\phi)$ its intersection with the zero section. In the full version [24] we show the following

► **Lemma 25.** *Let $\pi : E \rightarrow B$ be a real vector bundle over a compact manifold B . Let ϕ be a compact contractible multisection. Then there is a section s with $Z(s) = Z(\phi)$. In particular, if π has no nowhere zero section, then ϕ must intersect the zero section.*

The second lemma is about partitions of hyperplane arrangements.

► **Lemma 26.** *Let ρ be a depth measure for weighted hyperplanes whose depth regions are compact and contractible and let A be a weighted hyperplane arrangement in \mathbb{R}^d with $|A| \geq d + 2$. Then there exists a partition of A into strict subarrangements A_1 and A_2 whose median regions intersect.*

The proof is analogous to the proof of Lemma 9 in [23], replacing Proposition 1 from [29] with our Lemma 25. For a full proof we refer to the full version [24].

Proof of Theorem 24. Let A be a weighted arrangement of hyperplanes in \mathbb{R}^d . We prove the statement by induction on the number of hyperplanes in A . If A consists of at most $d + 1$ hyperplanes, it follows from condition (iii) that $\rho(A, q) \geq \frac{w(A)}{d+1}$ for some $q \in \mathbb{R}^d$: just take q as any point on a hyperplane of maximum weight. So assume that A consists of at least $d + 2$ hyperplanes. By assumption the depth regions are compact and contractible. Thus, by Lemma 26, we can partition A into strict subarrangements A_1 and A_2 whose median regions intersect. As both A_1 and A_2 are strict subarrangements, by the induction hypothesis for any point q in the intersection of their median regions we have $\rho(A_1, q) \geq \frac{w_1(A_1)}{d+1}$ and $\rho(A_2, q) \geq \frac{w_2(A_2)}{d+1}$. As ρ satisfies condition (iv), for any such point we thus have

$$\rho(A, q) \geq \rho(A_1, q) + \rho(A_2, q) \geq \frac{w_1(A_1) + w_2(A_2)}{d + 1} = \frac{w(A)}{d + 1}.$$



At this point, it is not clear how we can use Theorem 24 to prove the existence of centerpoints for regression depth. If we look at the depth regions of regression depth, we have seen in Section 5 that they are in general not contractible. To overcome this issue, we have introduced open regression depth and argued that the depth regions of open regression depth are contractible, and these arguments go through even if the arrangement is weighted. However, for a hyperplane arrangement in general position, these regions are by definition open, and thus not compact. Further, open regression depth is not a super-additive depth measure, as it does not satisfy condition (iii). In particular, if A consists of a single hyperplane, then the open regression depth is 0 everywhere, and so the base case of the proof of Theorem 24 fails. However, as we have seen in Lemma 23, if $k \leq \lceil \frac{|A|}{d+1} \rceil$ the region of regression depth at least k is contractible. Again, the involved arguments go through if the arrangement is weighted, implying that if $k \leq \frac{w(A)}{d+1}$, then the region of regression depth at least k is contractible. Thus, defining a new measure *truncated regression depth* by

$$\text{TRD}(A, q) := \min \left(\frac{w(A)}{d+1}, \text{RD}(A, q) \right),$$

we get a measure whose depth regions are closed and contractible. Clearly, the only unbounded regions are the ones containing an unbounded face of the arrangement, and we can make those compact by intersecting with a sufficiently large ball. Finally, as regression depth is super-additive, so is truncated regression depth, and by definition, truncated regression depth is bounded from above by regression depth. We thus have the following:

► **Lemma 27.** *Truncated regression depth is a super-additive depth measure for hyperplane arrangements which has compact and contractible depth regions. Further, for every arrangement A and every point q we have $\text{TRD}(A, q) \leq \text{RD}(A, q)$.*

It now follows from Theorem 24 that there is always a point of truncated regression depth $\text{TRD}(A, q) \geq \frac{w(A)}{d+1}$ and such a point also has regression depth $\text{RD}(A, q) \geq \frac{w(A)}{d+1}$.

7 A regression depth version of the center transversal theorem

Let A be an arrangement of hyperplanes in \mathbb{R}^d . Assume that the origin is not contained in any hyperplane in A . Let L be a k -dimensional linear subspace of \mathbb{R}^d . Then $A \cap L$ is a hyperplane arrangement in L . In particular, we can again study the depth of points $q \in L$ within the Euclidean space L with respect to the arrangement $A \cap L$. Note however that $A \cap L$ might have smaller cardinality than A , as some hyperplanes of A might be parallel to L . In fact, if all of them are parallel to L , then $A \cap L$ is empty. We define the regression depth of $q \in L$ with respect to $A \cap L$ as the minimum number of hyperplanes in A intersected by or parallel to any ray in L emanating from q , and denote it by $\text{RD}(A, q, L)$. In particular, if all hyperplanes in A are parallel to L , then $\text{RD}(A, q, L) = |A|$ for all $q \in L$. This definition extends to open regression depth and truncated regression depth, where we truncate at $\frac{|A_i|}{k+1}$.

► **Theorem 28.** *Let $1 \leq k \leq d$ be integers and A_1, \dots, A_{d-k+1} be $d-k+1$ finite arrangements of hyperplanes in \mathbb{R}^d . Then there exists a k -dimensional linear subspace L and a point $q \in L$ such that q has regression depth $\text{RD}(A_i, q, L) \geq \frac{|A_i|}{k+1}$ in L for every $i \in \{1, \dots, d-k+1\}$.*

Proof. We will prove the statement for truncated regression depth, which will imply the theorem as regression depth is bounded from below by truncated regression depth. Consider the Grassmann manifold $Gr_k(\mathbb{R}^d)$ of all k -dimensional subspaces of \mathbb{R}^d . Let γ_k^d be the canonical bundle over $Gr_k(\mathbb{R}^d)$, which has total space $E := \{(L, v) \mid v \in L\}$ and whose

projection $\pi : E \rightarrow Gr_k(\mathbb{R}^d)$ is given by $\pi((L, v)) = L$. For an arrangement A_i , let $R_i(L)$ be the set of points in L that have large depth, that is, $R_i(L) := \{v \in L \mid \text{TRD}(A_i, v, L) \geq \frac{|A_i|}{k+1}\}$. By Lemma 27, each $R_i(L)$ is compact and contractible. Further, when a hyperplane $h \in A_i$ becomes parallel to L , the depth of any point can only increase, thus $R_i(L') \subseteq R_i(L)$ for any L' in a small neighborhood of L . Thus, R_i is a compact contractible multisection. Define the negative multisection $-R_i$ by reflecting $R_i(L)$ at the origin for each L , and for each $i \in \{1, \dots, d-k\}$ consider $Q_i := R_{d-k+1} - R_i$, defined by taking the Minkowski sum of $R_{d-k+1}(L)$ and $-R_i(L)$ on each L . As Minkowski sums of compact and contractible sets are again compact and contractible, Q_i is again a compact contractible multisection. In particular, by Lemma 25, there are sections s_i whose zeroes coincide with the zeroes of Q_i . It was shown in [29], Prop. 2 (see also [5], Lem. 1), that any $d-k$ sections on γ_k^d must have a common zero, that is, there is a subspace L such that $s_1(L) = \dots = s_{d-k}(L) = 0$. By the definition of the sections s_i , this implies that there is a point $q \in L$ such that $q \in R_i(L)$ for all $i \in \{1, \dots, d-k+1\}$. In particular, $\text{TRD}(A_i, q, L) \geq \frac{|A_i|}{k+1}$ in L for every $i \in \{1, \dots, d-k+1\}$. ◀

Since there is a regression depth version of the center transversal theorem and of Tverberg's theorem, a natural question is if there is a generalization of both. This is still open in the case of finite families of points, since it was conjectured by Tverberg and Vrećica in 1993 [27].

► **Conjecture 29.** *Let $1 \leq k \leq d$ be integers and A_1, \dots, A_{d-k+1} be $d-k+1$ finite arrangements of hyperplanes in \mathbb{R}^d . Assume that $|A_i| = (k+1)(r_i-1) + 1$ for some positive integer r_i , for each $i = 1, \dots, d-k+1$. Then, there exists a k -dimensional subspace L , a point $q \in L$, and a partition of each A_i into r_i parts $A_i^{(1)}, \dots, A_i^{(r_i)}$ such that $\text{RD}(A_i^{(j)}, q, L) \geq 1$ for each $i = 1, \dots, d-k+1, j = 1, \dots, r_i$.*

The classic conjecture for families of points, which has similar parameters, has only been confirmed when all r_i are powers of the same prime p and pk is even [10].

References

- 1 Nina Amenta, Marshall Bern, David Eppstein, and S H Teng. Regression depth and center points. *Discrete & Computational Geometry*, 23(3):305–323, 2000.
- 2 Imre Bárány and Pablo Soberón. Tverberg's theorem is 50 years old: a survey. *Bulletin of the American Mathematical Society*, 55(4):459–492, 2018.
- 3 Sergey Bereg, Ferran Hurtado, Mikio Kano, Matias Korman, Dolores Lara, Carlos Seara, Rodrigo I Silveira, Jorge Urrutia, and Kevin Verbeek. Balanced partitions of 3-colored geometric sets in the plane. *Discrete Applied Mathematics*, 181:21–32, January 2015. doi:10.1016/j.dam.2014.10.015.
- 4 J. P. Carvalho and P. Soberón. Counterexamples to the colorful Tverberg conjecture for hyperplanes. *Acta Math. Hungar.*, 167(2):385–392, 2022. doi:10.1007/s10474-022-01249-8.
- 5 VL Dol'nikov. Transversals of families of sets in and a connection between the Helly and Borsuk theorems. *Russian Academy of Sciences. Sbornik Mathematics*, 79(1):93, 1994.
- 6 Ruy Fabila-Monroy and Clemens Huemer. Carathéodory's theorem in depth. *Discrete Comput. Geom.*, 58(1):51–66, 2017. doi:10.1007/s00454-017-9893-8.
- 7 Radoslav Fulek, Andreas F. Holmsen, and János Pach. Intersecting Convex Sets by Rays. *Discrete & Computational Geometry*, 42(3):343–358, 2009. doi:10.1007/s00454-009-9163-5.
- 8 Sarel Har-Peled and Timothy Zhou. Improved Approximation Algorithms for Tverberg Partitions. *arXiv preprint arXiv:2007.08717*, 2020.
- 9 Eduard Helly. Über Systeme abgeschlossener Mengen mit gemeinschaftlichen Punkten. *Monatshfte d. Mathematik*, 37:281–302, 1930.

- 10 Roman N. Karasev. Tverberg's Transversal Conjecture and Analogues of Nonembeddability Theorems for Transversals. *Discrete & Computational Geometry*, 38(3):513–525, December 2007. doi:10.1007/s00454-007-1355-2.
- 11 Roman N. Karasev. Dual theorems on central points and their generalizations. *Sbornik: Mathematics*, 199(10):1459–1479, 2008. doi:10.1070/sm2008v199n10abeh003968.
- 12 Roman N. Karasev. Tverberg-Type Theorems for Intersecting by Rays. *Discrete & Computational Geometry*, 45(2):340–347, 2011. doi:10.1007/s00454-010-9294-8.
- 13 Roman N. Karasev and Benjamin Matschke. Projective Center Point and Tverberg Theorems. *Discrete & Computational Geometry*, 52(1):88–101, 2014. doi:10.1007/s00454-014-9602-9.
- 14 Seunghun Lee and Kangmin Yoo. On a conjecture of Karasev. *Comput. Geom.*, 75:1–10, 2018. doi:10.1016/j.comgeo.2018.06.003.
- 15 Ivan Mizera. On depth and deep points: a calculus. *The Annals of Statistics*, 30(6):1681–1736, 2002.
- 16 János Pach. A Tverberg-type result on multicolored simplices. *Computational Geometry*, 10(2):71–76, 1998. doi:10.1016/s0925-7721(97)00022-9.
- 17 Richard Rado. A Theorem on General Measure. *Journal of the London Mathematical Society*, s1-21(4):291–300, 1946. doi:10.1112/jlms/s1-21.4.291.
- 18 Edgardo Roldán-Pensado and Pablo Soberón. A survey of mass partitions. *Bull. Amer. Math. Soc. (N.S.)*, 59(2):227–267, 2022. doi:10.1090/bull1/1725.
- 19 David Rolnick and Pablo Soberón. Algorithms for Tverberg's theorem via centerpoint theorems. *arXiv preprint arXiv:1601.03083*, 2016.
- 20 Jean-Pierre Roudneff. Partitions of points into simplices with k -dimensional intersection. part I: The conic Tverberg's theorem. *European Journal of Combinatorics*, 22(5):733–743, 2001.
- 21 Peter J. Rousseeuw and Mia Hubert. Depth in an arrangement of hyperplanes. *Discrete & Computational Geometry*, 22(2):167–176, 1999.
- 22 Peter J. Rousseeuw and Mia Hubert. Regression depth. *J. Amer. Statist. Assoc.*, 94(446):388–433, 1999. With discussion and a reply by the authors and Stefan Van Aelst. doi:10.2307/2670155.
- 23 Patrick Schnider. Enclosing depth and other depth measures. *arXiv preprint arXiv:2103.08421*, 2021.
- 24 Patrick Schnider and Pablo Soberón. Combinatorial depth measures for hyperplane arrangements, 2023. doi:10.48550/ARXIV.2302.07768.
- 25 John W. Tukey. Mathematics and the picturing of data. In *Proc. International Congress of Mathematicians*, pages 523–531, 1975.
- 26 Helge Tverberg. A generalization of Radon's theorem. *J. London Math. Soc.*, 41(1):123–128, 1966.
- 27 Helge Tverberg and Siniša T. Vrećica. On Generalizations of Radon's Theorem and the Ham Sandwich Theorem. *European Journal of Combinatorics*, 14(3):259–264, 1993. doi:10.1006/eujc.1993.1029.
- 28 Marc van Kreveld, Joseph S. B. Mitchell, Peter Rousseeuw, Micha Sharir, Jack Snoeyink, and Bettina Speckmann. Efficient algorithms for maximum regression depth. *Discrete Comput. Geom.*, 39(4):656–677, 2008. doi:10.1007/s00454-007-9046-6.
- 29 Rade T. Živaljević and Siniša T. Vrećica. An extension of the ham sandwich theorem. *Bulletin of the London Mathematical Society*, 22(2):183–186, 1990.