

Long-range and short-range dihadron angular correlations in central PbPb collisions at root $s(NN)=2.76$ TeV

Journal Article

Author(s):

CMS Collaboration; Chatrchyan, Serguei; Bortignon, Pierluigi; Caminada, Lea; Chanon, Nicolas; Chen, Zhiling; Cittolin, Sergio; Dissertori, Günther; Dittmar, Michael; Eugster, Jürg; Freudenreich, Klaus; Grab, Christoph; Hintz, Wieland; Lecomte, Pierre; Lustermann, Werner; Marchica, Carmelo; Martinez Ruiz del Arbol, Pablo; Meridiani, Paolo; Milenovic, Predrag; Moortgat, Filip; Nägeli, Christoph; Nef, Pascal; Nessi-Tedaldi, Francesca; Pape, Luc; Pauss, Felicitas; Punz, Thomas; Rizzi, Andrea; Ronga, Frédéric J.; Rossini, Marco; Sala, Leonardo; Sanchez, Ann-Karin; Sawley, Marie-Christine; Stieger, Benjamin; Tauscher, Ludwig; Thea, Alessandro; Theofilatos, Konstantinos; Treille, Daniel; Urscheler, Christina; Wallny, Rainer; Weber, Marc; Wehrli, Lukas; Weng, Joanna; et al.

Publication date:

2011-07

Permanent link:

<https://doi.org/10.3929/ethz-b-000039710>

Rights / license:

[Creative Commons Attribution-NonCommercial 4.0 International](#)

Originally published in:

Journal of High Energy Physics 2011(7), [https://doi.org/10.1007/JHEP07\(2011\)076](https://doi.org/10.1007/JHEP07(2011)076)

Long-range and short-range dihadron angular correlations in central PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$

The CMS collaboration

ABSTRACT: First measurements of dihadron correlations for charged particles are presented for central PbPb collisions at a nucleon-nucleon center-of-mass energy of 2.76 TeV over a broad range in relative pseudorapidity ($\Delta\eta$) and the full range of relative azimuthal angle ($\Delta\phi$). The data were collected with the CMS detector, at the LHC. A broadening of the away-side ($\Delta\phi \approx \pi$) azimuthal correlation is observed at all $\Delta\eta$, as compared to the measurements in pp collisions. Furthermore, long-range dihadron correlations in $\Delta\eta$ are observed for particles with similar ϕ values. This phenomenon, also known as the “ridge”, persists up to at least $|\Delta\eta| = 4$. For particles with transverse momenta (p_{T}) of 2-4 GeV/c, the ridge is found to be most prominent when these particles are correlated with particles of $p_{\text{T}} = 2-6 \text{ GeV}/c$, and to be much reduced when paired with particles of $p_{\text{T}} = 10-12 \text{ GeV}/c$.

KEYWORDS: Hadron-Hadron Scattering

Contents

1	Introduction	1
2	Data and analysis	2
3	Results	5
3.1	Associated yield distributions versus $\Delta\phi$	6
3.2	Integrated associated yield	8
3.3	Fourier decomposition of $\Delta\phi$ distributions	9
4	Summary	11
	The CMS collaboration	16

1 Introduction

Measurements of dihadron azimuthal correlations [1–7] have provided a powerful tool to study the properties of the strongly interacting medium created in ultrarelativistic nuclear collisions [8–11]. An early indication of strong jet-medium interactions at RHIC was the absence of high-transverse-momentum (high- p_T) back-to-back particle pairs in dihadron correlation measurements [1] and the corresponding enhancement of low- p_T hadrons recoiling from a high- p_T leading, or “trigger”, particle [3]. The recent observations of the suppression of high- p_T charged hadrons [12] and of asymmetric energies of reconstructed jets [13, 14] in PbPb collisions at the Large Hadron Collider (LHC) provide further evidence of jet quenching, suggesting a large energy loss for partons traversing the produced medium.

At RHIC, extending dihadron azimuthal correlation measurements to larger relative pseudorapidities resulted in the discovery of a ridge-shaped correlation in central AuAu collisions between particles with small relative azimuthal angles ($|\Delta\phi| \approx 0$), out to very large relative pseudorapidity ($|\Delta\eta|$) [2, 6]. Although the “ridge” has been qualitatively described by several different models [15–26], its origin is still not well understood. Some models attribute the ridge to jet-medium interactions, while others attribute it to the medium itself. The ridge has been observed for particles with transverse momenta from several hundred MeV/ c to a few GeV/ c . However, the character of the ridge for even higher- p_T particles, as well as its dependence on collision energy, is still poorly understood from the RHIC results [2]. Recently, a striking ridge structure has also been observed in very high multiplicity proton-proton (pp) collisions at a center-of-mass energy of 7 TeV at the LHC by the Compact Muon Solenoid (CMS) Collaboration [27], posing new challenges to the understanding of these long-range correlations.

This paper presents the first measurement of dihadron correlations for charged particles produced in the most central (0-5% centrality) PbPb collisions at a nucleon-nucleon center-of-mass energy ($\sqrt{s_{\text{NN}}}$) of 2.76 TeV over a large phase space. The results are presented in terms of the associated hadron yields as a function of pseudorapidity and azimuthal angle relative to trigger particles in different transverse momentum intervals. Traditionally, trigger particles have been utilized to represent the direction of the leading hadron in a jet, and were required to have a higher momentum than all the other associated particles in the jet [2, 6]. However, as shown in ref. [27], important information can also be obtained by studying the correlation of hadron pairs from the same p_{T} interval, which is particularly useful when addressing the properties of the medium itself. The current analysis employs both approaches. This measurement provides a unique examination of the ridge in the most central PbPb collisions at the highest energies reached so far in the laboratory over a wide range in transverse momentum (2-12 GeV/ c) and up to large relative pseudorapidity ($|\Delta\eta| \approx 4$), imposing further quantitative constraints on the possible origin of the ridge.

Details of event readout and analysis for extracting the correlation functions are described in section 2, the physics results found using the correlations are described in section 3, and a summary is given in section 4.

2 Data and analysis

The analysis reported in this paper is based on PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV collected during the LHC heavy-ion run in November and December 2010 with the CMS detector. The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter. Within the field volume are the inner tracker, the crystal electromagnetic calorimeter, and the brass/scintillator hadron calorimeter. Muons are measured in gas-ionization detectors embedded in the steel return yoke. In addition to the barrel and endcap detectors, CMS has extensive forward calorimetry. The nearly 4π solid-angle acceptance of the CMS detector is ideally suited for studies of both short- and long-range particle correlations. A detailed description of the CMS detector can be found in ref. [28]. CMS uses a right-handed coordinate system, with the origin at the nominal interaction point, the x axis pointing to the center of the LHC, the y axis pointing up (perpendicular to the LHC plane), and the z axis along the counterclockwise beam direction. The detector subsystem primarily used for the present analysis is the inner tracker that reconstructs the trajectories of charged particles with $p_{\text{T}} > 100$ MeV/ c , covering the pseudorapidity region $|\eta| < 2.5$, where $\eta = -\ln[\tan(\theta/2)]$ and θ is the polar angle relative to the beam direction. The inner tracker consists of 1440 silicon pixel and 15 148 silicon strip detector modules immersed in the 3.8 T axial magnetic field of the superconducting solenoid.

The event readout of the CMS detector for PbPb collisions is triggered by coincident signals in forward detectors located on both sides of the nominal collision point. In particular, minimum bias PbPb data are recorded based on coincident signals in the beam scintillator counters (BSC, $3.23 < |\eta| < 4.65$) or in the steel/quartz-fiber Cherenkov forward hadron calorimeters (HF, $2.9 < |\eta| < 5.2$) from both ends of the detector. In order to suppress events due to noise, cosmic rays, double-firing triggers, and beam backgrounds,

the minimum bias trigger used in this analysis is required to be in coincidence with bunches colliding in the interaction region. The trigger has an acceptance of $(97 \pm 3)\%$ for hadronic inelastic PbPb collisions [14].

Events are selected offline by requiring in addition at least three hits in the HF calorimeters at both ends of CMS, with at least 3 GeV of energy in each cluster, and the presence of a reconstructed primary vertex containing at least two tracks. These criteria further reduce background from single-beam interactions (e.g., beam gas and beam halo), cosmic muons, and large-impact-parameter, ultra-peripheral collisions that lead to the electromagnetic breakup of one or both of the Pb nuclei [29]. The reconstructed primary vertex is required to be located within 15 cm of the nominal collision point along the beam axis and within a radius of 0.02 cm relative to the average vertex position in the transverse plane.

This analysis is based on a data sample of PbPb collisions corresponding to an integrated luminosity of approximately $3.12 \mu\text{b}^{-1}$ [30, 31], which contains 24.1 million minimum bias collisions after all event selections are applied.

The energy released in the collisions is related to the centrality of heavy-ion interactions, i.e., the geometrical overlap of the incoming nuclei. In CMS, centrality is classified according to percentiles of the distribution of the energy deposited in the HF calorimeters. The centrality class used in this analysis corresponds to the 0-5% most central PbPb collisions, a total of 1.2 million events. More details on the centrality determination can be found in refs. [14, 32].

A reconstructed track is considered as a primary-track candidate if the significance of the separation along the beam axis between the track and the primary vertex, $d_z/\sigma(d_z)$, and the significance of the impact parameter relative to the primary vertex transverse to the beam, $d_{xy}/\sigma(d_{xy})$, are each less than 3. In order to remove tracks with potentially poorly reconstructed momentum values, the relative uncertainty of the momentum measurement, $\sigma(p_T)/p_T$, is required to be less than 5.0%. Requiring at least 12 hits on each track helps to reject misidentified ('fake') tracks. Systematic uncertainties related to the track selections have been evaluated as discussed below.

Trigger particles are defined as all charged particles originating from the primary vertex, with $|\eta| < 2.4$ and in a specified p_T^{trig} range. The number of trigger particles in the event is denoted by N_{trig} , which can be more than one per event. Hadron pairs are formed by associating with every trigger particle the remaining charged particles with $|\eta| < 2.4$ and in a specified p_T^{assoc} range. The per-trigger-particle associated yield distribution is then defined by:

$$\frac{1}{N_{\text{trig}}} \frac{d^2 N^{\text{pair}}}{d\Delta\eta d\Delta\phi} = B(0, 0) \times \frac{S(\Delta\eta, \Delta\phi)}{B(\Delta\eta, \Delta\phi)}, \quad (2.1)$$

where $\Delta\eta$ and $\Delta\phi$ are the differences in η and ϕ of the pair, respectively. The signal distribution, $S(\Delta\eta, \Delta\phi)$, is the measured per-trigger-particle distribution of same-event pairs, i.e.,

$$S(\Delta\eta, \Delta\phi) = \frac{1}{N_{\text{trig}}} \frac{d^2 N^{\text{same}}}{d\Delta\eta d\Delta\phi}. \quad (2.2)$$

The mixed-event background distribution,

$$B(\Delta\eta, \Delta\phi) = \frac{1}{N_{\text{trig}}} \frac{d^2 N^{\text{mix}}}{d\Delta\eta d\Delta\phi}, \quad (2.3)$$

is constructed by pairing the trigger particles in each event with the associated particles from 10 different random events, excluding the original event. The symbol N^{mix} denotes the number of pairs taken from the mixed event. The background distribution is used to account for random combinatorial background and pair-acceptance effects. The normalization factor $B(0,0)$ is the value of $B(\Delta\eta, \Delta\phi)$ at $\Delta\eta = 0$ and $\Delta\phi = 0$ (with a bin width of 0.3 in $\Delta\eta$ and $\pi/16$ in $\Delta\phi$), representing the mixed-event associated yield for both particles of the pair going in approximately the same direction, thus having full pair acceptance. Therefore, the ratio $B(0,0)/B(\Delta\eta, \Delta\phi)$ is the pair-acceptance correction factor used to derive the corrected per-trigger-particle associated yield distribution. Equation (2.1) is calculated in 2 cm wide bins of the vertex position (z_{vtx}) along the beam direction and averaged over the range $|z_{\text{vtx}}| < 15$ cm. To maximize the statistical precision, the absolute values of $\Delta\eta$ and $\Delta\phi$ are used to fill one quadrant of the $(\Delta\eta, \Delta\phi)$ histograms, with the other three quadrants filled (only for illustration purposes) by reflection. Therefore, the resulting distributions are symmetric about $(\Delta\eta, \Delta\phi) = (0, 0)$ by construction.

Each reconstructed track is weighted by the inverse of the efficiency factor, $\varepsilon_{\text{trk}}(\eta, p_{\text{T}})$, as a function of the track's pseudorapidity and transverse momentum. The efficiency weighting factor accounts for the detector acceptance $A(\eta, p_{\text{T}})$, the reconstruction efficiency $E(\eta, p_{\text{T}})$, and the fraction of misidentified tracks, $F(\eta, p_{\text{T}})$,

$$\varepsilon_{\text{trk}}(\eta, p_{\text{T}}) = \frac{AE}{1 - F}. \quad (2.4)$$

Studies with simulated Monte Carlo (MC) events show that the combined geometrical acceptance and reconstruction efficiency for the primary-track reconstruction reaches about 60% for the 0-5% most central PbPb collisions at $p_{\text{T}} > 2$ GeV/ c over the full CMS tracker acceptance ($|\eta| < 2.4$) and 65% for $|\eta| < 1.0$. The fraction of misidentified tracks is about 1-2% for $|\eta| < 1.0$, but increases to 10% at $|\eta| \approx 2.4$. The weighting changes the overall scale but not the shape of the associated yield distribution, which depends on the ratio of the signal to background distributions.

A closure test of the track-weighting procedure is performed on HYDJET [33] (version 1.6) MC events. The efficiency-weighted associated yield distribution from reconstructed tracks is found to agree with the generator-level correlation function to within $\pm 3.3\%$. In addition, systematic checks of the tracking efficiency, in which simulated MC tracks are embedded into data events, give results consistent with pure HYDJET simulations to within $\pm 5.0\%$. The tracking efficiency also depends on the vertex z position of the event. However, this dependence is negligible in this analysis, and its effects are taken into account in the systematic uncertainty by comparing the efficiency-corrected correlation functions for two different z_{vtx} ranges, $|z_{\text{vtx}}| < 15$ cm and $|z_{\text{vtx}}| < 5$ cm, which are found to differ by less than $\pm 2.2\%$. Additional uncertainties due to track quality cuts are examined by loosening or tightening the track selections described previously, and the final results are

Source	Systematic uncertainty of the per-trigger-particle associated yield (%)
Tracking weighting closure test	3.3
Tracking efficiency	5.0
Vertex dependence	2.2
Track selection dependence	2.0
Construction of the mixed-event background	2.9-3.6
Total	7.3-7.6

Table 1. Summary of systematic uncertainties.

found to be insensitive to the selections to within $\pm 2.0\%$. An independent analysis, using a somewhat different but well-established methodology [34, 35] in constructing the mixed-event background is performed as a cross-check, where 10 trigger particles from different events are selected first and combined to form a single event, and then correlated with particles from another event. It yields results within $\pm 2.9\text{-}3.6\%$ of the default values ($\pm 3.6\%$ for $|\Delta\eta| < 1$ and $\pm 2.9\%$ for $2 < |\Delta\eta| < 4$), with a slight dependence on $\Delta\eta$ and $\Delta\phi$. The other four sources of systematic uncertainty are largely independent of $\Delta\eta$ and $\Delta\phi$.

Table 1 summarizes the different systematic sources, whose corresponding uncertainties are added in quadrature becoming the quoted systematic uncertainties of the per-trigger-particle associated yield.

3 Results

The measured per-trigger-particle associated yield distribution of charged hadrons as a function of $|\Delta\eta|$ and $|\Delta\phi|$ in the 0-5% most central PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV is shown in figure 1a for trigger particles with $4 < p_{\text{T}}^{\text{trig}} < 6$ GeV/c and associated particles with $2 < p_{\text{T}}^{\text{assoc}} < 4$ GeV/c. To understand the effects of the hot, dense medium produced in the collisions, this distribution can be compared to that from a PYTHIA8 MC simulation [36] (version 8.135) of pp collisions at $\sqrt{s} = 2.76$ TeV, shown in figure 1b. This transverse momentum range, one of several studied later in this paper, is chosen for this figure since it illustrates the differences between correlations from PbPb data and PYTHIA8 pp MC events most clearly. The main features of the simulated pp MC distribution are a narrow jet-fragmentation peak at $(\Delta\eta, \Delta\phi) \approx (0, 0)$ and a back-to-back jet structure at $|\Delta\phi| = \pi$, but extended in $\Delta\eta$. In the 0-5% most central PbPb collisions, particle correlations are significantly modified, as shown in figure 1a. The away-side pairs ($\Delta\phi \approx \pi$) exhibit a correlation with similar amplitude compared to PYTHIA8, although the structure in PbPb data is much broader in both $\Delta\phi$ and $\Delta\eta$ so that it appears almost flat. On the near side ($\Delta\phi \approx 0$), besides the common jet-like particle production in both pp and PbPb at $(\Delta\eta, \Delta\phi) \approx (0, 0)$ due to jet fragmentation, a clear and significant ridge-like structure is observed in PbPb at $\Delta\phi \approx 0$, which extends all the way to the limit of the measurement of $|\Delta\eta| = 4$.




Figure 1. Two-dimensional (2-D) per-trigger-particle associated yield of charged hadrons as a function of $|\Delta\eta|$ and $|\Delta\phi|$ for $4 < p_T^{\text{trig}} < 6 \text{ GeV}/c$ and $2 < p_T^{\text{assoc}} < 4 \text{ GeV}/c$ from (a) 0-5% most central PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$, and (b) PYTHIA8 pp MC simulation at $\sqrt{s} = 2.76 \text{ TeV}$.

In relativistic heavy-ion collisions, the anisotropic hydrodynamic expansion of the produced medium is one possible source of long-range azimuthal correlations, driven by the event-by-event initial anisotropy of the collision zone [7, 37]. For non-central collisions, these correlations are dominated by the second-order Fourier component of the $|\Delta\phi|$ distribution, usually called elliptic flow or v_2 . Measurements of dihadron correlations at RHIC have frequently attempted to subtract or factorize the elliptic flow contribution based on direct v_2 measurements, in order to reveal other features of particle correlations that may provide insight into the interactions between the jets and the medium. However, recent theoretical developments indicate that the interplay between initial-state fluctuations and the subsequent hydrodynamic expansion gives rise to additional Fourier components in the azimuthal particle correlations [25, 26, 38–42]. These components need to be treated on equal footing with the elliptic flow component. In particular for the 0-5% most central PbPb collisions, the elliptic flow contribution to the azimuthal correlations is not expected to be dominant [43]. Therefore, the original unsubtracted correlation functions are presented in this paper, containing the full information necessary for the comparison with theoretical calculations.

3.1 Associated yield distributions versus $\Delta\phi$

To quantitatively examine the features of short-range and long-range azimuthal correlations, one dimensional (1-D) $\Delta\phi$ correlation functions are calculated by averaging the 2-D distributions over a limited region in $\Delta\eta$ from $\Delta\eta_{\text{min}}$ to $\Delta\eta_{\text{max}}$:

$$\frac{1}{N_{\text{trig}}} \frac{dN^{\text{pair}}}{d\Delta\phi} = \frac{1}{\Delta\eta_{\text{max}} - \Delta\eta_{\text{min}}} \int_{\Delta\eta_{\text{min}}}^{\Delta\eta_{\text{max}}} \frac{1}{N_{\text{trig}}} \frac{d^2N^{\text{pair}}}{d\Delta\eta d\Delta\phi} d\Delta\eta. \quad (3.1)$$




Figure 2. Short-range ($|\Delta\eta| < 1$) per-trigger-particle associated yields of charged hadrons as a function of $|\Delta\phi|$ from the 0-5% most central PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, requiring $2 < p_T^{\text{assoc}} < 4$ GeV/c, for five different intervals of p_T^{trig} . The PYTHIA8 pp MC results (solid histograms) are also shown, shifted up by a constant value to match the PbPb data at $\Delta\phi = 1$ for ease of comparison. The error bars are statistical only and are too small to be visible in most of the panels. The systematic uncertainty of 7.6% for all data points is not shown in the plots.




Figure 3. Long-range ($2 < |\Delta\eta| < 4$) per-trigger-particle associated yields of charged hadrons as a function of $|\Delta\phi|$ under the same conditions as in figure 2. The systematic uncertainty of 7.3% for all data points is not shown in the plots. Dashed lines show the fits by the first five terms in the Fourier series, as discussed in section 3.3.

The results of extracting the 1-D $\Delta\phi$ correlations for the 0-5% most central PbPb collisions are shown in figures 2 and 3. The associated yield per trigger particle in the range of $2 < p_T^{\text{assoc}} < 4$ GeV/c is extracted for five different p_T^{trig} intervals (2-4, 4-6, 6-8, 8-10, and 10-12 GeV/c) and two ranges in $\Delta\eta$. Figure 2 gives the short-range pseudorapidity result, i.e., averaged over the region $|\Delta\eta| < 1$. Figure 3 shows the same comparison for long-range correlations, i.e., averaged over the region $2 < |\Delta\eta| < 4$. A comparison to PYTHIA8 pp MC events at $\sqrt{s} = 2.76$ TeV is also shown, with a constant added to match the PbPb results at $\Delta\phi = 1$ in order to facilitate the comparison. In this projection, only the range $0 < \Delta\phi < \pi$ is shown, as the $\Delta\phi$ correlation function is symmetric around $\Delta\phi = 0$ by construction.

The panels of figures 2 and 3 show a modified shape of the away-side associated yield distribution for $|\Delta\phi| > \pi/2$ in central PbPb collisions that is not present in the PYTHIA8 pp MC simulation. This modification is most pronounced for lower p_T^{trig} values and is characterized by a similarly broad distribution as seen in lower-energy RHIC measurements,

where it was attributed to jet quenching and related phenomena [3, 5]. The near-side associated yield in PbPb data includes the contributions from both the jet-like peak and the ridge structure seen in figure 1a. Thus, it is not directly comparable to PYTHIA8 pp MC events, where the ridge component is absent. A more quantitative comparison of the jet-like component between PbPb data and PYTHIA8 pp MC events will be discussed below.

For azimuthal correlations at large values of $\Delta\eta$ (figure 3), a clear maximum at $\Delta\phi \approx 0$ is observed, which corresponds to the ridge structure seen in the 2-D distribution of figure 1a. This feature of the long-range azimuthal correlation function is not present in the PYTHIA8 pp MC simulation for any p_T^{trig} bin. In PbPb, the height of the ridge structure decreases as p_T^{trig} increases (figure 3) and has largely vanished for $p_T^{\text{trig}} \approx 10\text{--}12\text{ GeV}/c$. The diminishing height of the ridge with increasing p_T^{trig} was not evident from previous measurements at RHIC in AuAu collisions.

3.2 Integrated associated yield

The strengths of the jet peak and ridge on the near side, as well as their dependences on $\Delta\eta$ and p_T^{trig} , can be quantified by the integrated associated yields. In the presence of multiple sources of correlations, the correlation of interest is commonly estimated using an implementation of the zero-yield-at-minimum (ZYAM) method [17]. However, as mentioned previously, the possible contribution of elliptic flow is not taken into account as it is not the dominant effect for the most central PbPb collisions considered here. The ZYAM method is implemented as follows. A second-order polynomial is fitted to the $|\Delta\phi|$ distributions in the region $0.5 < |\Delta\phi| < 1.5$. The location of the minimum of the polynomial in this region is denoted as $\Delta\phi_{\text{ZYAM}}$. Using the position of the minimum, the associated yield is then calculated as the integral of the $|\Delta\phi|$ distribution minus its value at $\Delta\phi_{\text{ZYAM}}$ between $|\Delta\phi| = 0$ and $\Delta\phi_{\text{ZYAM}}$. The uncertainty on the minimum level obtained by the ZYAM procedure, combined with the uncertainty arising from the choice of fit range in $|\Delta\phi|$, results in an uncertainty on the absolute associated yield that is constant with a value of 0.012 over all $\Delta\eta$ and p_T^{trig} bins.

The ZYAM procedure enables the direct extraction of integrated yields of the $\Delta\phi$ -projected distributions in a well-defined manner. Figure 4 shows the resulting near-side associated yield as a function of $|\Delta\eta|$ (in slices of 0.6 units) in both the central PbPb collisions and PYTHIA8 pp MC. The near-side associated yield for PYTHIA8 pp shows a strong peak at $|\Delta\eta| = 0$, which corresponds to the expected correlations within jets. This near-side peak diminishes rapidly with increasing $|\Delta\eta|$. The PbPb data also exhibit a jet-like correlation peak in the yield for small $|\Delta\eta|$, but in contrast, the PbPb data clearly show that the ridge extends to the highest $|\Delta\eta|$ values measured.

Figure 5 presents the integrated associated yield for the jet region ($|\Delta\eta| < 1$) and the ridge region ($2 < |\Delta\eta| < 4$) with $2 < p_T^{\text{assoc}} < 4\text{ GeV}/c$, as a function of p_T^{trig} in the 0-5% most central PbPb collisions. The ridge-region yield is defined as the integral of the near side in long-range $\Delta\phi$ azimuthal correlation functions (figure 3), while the jet-region yield is determined by the difference between the short- and long-range near-side integral, as the ridge is found to be approximately constant in $\Delta\eta$ (figure 4). While the jet-region yield shows an increase with p_T^{trig} due to the increasing jet transverse energy, the ridge-region yield is most prominent for $2 < p_T^{\text{trig}} < 6\text{ GeV}/c$ and tends to drop to almost zero when




Figure 4. Integrated near-side ($|\Delta\phi| < \Delta\phi_{\text{ZYAM}}$) associated yield for $4 < p_{\text{T}}^{\text{trig}} < 6 \text{ GeV}/c$ and $2 < p_{\text{T}}^{\text{assoc}} < 4 \text{ GeV}/c$, above the minimum level found by the ZYAM procedure, as a function of $|\Delta\eta|$ for the 0-5% most central PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$. The error bars correspond to statistical uncertainties, while the brackets denote the systematic uncertainties. The solid line shows the prediction from the PYTHIA8 simulation of pp collisions at $\sqrt{s} = 2.76 \text{ TeV}$.

$p_{\text{T}}^{\text{trig}}$ reaches 10-12 GeV/c. The bars in figure 5 correspond to the statistical uncertainties, while the brackets around the data points denote the systematic uncertainties, which are dominated by the tracking performance, as discussed earlier. Results from the PYTHIA8 pp MC simulation, displayed as solid lines in figure 5, are consistent with zero for all $p_{\text{T}}^{\text{trig}}$ bins in the ridge region and have qualitatively the same trend with $p_{\text{T}}^{\text{trig}}$ in the jet region as the data. It has been seen from previous studies that the ridge is absent in both minimum bias pp data and PYTHIA8 MC simulations. However, the PYTHIA8 program does not fully describe the jet-like correlations observed in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ [27]. A quantitative comparison of the correlations in PbPb and pp collisions will therefore be deferred until pp results at $\sqrt{s} = 2.76 \text{ TeV}$ become available.

3.3 Fourier decomposition of $\Delta\phi$ distributions

Motivated by recent theoretical developments in understanding the long-range ridge effect in the context of higher-order hydrodynamic flow induced by initial geometric fluctuations, such as the “triangular flow” effect [26, 38–42], an alternative way of quantifying the observed long-range correlations is investigated in this paper. The 1-D $\Delta\phi$ -projected distribution for $2 < |\Delta\eta| < 4$ is decomposed in a Fourier series as

$$\frac{1}{N_{\text{trig}}} \frac{dN^{\text{pair}}}{d\Delta\phi} = \frac{N_{\text{assoc}}}{2\pi} \left\{ 1 + \sum_{n=1}^{\infty} 2V_{n\Delta} \cos(n\Delta\phi) \right\}, \quad (3.2)$$




Figure 5. Integrated near-side ($|\Delta\phi| < \Delta\phi_{\text{ZYAM}}$) associated yield above the minimum level found by the ZYAM procedure for (a) the short-range jet region ($|\Delta\eta| < 1$) and (b) the long-range ridge region ($2 < |\Delta\eta| < 4$), as a function of $p_{\text{T}}^{\text{trig}}$ in the 0-5% most central PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV with $2 < p_{\text{T}}^{\text{assoc}} < 4$ GeV/c. The error bars correspond to statistical uncertainties, while the brackets around the data points denote the systematic uncertainties. The solid line shows the prediction from the PYTHIA8 simulation of pp collisions at $\sqrt{s} = 2.76$ TeV.

where N_{assoc} represents the total number of hadron pairs per trigger particle for a given $|\Delta\eta|$ range and $(p_{\text{T}}^{\text{trig}}, p_{\text{T}}^{\text{assoc}})$ bin. The 1-D $\Delta\phi$ projections displayed in figure 3 are fitted by the first five terms in the Fourier series, the resulting fits being shown as the dashed lines in figure 3. The data are well described by the first five terms of the Fourier series. Figure 6 presents the first five Fourier coefficients from the fit as functions of $p_{\text{T}}^{\text{trig}}$ for $2 < p_{\text{T}}^{\text{assoc}} < 4$ GeV/c for the 0-5% most central PbPb collisions and for the PYTHIA8 pp MC simulation. The PYTHIA8 results are scaled by the ratio of $N_{\text{assoc}}^{\text{PYTHIA8}}/N_{\text{assoc}}^{\text{PbPb}}$ in order to remove the trivial effect of the multiplicity dependence. The error bars are statistical only, while the systematic uncertainties are found to be negligible because the Fourier coefficients characterize the overall shape of the correlation functions, and thus are not sensitive to the absolute scale. The coefficients from the fit are also found to be largely independent of each other (correlation coefficients typical below 5%).

All the Fourier coefficients found from fitting the PbPb data show a similar dependence on $p_{\text{T}}^{\text{trig}}$ except for the $V_{1\Delta}$ term, which contains an additional negative contribution that grows toward higher $p_{\text{T}}^{\text{trig}}$. This negative $V_{1\Delta}$ component is consistent with a contribution from momentum conservation or back-to-back dijets [44]. If the observed correlation is purely driven by the single-particle azimuthal anisotropy arising from the hydrodynamic




Figure 6. Fourier coefficients, $V_{1\Delta}$, $V_{2\Delta}$, $V_{3\Delta}$, $V_{4\Delta}$, and $V_{5\Delta}$, extracted as functions of p_T^{trig} for $2 < p_T^{\text{assoc}} < 4 \text{ GeV}/c$ for the 0-5% most central PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$. The error bars represent statistical uncertainties only. The solid lines show the predictions from the PYTHIA8 simulation of pp collisions at $\sqrt{s} = 2.76 \text{ TeV}$.

expansion of the medium [45], the extracted $V_{n\Delta}$ components would be related to the flow coefficients v_n (i.e., v_2 for anisotropic elliptic flow) via

$$V_{n\Delta} \sim v_n^{\text{trig}} \times v_n^{\text{assoc}}, \quad (3.3)$$

where v_n^{trig} and v_n^{assoc} are the flow coefficients for the trigger and associated particles [26]. The flow coefficients, and especially the higher-order terms, are sensitive to the initial conditions and viscosity of the hot, dense medium [38, 46]. This Fourier analysis serves as an alternative way of quantifying the azimuthal correlation functions in heavy-ion collisions and potentially provides new insights in understanding the origin of the ridge effect in these collisions. Analysis of larger data samples using wider ranges of centrality, p_T^{trig} and p_T^{assoc} would provide a direct test of the relation in eq. (3.3) and thus allow a more detailed comparison to theoretical calculations of hydrodynamics, for which the Fourier components provide a concise description of the data.

4 Summary

The CMS detector at the LHC has been used to measure angular correlations between charged particles in $\Delta\eta$ and $\Delta\phi$ up to $|\Delta\eta| \approx 4$ and over the full range of $\Delta\phi$ in the 0-5% most central PbPb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$. This is the first study of long-range azimuthal correlations over a large difference in pseudorapidity in PbPb interactions at the LHC energy. The extracted 2-D associated yield distributions show a variety of characteristic features in heavy-ion collisions that are not present in minimum bias pp interactions. Short- and long-range azimuthal correlations have been studied as a function of the transverse momentum of the trigger particles. The observed long-range ridge-like structure for approximately equal azimuthal angles ($\Delta\phi \approx 0$) is most evident in the intermediate transverse momentum range, $2 < p_T^{\text{trig}} < 6 \text{ GeV}/c$, and decreases to almost zero for p_T^{trig} above 10-12 GeV/c. A qualitatively similar dependence of the ridge on transverse momentum has also been observed in high-multiplicity pp events, indicating a potentially similar physical origin of the effect. A Fourier decomposition of the 1-D $\Delta\phi$ -projected correlation functions

in the ridge region ($2 < |\Delta\eta| < 4$) has been presented. This alternative way of quantifying the correlation data provides valuable information to test a wide range of theoretical models, including recent hydrodynamic calculations of higher-order Fourier components. The very broad solid-angle coverage of the CMS detector and the statistical accuracy of the sample analyzed in this paper provide significantly improved observations of short- and long-range particle correlations over previously available measurements.

Acknowledgments

We wish to congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC machine. We thank the technical and administrative staff at CERN and other CMS institutes, and acknowledge support from: FMSR (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, and FAPESP (Brazil); MES (Bulgaria); CERN; CAS, MoST, and NSFC (China); COLCIENCIAS (Colombia); MSES (Croatia); RPF (Cyprus); Academy of Sciences and NICPB (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRT (Greece); OTKA and NKTH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); NRF and WCU (Korea); LAS (Lithuania); CINVESTAV, CONACYT, SEP, and UASLP-FAI (Mexico); MSI (New Zealand); PAEC (Pakistan); SCSR (Poland); FCT (Portugal); JINR (Armenia, Belarus, Georgia, Ukraine, Uzbekistan); MST and MAE (Russia); MSTD (Serbia); MICINN and CPAN (Spain); Swiss Funding Agencies (Switzerland); NSC (Taipei); TUBITAK and TAEK (Turkey); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie programme and the European Research Council (European Union); the Leventis Foundation; the A. P. Sloan Foundation; the Alexander von Humboldt Foundation; the Associazione per lo Sviluppo Scientifico e Tecnologico del Piemonte (Italy); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); and the Council of Science and Industrial Research, India.

Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] STAR collaboration, C. Adler et al., *Disappearance of back-to-back high p_T hadron correlations in central Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV*, *Phys. Rev. Lett.* **90** (2003) 082302 [[nucl-ex/0210033](#)] [[SPIRES](#)].
- [2] STAR collaboration, B.I. Abelev et al., *Long range rapidity correlations and jet production in high energy nuclear collisions*, *Phys. Rev. C* **80** (2009) 064912 [[arXiv:0909.0191](#)] [[SPIRES](#)].

- [3] STAR collaboration, J. Adams et al., *Distributions of charged hadrons associated with high transverse momentum particles in p p and Au + Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV*, *Phys. Rev. Lett.* **95** (2005) 152301 [[nucl-ex/0501016](#)] [[SPIRES](#)].
- [4] PHENIX collaboration, A. Adare et al., *System size and energy dependence of jet-induced hadron pair correlation shapes in Cu + Cu and Au + Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV and 62.4 GeV*, *Phys. Rev. Lett.* **98** (2007) 232302 [[nucl-ex/0611019](#)] [[SPIRES](#)].
- [5] PHENIX collaboration, A. Adare et al., *Dihadron azimuthal correlations in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV*, *Phys. Rev. C* **78** (2008) 014901 [[arXiv:0801.4545](#)] [[SPIRES](#)].
- [6] PHOBOS collaboration, B. Alver et al., *High transverse momentum triggered correlations over a large pseudorapidity acceptance in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV*, *Phys. Rev. Lett.* **104** (2010) 062301 [[arXiv:0903.2811](#)] [[SPIRES](#)].
- [7] PHOBOS collaboration, B. Alver et al., *System size dependence of cluster properties from two-particle angular correlations in Cu+Cu and Au+Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV*, *Phys. Rev. C* **81** (2010) 024904 [[arXiv:0812.1172](#)] [[SPIRES](#)].
- [8] STAR collaboration, J. Adams et al., *Experimental and theoretical challenges in the search for the quark gluon plasma: the STAR collaboration's critical assessment of the evidence from RHIC collisions*, *Nucl. Phys. A* **757** (2005) 102 [[nucl-ex/0501009](#)] [[SPIRES](#)].
- [9] PHENIX collaboration, K. Adcox et al., *Formation of dense partonic matter in relativistic nucleus nucleus collisions at RHIC: experimental evaluation by the PHENIX collaboration*, *Nucl. Phys. A* **757** (2005) 184 [[nucl-ex/0410003](#)] [[SPIRES](#)].
- [10] B.B. Back et al., *The PHOBOS perspective on discoveries at RHIC*, *Nucl. Phys. A* **757** (2005) 28 [[nucl-ex/0410022](#)] [[SPIRES](#)].
- [11] BRAHMS collaboration, I. Arsene et al., *Quark gluon plasma an color glass condensate at RHIC? The perspective from the BRAHMS experiment*, *Nucl. Phys. A* **757** (2005) 1 [[nucl-ex/0410020](#)] [[SPIRES](#)].
- [12] ALICE collaboration, K. Aamodt et al., *Suppression of charged particle production at large transverse momentum in central Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV*, *Phys. Lett. B* **696** (2011) 30 [[arXiv:1012.1004](#)] [[SPIRES](#)].
- [13] ATLAS collaboration, G. Aad et al., *Observation of a centrality-dependent dijet asymmetry in lead-lead collisions at $\sqrt{(S_{\text{NN}})} = 2.76$ TeV with the ATLAS detector at the LHC*, *Phys. Rev. Lett.* **105** (2010) 252303 [[arXiv:1011.6182](#)] [[SPIRES](#)].
- [14] CMS collaboration, S. Chatrchyan et al., *Observation and studies of jet quenching in Pb-Pb collisions at nucleon-nucleon center-of-mass energy = 2.76 TeV*, submitted to *Phys. Rev. C* (2011).
- [15] N. Armesto, C.A. Salgado and U.A. Wiedemann, *Measuring the collective flow with jets*, *Phys. Rev. Lett.* **93** (2004) 242301 [[hep-ph/0405301](#)] [[SPIRES](#)].
- [16] A. Majumder, B. Müller and S.A. Bass, *Longitudinal broadening of quenched jets in turbulent color fields*, *Phys. Rev. Lett.* **99** (2007) 042301 [[hep-ph/0611135](#)] [[SPIRES](#)].
- [17] C.B. Chiu and R.C. Hwa, *Pedestal and peak structure in jet correlation*, *Phys. Rev. C* **72** (2005) 034903 [[nucl-th/0505014](#)] [[SPIRES](#)].
- [18] C.-Y. Wong, *The momentum kick model description of the near-side ridge and jet quenching*, *Phys. Rev. C* **78** (2008) 064905 [[arXiv:0806.2154](#)] [[SPIRES](#)].

- [19] S.A. Voloshin, *Transverse radial expansion in nuclear collisions and two particle correlations*, *Phys. Lett. B* **632** (2006) 490 [[nucl-th/0312065](#)] [[SPIRES](#)].
- [20] P. Romatschke, *Momentum broadening in an anisotropic plasma*, *Phys. Rev. C* **75** (2007) 014901 [[hep-ph/0607327](#)] [[SPIRES](#)].
- [21] E.V. Shuryak, *On the origin of the ‘ridge’ phenomenon induced by jets in heavy ion collisions*, *Phys. Rev. C* **76** (2007) 047901 [[arXiv:0706.3531](#)] [[SPIRES](#)].
- [22] A. Dumitru, F. Gelis, L. McLerran and R. Venugopalan, *Glasma flux tubes and the near side ridge phenomenon at RHIC*, *Nucl. Phys. A* **810** (2008) 91 [[arXiv:0804.3858](#)] [[SPIRES](#)].
- [23] S. Gavin, L. McLerran and G. Moschelli, *Long range correlations and the soft ridge in relativistic nuclear collisions*, *Phys. Rev. C* **79** (2009) 051902 [[arXiv:0806.4718](#)] [[SPIRES](#)].
- [24] K. Dusling, D. Fernandez-Fraile and R. Venugopalan, *Three-particle correlation from glasma flux tubes*, *Nucl. Phys. A* **828** (2009) 161 [[arXiv:0902.4435](#)].
- [25] Y. Hama, R.P.G. Andrade, F. Grassi and W.-L. Qian, *Trying to understand the ridge effect in hydrodynamic model*, *Nonlin. Phenom. Complex Syst.* **12** (2009) 466 [[arXiv:0911.0811](#)] [[SPIRES](#)].
- [26] B. Alver and G. Roland, *Collision geometry fluctuations and triangular flow in heavy-ion collisions*, *Phys. Rev. C* **81** (2010) 054905 [[arXiv:1003.0194](#)] [[SPIRES](#)].
- [27] CMS collaboration, V. Khachatryan et al., *Observation of long-range near-side angular correlations in proton-proton collisions at the LHC*, *JHEP* **09** (2010) 091 [[arXiv:1009.4122](#)] [[SPIRES](#)].
- [28] CMS collaboration, *The CMS experiment at the CERN LHC*, *JINST* **03** (2008) S08004.
- [29] O. Djuvslund and J. Nystrand, *Single and double photonuclear excitations in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV at the CERN Large Hadron Collider*, *Phys. Rev. C* **83** (2011) 041901 [[arXiv:1011.4908](#)] [[SPIRES](#)].
- [30] CMS collaboration, *Measurement of CMS luminosity*, *CMS Physics Analysis Summary CMS-PAS-EWK-10-004* (2010).
- [31] CMS collaboration, *Absolute luminosity normalization*, *CMS Detector Performance Summary CMS-DP-2011-002* (2011).
- [32] CMS collaboration, D.G. d’Enterria, (Ed.) et al., *CMS physics technical design report: addendum on high density QCD with heavy ions*, *J. Phys. G* **34** (2007) 2307 [[SPIRES](#)].
- [33] I.P. Lokhtin and A.M. Snigirev, *A model of jet quenching in ultrarelativistic heavy ion collisions and high- p_T hadron spectra at RHIC*, *Eur. Phys. J. C* **45** (2006) 211 [[hep-ph/0506189](#)] [[SPIRES](#)].
- [34] STAR collaboration, M.M. Aggarwal et al., *Azimuthal di-hadron correlations in d+Au and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV from STAR*, *Phys. Rev. C* **82** (2010) 024912 [[arXiv:1004.2377](#)] [[SPIRES](#)].
- [35] V. Chetluru and O. Barannikova, *Jet studies for 5.5 TeV Pb+Pb collisions via di-hadron correlations*, *Indian J. Phys.* **84** (2011) 1807.
- [36] T. Sjöstrand, S. Mrenna and P.Z. Skands, *A brief introduction to PYTHIA 8.1*, *Comput. Phys. Commun.* **178** (2008) 852 [[arXiv:0710.3820](#)] [[SPIRES](#)].
- [37] PHOBOS collaboration, B. Alver et al., *System size, energy, pseudorapidity and centrality dependence of elliptic flow*, *Phys. Rev. Lett.* **98** (2007) 242302 [[nucl-ex/0610037](#)] [[SPIRES](#)].

- [38] B.H. Alver, C. Gombeaud, M. Luzum and J.-Y. Ollitrault, *Triangular flow in hydrodynamics and transport theory*, *Phys. Rev. C* **82** (2010) 034913 [[arXiv:1007.5469](#)] [[SPIRES](#)].
- [39] B. Schenke, S. Jeon and C. Gale, *Elliptic and triangular flow in event-by-event (3+1)D viscous hydrodynamics*, *Phys. Rev. Lett.* **106** (2011) 042301 [[arXiv:1009.3244](#)] [[SPIRES](#)].
- [40] H. Petersen, G.-Y. Qin, S.A. Bass and B. Müller, *Triangular flow in event-by-event ideal hydrodynamics in Au+Au collisions at $\sqrt{s_{NN}} = 200A$ GeV*, *Phys. Rev. C* **82** (2010) 041901 [[arXiv:1008.0625](#)] [[SPIRES](#)].
- [41] J. Xu and C.M. Ko, *The effect of triangular flow on di-hadron azimuthal correlations in relativistic heavy ion collisions*, *Phys. Rev. C* **83** (2011) 021903 [[arXiv:1011.3750](#)] [[SPIRES](#)].
- [42] D. Teaney and L. Yan, *Triangularity and dipole asymmetry in heavy ion collisions*, [arXiv:1010.1876](#) [[SPIRES](#)].
- [43] THE ALICE collaboration, K. Aamodt et al., *Elliptic flow of charged particles in Pb-Pb collisions at 2.76 TeV*, [arXiv:1011.3914](#) [[SPIRES](#)].
- [44] M. Luzum and J.-Y. Ollitrault, *Directed flow at midrapidity in heavy-ion collisions*, *Phys. Rev. Lett.* **106** (2011) 102301 [[arXiv:1011.6361](#)] [[SPIRES](#)].
- [45] S. Voloshin and Y. Zhang, *Flow study in relativistic nuclear collisions by Fourier expansion of Azimuthal particle distributions*, *Z. Phys. C* **70** (1996) 665 [[hep-ph/9407282](#)] [[SPIRES](#)].
- [46] P. Staig and E. Shuryak, *The fate of the initial state fluctuations in heavy ion collisions. II the fluctuations and sounds*, [arXiv:1008.3139](#) [[SPIRES](#)].

The CMS collaboration

Yerevan Physics Institute, Yerevan, Armenia

S. Chatrchyan, V. Khachatryan, A.M. Sirunyan, A. Tumasyan

Institut für Hochenergiephysik der OeAW, Wien, Austria

W. Adam, T. Bergauer, M. Dragicevic, J. Erö, C. Fabjan, M. Friedl, R. Frühwirth, V.M. Ghete, J. Hammer¹, S. Häsnel, M. Hoch, N. Hörmann, J. Hrubec, M. Jeitler, W. Kiesenhofer, M. Krammer, D. Liko, I. Mikulec, M. Pernicka, H. Rohringer, R. Schöffbeck, J. Strauss, A. Taurok, F. Teischinger, P. Wagner, W. Waltenberger, G. Walzel, E. Widl, C.-E. Wulz

National Centre for Particle and High Energy Physics, Minsk, Belarus

V. Mossolov, N. Shumeiko, J. Suarez Gonzalez

Universiteit Antwerpen, Antwerpen, Belgium

S. Bansal, L. Benucci, E.A. De Wolf, X. Janssen, J. Maes, T. Maes, L. Mucibello, S. Ochesanu, B. Roland, R. Rougny, M. Selvaggi, H. Van Haevermaet, P. Van Mechelen, N. Van Remortel

Vrije Universiteit Brussel, Brussel, Belgium

F. Blekman, S. Blyweert, J. D'Hondt, O. Devroede, R. Gonzalez Suarez, A. Kalogeropoulos, M. Maes, W. Van Doninck, P. Van Mulders, G.P. Van Onsem, I. Villella

Université Libre de Bruxelles, Bruxelles, Belgium

O. Charaf, B. Clerbaux, G. De Lentdecker, V. Dero, A.P.R. Gay, G.H. Hammad, T. Hreus, P.E. Marage, L. Thomas, C. Vander Velde, P. Vanlaer

Ghent University, Ghent, Belgium

V. Adler, A. Cimmino, S. Costantini, M. Grunewald, B. Klein, J. Lellouch, A. Marinov, J. Mccartin, D. Ryckbosch, F. Thyssen, M. Tytgat, L. Vanelderen, P. Verwilligen, S. Walsh, N. Zaganidis

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

S. Basegmez, G. Bruno, J. Caudron, L. Ceard, E. Cortina Gil, J. De Favereau De Jeneret, C. Delaere¹, D. Favart, A. Giammanco, G. Grégoire, J. Hollar, V. Lemaitre, J. Liao, O. Militaru, S. Oryn, D. Pagano, A. Pin, K. Piotrkowski, N. Schul

Université de Mons, Mons, Belgium

N. Bely, T. Caebergs, E. Daubie

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves, D. De Jesus Damiao, M.E. Pol, M.H.G. Souza

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W. Carvalho, E.M. Da Costa, C. De Oliveira Martins, S. Fonseca De Souza, L. Mundim, H. Nogima, V. Oguri, W.L. Prado Da Silva, A. Santoro, S.M. Silva Do Amaral, A. Sznajder

Instituto de Fisica Teorica, Universidade Estadual Paulista, Sao Paulo, Brazil

C.A. Bernardes², F.A. Dias, T.R. Fernandez Perez Tomei, E. M. Gregores², C. Lagana, F. Marinho, P.G. Mercadante², S.F. Novaes, Sandra S. Padula

Institute for Nuclear Research and Nuclear Energy, Sofia, Bulgaria

N. Darmenov¹, L. Dimitrov, V. Genchev¹, P. Iaydjiev¹, S. Piperov, M. Rodozov, S. Stoykova, G. Sultanov, V. Tcholakov, R. Trayanov, I. Vankov

University of Sofia, Sofia, Bulgaria

A. Dimitrov, R. Hadjiiska, A. Karadzhinova, V. Kozhuharov, L. Litov, M. Mateev, B. Pavlov, P. Petkov

Institute of High Energy Physics, Beijing, China

J.G. Bian, G.M. Chen, H.S. Chen, C.H. Jiang, D. Liang, S. Liang, X. Meng, J. Tao, J. Wang, J. Wang, X. Wang, Z. Wang, H. Xiao, M. Xu, J. Zang, Z. Zhang

State Key Lab. of Nucl. Phys. and Tech., Peking University, Beijing, China

Y. Ban, S. Guo, Y. Guo, W. Li, Y. Mao, S.J. Qian, H. Teng, L. Zhang, B. Zhu, W. Zou

Universidad de Los Andes, Bogota, Colombia

A. Cabrera, B. Gomez Moreno, A.A. Ocampo Rios, A.F. Osorio Oliveros, J.C. Sanabria

Technical University of Split, Split, Croatia

N. Godinovic, D. Lelas, K. Lelas, R. Plestina³, D. Polic, I. Puljak

University of Split, Split, Croatia

Z. Antunovic, M. Dzelalija

Institute Rudjer Boskovic, Zagreb, Croatia

V. Brigljevic, S. Duric, K. Kadija, S. Morovic

University of Cyprus, Nicosia, Cyprus

A. Attikis, M. Galanti, J. Mousa, C. Nicolaou, F. Ptochos, P.A. Razis

Charles University, Prague, Czech Republic

M. Finger, M. Finger Jr.

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

Y. Assran⁴, S. Khalil⁵, M.A. Mahmoud⁶

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

A. Hektor, M. Kadastik, M. Müntel, M. Raidal, L. Rebane

Department of Physics, University of Helsinki, Helsinki, Finland

V. Azzolini, P. Eerola, G. Fedi

Helsinki Institute of Physics, Helsinki, Finland

S. Czellar, J. Härkönen, A. Heikkinen, V. Karimäki, R. Kinnunen, M.J. Kortelainen, T. Lampén, K. Lassila-Perini, S. Lehti, T. Lindén, P. Luukka, T. Mäenpää, E. Tuominen, J. Tuominiemi, E. Tuovinen, D. Ungaro, L. Wendland

Lappeenranta University of Technology, Lappeenranta, Finland

K. Banzuzi, A. Korpela, T. Tuuva

Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France

D. Sillou

DSM/IRFU, CEA/Saclay, Gif-sur-Yvette, France

M. Besancon, S. Choudhury, M. Dejardin, D. Denegri, B. Fabbro, J.L. Faure, F. Ferri, S. Ganjour, F.X. Gentit, A. Givernaud, P. Gras, G. Hamel de Monchenault, P. Jarry, E. Locci, J. Malcles, M. Marionneau, L. Millischer, J. Rander, A. Rosowsky, I. Shreyber, M. Titov, P. Verrecchia

Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France

S. Baffioni, F. Beaudette, L. Benhabib, L. Bianchini, M. Bluj⁷, C. Broutin, P. Busson, C. Charlot, T. Dahms, L. Dobrzynski, S. Elgammal, R. Granier de Cassagnac, M. Haguenaer, P. Miné, C. Mironov, C. Ochando, P. Paganini, D. Sabes, R. Salerno, Y. Sirois, C. Thiebaux, B. Wyslouch⁸, A. Zabi

Institut Pluridisciplinaire Hubert Curien, Université de Strasbourg, Université de Haute Alsace Mulhouse, CNRS/IN2P3, Strasbourg, France

J.-L. Agram⁹, J. Andrea, D. Bloch, D. Bodin, J.-M. Brom, M. Cardaci, E.C. Chabert, C. Collard, E. Conte⁹, F. Drouhin⁹, C. Ferro, J.-C. Fontaine⁹, D. Gelé, U. Goerlach, S. Greder, P. Juillot, M. Karim⁹, A.-C. Le Bihan, Y. Mikami, P. Van Hove

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules (IN2P3), Villeurbanne, France

F. Fassi, D. Mercier

Université de Lyon, Université Claude Bernard Lyon 1, CNRS-IN2P3, Institut de Physique Nucléaire de Lyon, Villeurbanne, France

C. Baty, S. Beauceron, N. Beaupere, M. Bedjidian, O. Bondu, G. Boudoul, D. Boumediene, H. Brun, J. Chasserat, R. Chierici, D. Contardo, P. Depasse, H. El Mamouni, J. Fay, S. Gascon, B. Ille, T. Kurca, T. Le Grand, M. Lethuillier, L. Mirabito, S. Perries, V. Sordini, S. Tosi, Y. Tschudi, P. Verdier

Institute of High Energy Physics and Informatization, Tbilisi State University, Tbilisi, Georgia

D. Lomidze

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

G. Anagnostou, M. Edelhoff, L. Feld, N. Heracleous, O. Hindrichs, R. Jussen, K. Klein, J. Merz, N. Mohr, A. Ostapchuk, A. Perieanu, F. Raupach, J. Sammet, S. Schael, D. Sprenger, H. Weber, M. Weber, B. Wittmer

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

M. Ata, W. Bender, E. Dietz-Laursonn, M. Erdmann, J. Frangenheim, T. Hebbeker, A. Hinzmann, K. Hoepfner, T. Klimkovich, D. Klingebiel, P. Kreuzer, D. Lanske[†], C. Magass, M. Merschmeyer, A. Meyer, P. Papacz, H. Pieta, H. Reithler, S.A. Schmitz, L. Sonnenschein, J. Steggemann, D. Teyssier

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

M. Bontenackels, M. Davids, M. Duda, G. Flügge, H. Geenen, M. Giffels, W. Haj Ahmad, D. Heydhausen, T. Kress, Y. Kuessel, A. Linn, A. Nowack, L. Perchalla, O. Pooth, J. Rennefeld, P. Sauerland, A. Stahl, M. Thomas, D. Tornier, M.H. Zoeller

Deutsches Elektronen-Synchrotron, Hamburg, Germany

M. Aldaya Martin, W. Behrenhoff, U. Behrens, M. Bergholz¹⁰, A. Bethani, K. Borrás, A. Cakir, A. Campbell, E. Castro, D. Dammann, G. Eckerlin, D. Eckstein, A. Flossdorf, G. Flucke, A. Geiser, J. Hauk, H. Jung¹, M. Kasemann, I. Katkov¹¹, P. Katsas, C. Kleinwort, H. Kluge, A. Knutsson, M. Krämer, D. Krücker, E. Kuznetsova, W. Lange, W. Lohmann¹⁰, R. Mankel, M. Marienfeld, I.-A. Melzer-Pellmann, A.B. Meyer, J. Mnich, A. Mussgiller, J. Olzem, D. Pitzl, A. Raspereza, A. Raval, M. Rosin, R. Schmidt¹⁰, T. Schoerner-Sadenius, N. Sen, A. Spiridonov, M. Stein, J. Tomaszewska, R. Walsh, C. Wissing

University of Hamburg, Hamburg, Germany

C. Autermann, V. Blobel, S. Bobrovskiy, J. Draeger, H. Enderle, U. Gebbert, K. Kaschube, G. Kaussen, R. Klanner, J. Lange, B. Mura, S. Naumann-Emme, F. Nowak, N. Pietsch, C. Sander, H. Schettler, P. Schleper, M. Schröder, T. Schum, J. Schwandt, H. Stadie, G. Steinbrück, J. Thomsen

Institut für Experimentelle Kernphysik, Karlsruhe, Germany

C. Barth, J. Bauer, J. Berger, V. Buege, T. Chwalek, W. De Boer, A. Dierlamm, G. Dirkes, M. Feindt, J. Gruschke, C. Hackstein, F. Hartmann, M. Heinrich, H. Held, K.H. Hoffmann, S. Honc, J.R. Komaragiri, T. Kuhr, D. Martschei, S. Mueller, Th. Müller, M. Niegel, O. Oberst, A. Oehler, J. Ott, T. Peiffer, G. Quast, K. Rabbertz, F. Ratnikov, N. Ratnikova, M. Renz, C. Saout, A. Scheurer, P. Schieferdecker, F.-P. Schilling, G. Schott, H.J. Simonis, F.M. Stober, D. Troendle, J. Wagner-Kuhr, T. Weiler, M. Zeise, V. Zhukov¹¹, E.B. Ziebarth

Institute of Nuclear Physics “Demokritos”, Aghia Paraskevi, Greece

G. Daskalakis, T. Geralis, S. Kesisoglou, A. Kyriakis, D. Loukas, I. Manolakos, A. Markou, C. Markou, C. Mavrommatis, E. Ntomari, E. Petrakou

University of Athens, Athens, Greece

L. Gouskos, T.J. Mertzimekis, A. Panagiotou, E. Stiliaris

University of Ioánnina, Ioánnina, Greece

I. Evangelou, C. Foudas, P. Kokkas, N. Manthos, I. Papadopoulos, V. Patras, F.A. Triantis

KFKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary

A. Aranyi, G. Bencze, L. Boldizsar, C. Hajdu¹, P. Hidas, D. Horvath¹², A. Kapusi, K. Krajczar¹³, F. Sikler¹, G.I. Veres¹³, G. Vesztergombi¹³

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

N. Beni, J. Molnar, J. Palinkas, Z. Szillasi, V. Veszpremi

University of Debrecen, Debrecen, Hungary

P. Raics, Z.L. Trocsanyi, B. Ujvari

Panjab University, Chandigarh, India

S.B. Beri, V. Bhatnagar, N. Dhingra, R. Gupta, M. Jindal, M. Kaur, J.M. Kohli, M.Z. Mehta, N. Nishu, L.K. Saini, A. Sharma, A.P. Singh, J.B. Singh, S.P. Singh

University of Delhi, Delhi, India

S. Ahuja, S. Bhattacharya, B.C. Choudhary, B. Gomber, P. Gupta, S. Jain, S. Jain, R. Khurana, A. Kumar, M. Naimuddin, K. Ranjan, R.K. Shivpuri

Bhabha Atomic Research Centre, Mumbai, India

R.K. Choudhury, D. Dutta, S. Kailas, V. Kumar, P. Mehta, A.K. Mohanty¹, L.M. Pant, P. Shukla

Tata Institute of Fundamental Research - EHEP, Mumbai, India

T. Aziz, M. Guchait¹⁴, A. Gurtu, M. Maity¹⁵, D. Majumder, G. Majumder, K. Mazumdar, G.B. Mohanty, A. Saha, K. Sudhakar, N. Wickramage

Tata Institute of Fundamental Research - HECR, Mumbai, India

S. Banerjee, S. Dugad, N.K. Mondal

Institute for Research and Fundamental Sciences (IPM), Tehran, Iran

H. Arfaei, H. Bakhshiansohi¹⁶, S.M. Etesami, A. Fahim¹⁶, M. Hashemi, A. Jafari¹⁶, M. Khakzad, A. Mohammadi¹⁷, M. Mohammadi Najafabadi, S. Paktinat Mehdiabadi, B. Safarzadeh, M. Zeinali¹⁸

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia^{a,b}, L. Barbone^{a,b}, C. Calabria^{a,b}, A. Colaleo^a, D. Creanza^{a,c}, N. De Filippis^{a,c,1}, M. De Palma^{a,b}, L. Fiore^a, G. Iaselli^{a,c}, L. Lusito^{a,b}, G. Maggi^{a,c}, M. Maggi^a, N. Manna^{a,b}, B. Marangelli^{a,b}, S. My^{a,c}, S. Nuzzo^{a,b}, N. Pacifico^{a,b}, G.A. Pierro^a, A. Pompili^{a,b}, G. Pugliese^{a,c}, F. Romano^{a,c}, G. Roselli^{a,b}, G. Selvaggi^{a,b}, L. Silvestris^a, R. Trentadue^a, S. Tupputi^{a,b}, G. Zito^a

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi^a, A.C. Benvenuti^a, D. Bonacorsi^a, S. Braibant-Giacomelli^{a,b}, L. Brigliadori^a, P. Capiluppi^{a,b}, A. Castro^{a,b}, F.R. Cavallo^a, M. Cuffiani^{a,b}, G.M. Dallavalle^a, F. Fabbri^a, A. Fanfani^{a,b}, D. Fasanella^a, P. Giacomelli^a, M. Giunta^a, C. Grandi^a, S. Marcellini^a, G. Masetti^b, M. Meneghelli^{a,b}, A. Montanari^a, F.L. Navarria^{a,b}, F. Odorici^a, A. Perrotta^a, F. Primavera^a, A.M. Rossi^{a,b}, T. Rovelli^{a,b}, G. Siroli^{a,b}, R. Travaglini^{a,b}

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Albergo^{a,b}, G. Cappello^{a,b}, M. Chiorboli^{a,b,1}, S. Costa^{a,b}, A. Tricomi^{a,b}, C. Tuve^a

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

G. Barbagli^a, V. Ciulli^{a,b}, C. Civinini^a, R. D'Alessandro^{a,b}, E. Focardi^{a,b}, S. Frosali^{a,b}, E. Gallo^a, S. Gonzi^{a,b}, P. Lenzi^{a,b}, M. Meschini^a, S. Paoletti^a, G. Sguazzoni^a, A. Tropiano^{a,1}

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi, S. Bianco, S. Colafranceschi¹⁹, F. Fabbri, D. Piccolo

INFN Sezione di Genova, Genova, Italy

P. Fabbriatore, R. Musenich

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^{a,b}, F. De Guio^{a,b,1}, L. Di Matteo^{a,b}, S. Gennai¹, A. Ghezzi^{a,b}, S. Malvezzi^a, A. Martelli^{a,b}, A. Massironi^{a,b}, D. Menasce^a, L. Moroni^a, M. Paganoni^{a,b}, D. Pedrini^a, S. Ragazzi^{a,b}, N. Redaelli^a, S. Sala^a, T. Tabarelli de Fatis^{a,b}

INFN Sezione di Napoli^a, Università di Napoli “Federico II”^b, Napoli, Italy

S. Buontempo^a, C.A. Carrillo Montoya^{a,1}, N. Cavallo^{a,20}, A. De Cosa^{a,b}, F. Fabozzi^{a,20}, A.O.M. Iorio^{a,1}, L. Lista^a, M. Merola^{a,b}, P. Paolucci^a

INFN Sezione di Padova^a, Università di Padova^b, Università di Trento (Trento)^c, Padova, Italy

P. Azzi^a, N. Bacchetta^a, P. Bellan^{a,b}, M. Bellato^a, M. Biasotto^{a,21}, D. Bisello^{a,b}, A. Branca^a, P. Checchia^a, M. De Mattia^{a,b}, T. Dorigo^a, F. Gasparini^{a,b}, F. Gonella^a, A. Gozzelino, M. Gulmini^{a,21}, S. Lacaprara^{a,21}, I. Lazzizzera^{a,c}, M. Margoni^{a,b}, G. Maron^{a,21}, A.T. Meneguzzo^{a,b}, M. Nespolo^{a,1}, M. Passaseo^a, L. Perrozzi^{a,1}, N. Pozzobon^{a,b}, P. Ronchese^{a,b}, F. Simonetto^{a,b}, E. Torassa^a, M. Tosi^{a,b}, A. Triossi^a, S. Vanini^{a,b}

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

P. Baesso^{a,b}, U. Berzano^a, S.P. Ratti^{a,b}, C. Riccardi^{a,b}, P. Torre^{a,b}, P. Vitulo^{a,b}, C. Viviani^{a,b}

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

M. Biasini^{a,b}, G.M. Bilei^a, B. Caponeri^{a,b}, L. Fanò^{a,b}, P. Lariccia^{a,b}, A. Lucaroni^{a,b,1}, G. Mantovani^{a,b}, M. Menichelli^a, A. Nappi^{a,b}, F. Romeo^{a,b}, A. Santocchia^{a,b}, S. Taroni^{a,b,1}, M. Valdata^{a,b}

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy

P. Azzurri^{a,c}, G. Bagliesi^a, J. Bernardini^{a,b}, T. Boccali^{a,1}, G. Broccolo^{a,c}, R. Castaldi^a, R.T. D’Agnolo^{a,c}, R. Dell’Orso^a, F. Fiori^{a,b}, L. Foà^{a,c}, A. Giassi^a, A. Kraan^a, F. Ligabue^{a,c}, T. Lomtadze^a, L. Martini^{a,22}, A. Messineo^{a,b}, F. Palla^a, G. Segneri^a, A.T. Serban^a, P. Spagnolo^a, R. Tenchini^a, G. Tonelli^{a,b,1}, A. Venturi^{a,1}, P.G. Verdini^a

INFN Sezione di Roma^a, Università di Roma “La Sapienza”^b, Roma, Italy

L. Barone^{a,b}, F. Cavallari^a, D. Del Re^{a,b}, E. Di Marco^{a,b}, M. Diemoz^a, D. Franci^{a,b}, M. Grassi^{a,1}, E. Longo^{a,b}, S. Nourbakhsh^a, G. Organtini^{a,b}, F. Pandolfi^{a,b,1}, R. Paramatti^a, S. Rahatlou^{a,b}, C. Rovelli¹

INFN Sezione di Torino^a, Università di Torino^b, Università del Piemonte Orientale (Novara)^c, Torino, Italy

N. Amapane^{a,b}, R. Arcidiacono^{a,c}, S. Argiro^{a,b}, M. Arneodo^{a,c}, C. Biino^a, C. Botta^{a,b,1}, N. Cartiglia^a, R. Castello^{a,b}, M. Costa^{a,b}, N. Demaria^a, A. Graziano^{a,b,1}, C. Mariotti^a, M. Marone^{a,b}, S. Maselli^a, E. Migliore^{a,b}, G. Mila^{a,b}, V. Monaco^{a,b}, M. Musich^{a,b}, M.M. Obertino^{a,c}, N. Pastrone^a, M. Pelliccioni^{a,b}, A. Romero^{a,b}, M. Ruspa^{a,c}, R. Sacchi^{a,b}, V. Sola^{a,b}, A. Solano^{a,b}, A. Staiano^a, A. Vilela Pereira^a

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte^a, F. Cossutti^a, G. Della Ricca^{a,b}, B. Gobbo^a, D. Montanino^{a,b}, A. Penzo^a

Kangwon National University, Chunchon, Korea

S.G. Heo, S.K. Nam

Kyungpook National University, Daegu, Korea

S. Chang, J. Chung, D.H. Kim, G.N. Kim, J.E. Kim, D.J. Kong, H. Park, S.R. Ro, D. Son, D.C. Son, T. Son

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

Zero Kim, J.Y. Kim, S. Song

Korea University, Seoul, Korea

S. Choi, B. Hong, M.S. Jeong, M. Jo, H. Kim, J.H. Kim, T.J. Kim, K.S. Lee, D.H. Moon, S.K. Park, H.B. Rhee, E. Seo, S. Shin, K.S. Sim

University of Seoul, Seoul, Korea

M. Choi, S. Kang, H. Kim, C. Park, I.C. Park, S. Park, G. Ryu

Sungkyunkwan University, Suwon, Korea

Y. Choi, Y.K. Choi, J. Goh, M.S. Kim, E. Kwon, J. Lee, S. Lee, H. Seo, I. Yu

Vilnius University, Vilnius, Lithuania

M.J. Bilinskas, I. Grigelionis, M. Janulis, D. Martisiute, P. Petrov, T. Sabonis

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

H. Castilla-Valdez, E. De La Cruz-Burelo, I. Heredia-de La Cruz, R. Lopez-Fernandez, R. Magaña Villalba, A. Sánchez-Hernández, L.M. Villasenor-Cendejas

Universidad Iberoamericana, Mexico City, Mexico

S. Carrillo Moreno, F. Vazquez Valencia

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

H.A. Salazar Ibarguen

Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico

E. Casimiro Linares, A. Morelos Pineda, M.A. Reyes-Santos

University of Auckland, Auckland, New Zealand

D. Krofcheck, J. Tam

University of Canterbury, Christchurch, New Zealand

P.H. Butler, R. Doesburg, H. Silverwood

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

M. Ahmad, I. Ahmed, M.I. Asghar, H.R. Hoorani, W.A. Khan, T. Khurshid, S. Qazi

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

G. Brona, M. Cwiok, W. Dominik, K. Doroba, A. Kalinowski, M. Konecki, J. Krolikowski

Soltan Institute for Nuclear Studies, Warsaw, Poland

T. Frueboes, R. Gokieli, M. Górski, M. Kazana, K. Nawrocki, K. Romanowska-Rybinska, M. Szleper, G. Wrochna, P. Zalewski

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

N. Almeida, P. Bargassa, A. David, P. Faccioli, P.G. Ferreira Parracho, M. Gallinaro, P. Musella, A. Nayak, P.Q. Ribeiro, J. Seixas, J. Varela

Joint Institute for Nuclear Research, Dubna, Russia

S. Afanasiev, I. Belotelov, P. Bunin, I. Golutvin, A. Kamenev, V. Karjavin, G. Kozlov, A. Lanev, P. Moisenz, V. Palichik, V. Perelygin, S. Shmatov, V. Smirnov, A. Volodko, A. Zarubin

Petersburg Nuclear Physics Institute, Gatchina (St Petersburg), Russia

V. Golovtsov, Y. Ivanov, V. Kim, P. Levchenko, V. Murzin, V. Oreshkin, I. Smirnov, V. Sulimov, L. Uvarov, S. Vavilov, A. Vorobyev, A. Vorobyev

Institute for Nuclear Research, Moscow, Russia

Yu. Andreev, A. Dermenev, S. Gninenko, N. Golubev, M. Kirsanov, N. Krasnikov, V. Matveev, A. Pashenkov, A. Toropin, S. Troitsky

Institute for Theoretical and Experimental Physics, Moscow, Russia

V. Epshteyn, V. Gavrilov, V. Kaftanov[†], M. Kossov¹, A. Krokhotin, N. Lychkovskaya, V. Popov, G. Safronov, S. Semenov, V. Stolin, E. Vlasov, A. Zhokin

Moscow State University, Moscow, Russia

E. Boos, A. Ershov, A. Gribushin, O. Kodolova, V. Korotkikh, I. Lokhtin, A. Markina, S. Obraztsov, M. Perfilov, S. Petrushanko, L. Sarycheva, V. Savrin, A. Snigirev, I. Vardanyan

P.N. Lebedev Physical Institute, Moscow, Russia

V. Andreev, M. Azarkin, I. Dremin, M. Kirakosyan, A. Leonidov, S.V. Rusakov, A. Vinogradov

State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, Russia

I. Azhgirey, S. Bitioukov, V. Grishin¹, V. Kachanov, D. Konstantinov, A. Korablev, V. Krychkine, V. Petrov, R. Ryutin, S. Slabospitsky, A. Sobol, L. Tourtchanovitch, S. Troshin, N. Tyurin, A. Uzunian, A. Volkov

University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia

P. Adzic²³, M. Djordjevic, D. Krpic²³, J. Milosevic

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

M. Aguilar-Benitez, J. Alcaraz Maestre, P. Arce, C. Battilana, E. Calvo, M. Cepeda, M. Cerrada, M. Chamizo Llatas, N. Colino, B. De La Cruz, A. Delgado Peris, C. Diez

Pardos, D. Domínguez Vázquez, C. Fernandez Bedoya, J.P. Fernández Ramos, A. Ferrando, J. Flix, M.C. Fouz, P. Garcia-Abia, O. Gonzalez Lopez, S. Goy Lopez, J.M. Hernandez, M.I. Josa, G. Merino, J. Puerta Pelayo, I. Redondo, L. Romero, J. Santaolalla, M.S. Soares, C. Willmott

Universidad Autónoma de Madrid, Madrid, Spain

C. Albajar, G. Codispoti, J.F. de Trocóniz

Universidad de Oviedo, Oviedo, Spain

J. Cuevas, J. Fernandez Menendez, S. Folgueras, I. Gonzalez Caballero, L. Lloret Iglesias, J.M. Vizan Garcia

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

J.A. Brochero Cifuentes, I.J. Cabrillo, A. Calderon, S.H. Chuang, J. Duarte Campderros, M. Felcini²⁴, M. Fernandez, G. Gomez, J. Gonzalez Sanchez, C. Jorda, P. Lobelle Pardo, A. Lopez Virto, J. Marco, R. Marco, C. Martinez Rivero, F. Matorras, F.J. Munoz Sanchez, J. Piedra Gomez²⁵, T. Rodrigo, A.Y. Rodríguez-Marrero, A. Ruiz-Jimeno, L. Scodellaro, M. Sobron Sanudo, I. Vila, R. Vilar Cortabitarte

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo, E. Auffray, G. Auzinger, P. Baillon, A.H. Ball, D. Barney, A.J. Bell²⁶, D. Benedetti, C. Bernet³, W. Bialas, P. Bloch, A. Bocci, S. Bolognesi, M. Bona, H. Breuker, K. Bunkowski, T. Camporesi, G. Cerminara, J.A. Coarasa Perez, B. Curé, D. D'Enterria, A. De Roeck, S. Di Guida, N. Dupont-Sagorin, A. Elliott-Peisert, B. Frisch, W. Funk, A. Gaddi, G. Georgiou, H. Gerwig, D. Gigi, K. Gill, D. Giordano, F. Glege, R. Gomez-Reino Garrido, M. Gouzevitch, P. Govoni, S. Gowdy, L. Guiducci, M. Hansen, C. Hartl, J. Harvey, J. Hegeman, B. Hegner, H.F. Hoffmann, A. Honma, V. Innocente, P. Janot, K. Kaadze, E. Karavakis, P. Lecoq, C. Lourenço, T. Mäki, M. Malberti, L. Malgeri, M. Mannelli, L. Masetti, A. Maurisset, F. Meijers, S. Mersi, E. Meschi, R. Moser, M.U. Mozer, M. Mulders, E. Nesvold¹, M. Nguyen, T. Orimoto, L. Orsini, E. Perez, A. Petrilli, A. Pfeiffer, M. Pierini, M. Pimiä, D. Piparo, G. Polese, A. Racz, J. Rodrigues Antunes, G. Rolandi²⁷, T. Rommerskirchen, M. Rovere, H. Sakulin, C. Schäfer, C. Schwick, I. Segoni, A. Sharma, P. Siegrist, M. Simon, P. Sphicas²⁸, M. Spiropulu²⁹, M. Stoye, M. Tadel, P. Tropea, A. Tsiros, P. Vichoudis, M. Voutilainen, W.D. Zeuner

Paul Scherrer Institut, Villigen, Switzerland

W. Bertl, K. Deiters, W. Erdmann, K. Gabathuler, R. Horisberger, Q. Ingram, H.C. Kaestli, S. König, D. Kotlinski, U. Langenegger, F. Meier, D. Renker, T. Rohe, J. Sibille³⁰, A. Starodumov³¹

Institute for Particle Physics, ETH Zurich, Zurich, Switzerland

P. Bortignon, L. Caminada³², N. Chanon, Z. Chen, S. Cittolin, G. Dissertori, M. Dittmar, J. Eugster, K. Freudenreich, C. Grab, W. Hintz, P. Lecomte, W. Luster, C. Marchica³², P. Martinez Ruiz del Arbol, P. Meridiani, P. Milenovic³³, F. Moortgat, C. Nägeli³², P. Nef, F. Nessi-Tedaldi, L. Pape, F. Pauss, T. Punz, A. Rizzi, F.J. Ronga,

M. Rossini, L. Sala, A.K. Sanchez, M.-C. Sawley, B. Stieger, L. Tauscher[†], A. Thea, K. Theofilatos, D. Treille, C. Urscheler, R. Wallny, M. Weber, L. Wehrli, J. Weng

Universität Zürich, Zurich, Switzerland

E. Aguiló, C. Amsler, V. Chiochia, S. De Visscher, C. Favaro, M. Ivova Rikova, B. Millan Mejias, P. Otiougova, C. Regenfus, P. Robmann, A. Schmidt, H. Snoek

National Central University, Chung-Li, Taiwan

Y.H. Chang, K.H. Chen, S. Dutta, C.M. Kuo, S.W. Li, W. Lin, Z.K. Liu, Y.J. Lu, D. Mekterovic, R. Volpe, J.H. Wu, S.S. Yu

National Taiwan University (NTU), Taipei, Taiwan

P. Bartalini, P. Chang, Y.H. Chang, Y.W. Chang, Y. Chao, K.F. Chen, W.-S. Hou, Y. Hsiung, K.Y. Kao, Y.J. Lei, R.-S. Lu, J.G. Shiu, Y.M. Tzeng, M. Wang

Cukurova University, Adana, Turkey

A. Adiguzel, M.N. Bakirci³⁴, S. Cerci³⁵, C. Dozen, I. Dumanoglu, A. Ekenel, E. Eskut, S. Girgis, G. Gokbulut, I. Hos, E.E. Kangal, A. Kayis Topaksu, G. Onengut, K. Ozdemir, S. Ozturk, A. Polatoz, K. Sogut³⁶, D. Sunar Cerci³⁵, B. Tali³⁵, H. Topakli³⁴, D. Uzun, L.N. Vergili, M. Vergili, S. Yilmaz

Middle East Technical University, Physics Department, Ankara, Turkey

I.V. Akin, T. Aliev, S. Bilmis, M. Deniz, H. Gamsizkan, A.M. Guler, K. Ocalan, A. Ozpineci, M. Serin, R. Sever, U.E. Surat, E. Yildirim, M. Zeyrek

Bogazici University, Istanbul, Turkey

M. Delimeroglu, D. Demir³⁷, E. Gülmez, B. Isildak, M. Kaya³⁸, O. Kaya³⁸, S. Ozkorucuklu³⁹, N. Sonmez⁴⁰

National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine

L. Levchuk

University of Bristol, Bristol, United Kingdom

F. Bostock, J.J. Brooke, T.L. Cheng, E. Clement, D. Cussans, R. Frazier, J. Goldstein, M. Grimes, M. Hansen, D. Hartley, G.P. Heath, H.F. Heath, L. Kreczko, S. Metson, D.M. Newbold⁴¹, K. Nirunpong, A. Poll, S. Senkin, V.J. Smith, S. Ward

Rutherford Appleton Laboratory, Didcot, United Kingdom

L. Basso⁴², A. Belyaev⁴², C. Brew, R.M. Brown, B. Camanzi, D.J.A. Cockerill, J.A. Coughlan, K. Harder, S. Harper, J. Jackson, B.W. Kennedy, E. Olaiya, D. Petyt, B.C. Radburn-Smith, C.H. Shepherd-Themistocleous, I.R. Tomalin, W.J. Womersley, S.D. Worm

Imperial College, London, United Kingdom

R. Bainbridge, G. Ball, J. Ballin, R. Beuselinck, O. Buchmuller, D. Colling, N. Cripps, M. Cutajar, G. Davies, M. Della Negra, W. Ferguson, J. Fulcher, D. Futyan, A. Gilbert, A. Guneratne Bryer, G. Hall, Z. Hatherell, J. Hays, G. Iles, M. Jarvis, G. Karapostoli,

L. Lyons, B.C. MacEvoy, A.-M. Magnan, J. Marrouche, B. Mathias, R. Nandi, J. Nash, A. Nikitenko³¹, A. Papageorgiou, M. Pesaresi, K. Petridis, M. Pioppi⁴³, D.M. Raymond, S. Rogerson, N. Rompotis, A. Rose, M.J. Ryan, C. Seez, P. Sharp, A. Sparrow, A. Tapper, S. Tourneur, M. Vazquez Acosta, T. Virdee, S. Wakefield, N. Wardle, D. Wardrope, T. Whyntie

Brunel University, Uxbridge, United Kingdom

M. Barrett, M. Chadwick, J.E. Cole, P.R. Hobson, A. Khan, P. Kyberd, D. Leslie, W. Martin, I.D. Reid, L. Teodorescu

Baylor University, Waco, USA

K. Hatakeyama, H. Liu

Boston University, Boston, USA

T. Bose, E. Carrera Jarrin, C. Fantasia, A. Heister, J. St. John, P. Lawson, D. Lazic, J. Rohlf, D. Sperka, L. Sulak

Brown University, Providence, USA

A. Avetisyan, S. Bhattacharya, J.P. Chou, D. Cutts, A. Ferapontov, U. Heintz, S. Jabeen, G. Kukartsev, G. Landsberg, M. Luk, M. Narain, D. Nguyen, M. Segala, T. Sinthuprasith, T. Speer, K.V. Tsang

University of California, Davis, Davis, USA

R. Breedon, M. Calderon De La Barca Sanchez, S. Chauhan, M. Chertok, J. Conway, P.T. Cox, J. Dolen, R. Erbacher, E. Friis, W. Ko, A. Kopecky, R. Lander, H. Liu, S. Maruyama, T. Miceli, M. Nikolic, D. Pellett, J. Robles, S. Salur, T. Schwarz, M. Searle, J. Smith, M. Squires, M. Tripathi, R. Vasquez Sierra, C. Veelken

University of California, Los Angeles, Los Angeles, USA

V. Andreev, K. Arisaka, D. Cline, R. Cousins, A. Deisher, J. Duris, S. Erhan, C. Farrell, J. Hauser, M. Ignatenko, C. Jarvis, C. Plager, G. Rakness, P. Schlein[†], J. Tucker, V. Valuev

University of California, Riverside, Riverside, USA

J. Babb, A. Chandra, R. Clare, J. Ellison, J.W. Gary, F. Giordano, G. Hanson, G.Y. Jeng, S.C. Kao, F. Liu, H. Liu, O.R. Long, A. Luthra, H. Nguyen, B.C. Shen[†], R. Stringer, J. Sturdy, S. Sumowidagdo, R. Wilken, S. Wimpenny

University of California, San Diego, La Jolla, USA

W. Andrews, J.G. Branson, G.B. Cerati, D. Evans, F. Golf, A. Holzner, R. Kelley, M. Lebourgeois, J. Letts, B. Mangano, S. Padhi, C. Palmer, G. Petrucciani, H. Pi, M. Pieri, R. Ranieri, M. Sani, V. Sharma, S. Simon, E. Sudano, Y. Tu, A. Vartak, S. Wasserbaech⁴⁴, F. Würthwein, A. Yagil, J. Yoo

University of California, Santa Barbara, Santa Barbara, USA

D. Barge, R. Bellan, C. Campagnari, M. D'Alfonso, T. Danielson, K. Flowers, P. Geffert, J. Incandela, C. Justus, P. Kalavase, S.A. Koay, D. Kovalskyi, V. Krutelyov, S. Lowette, N. Mccoll, V. Pavlunin, F. Rebassoo, J. Ribnik, J. Richman, R. Rossin, D. Stuart, W. To, J.R. Vlimant

California Institute of Technology, Pasadena, USA

A. Apresyan, A. Bornheim, J. Bunn, Y. Chen, M. Gataullin, Y. Ma, A. Mott, H.B. Newman, C. Rogan, K. Shin, V. Timciuc, P. Traczyk, J. Veverka, R. Wilkinson, Y. Yang, R.Y. Zhu

Carnegie Mellon University, Pittsburgh, USA

B. Akgun, R. Carroll, T. Ferguson, Y. Iiyama, D.W. Jang, S.Y. Jun, Y.F. Liu, M. Paulini, J. Russ, H. Vogel, I. Vorobiev

University of Colorado at Boulder, Boulder, USA

J.P. Cumalat, M.E. Dinardo, B.R. Drell, C.J. Edelmaier, W.T. Ford, A. Gaz, B. Heyburn, E. Luiggi Lopez, U. Nauenberg, J.G. Smith, K. Stenson, K.A. Ulmer, S.R. Wagner, S.L. Zang

Cornell University, Ithaca, USA

L. Agostino, J. Alexander, D. Cassel, A. Chatterjee, S. Das, N. Eggert, L.K. Gibbons, B. Heltsley, W. Hopkins, A. Khukhunaishvili, B. Kreis, G. Nicolas Kaufman, J.R. Patterson, D. Puigh, A. Ryd, E. Salvati, X. Shi, W. Sun, W.D. Teo, J. Thom, J. Thompson, J. Vaughan, Y. Weng, L. Winstrom, P. Wittich

Fairfield University, Fairfield, USA

A. Biselli, G. Cirino, D. Winn

Fermi National Accelerator Laboratory, Batavia, USA

S. Abdullin, M. Albrow, J. Anderson, G. Apollinari, M. Atac, J.A. Bakken, S. Banerjee, L.A.T. Bauerdick, A. Beretvas, J. Berryhill, P.C. Bhat, I. Bloch, F. Borchering, K. Burkett, J.N. Butler, V. Chetluru, H.W.K. Cheung, F. Chlebana, S. Cihangir, W. Cooper, D.P. Eartly, V.D. Elvira, S. Esen, I. Fisk, J. Freeman, Y. Gao, E. Gottschalk, D. Green, K. Gunthoti, O. Gutsche, J. Hanlon, R.M. Harris, J. Hirschauer, B. Hooberman, H. Jensen, M. Johnson, U. Joshi, R. Khatiwada, B. Klima, K. Kousouris, S. Kunori, S. Kwan, C. Leonidopoulos, P. Limon, D. Lincoln, R. Lipton, J. Lykken, K. Maeshima, J.M. Marraffino, D. Mason, P. McBride, T. Miao, K. Mishra, S. Mrenna, Y. Musienko⁴⁵, C. Newman-Holmes, V. O'Dell, R. Pordes, O. Prokofyev, N. Saoulidou, E. Sexton-Kennedy, S. Sharma, W.J. Spalding, L. Spiegel, P. Tan, L. Taylor, S. Tkaczyk, L. Uplegger, E.W. Vaandering, R. Vidal, J. Whitmore, W. Wu, F. Yang, F. Yumiceva, J.C. Yun

University of Florida, Gainesville, USA

D. Acosta, P. Avery, D. Bourilkov, M. Chen, M. De Gruttola, G.P. Di Giovanni, D. Dobur, A. Drozdetskiy, R.D. Field, M. Fisher, Y. Fu, I.K. Furic, J. Gartner, B. Kim, J. Konigsberg, A. Korytov, A. Kropivnitskaya, T. Kypreos, K. Matchev, G. Mitselmakher, L. Muniz, C. Prescott, R. Remington, M. Schmitt, B. Scurlock, P. Sellers, N. Skhirtladze, M. Snowball, D. Wang, J. Yelton, M. Zakaria

Florida International University, Miami, USA

C. Ceron, V. Gaultney, L. Kramer, L.M. Lebolo, S. Linn, P. Markowitz, G. Martinez, D. Mesa, J.L. Rodriguez

Florida State University, Tallahassee, USA

T. Adams, A. Askew, J. Bochenek, J. Chen, B. Diamond, S.V. Gleyzer, J. Haas, S. Hagopian, V. Hagopian, M. Jenkins, K.F. Johnson, H. Prosper, L. Quertenmont, S. Sekmen, V. Veeraraghavan

Florida Institute of Technology, Melbourne, USA

M.M. Baarmand, B. Dorney, S. Guragain, M. Hohlmann, H. Kalakhety, R. Ralich, I. Vodopiyanov

University of Illinois at Chicago (UIC), Chicago, USA

M.R. Adams, I.M. Anghel, L. Apanasevich, Y. Bai, V.E. Bazterra, R.R. Betts, J. Callner, R. Cavanaugh, C. Dragoiu, L. Gauthier, C.E. Gerber, S. Hamdan, D.J. Hofman, S. Khalatyan, G.J. Kunde⁴⁶, F. Lacroix, M. Malek, C. O'Brien, C. Silvestre, A. Smoron, D. Strom, N. Varelas

The University of Iowa, Iowa City, USA

U. Akgun, E.A. Albayrak, B. Bilki, W. Clarida, F. Duru, C.K. Lae, E. McCliment, J.-P. Merlo, H. Mermerkaya⁴⁷, A. Mestvirishvili, A. Moeller, J. Nachtman, C.R. Newsom, E. Norbeck, J. Olson, Y. Onel, F. Ozok, S. Sen, J. Wetzel, T. Yetkin, K. Yi

Johns Hopkins University, Baltimore, USA

B.A. Barnett, B. Blumenfeld, A. Bonato, C. Eskew, D. Fehling, G. Giurgiu, A.V. Gritsan, Z.J. Guo, G. Hu, P. Maksimovic, S. Rappoccio, M. Swartz, N.V. Tran, A. Whitbeck

The University of Kansas, Lawrence, USA

P. Baringer, A. Bean, G. Benelli, O. Grachov, R.P. Kenny Iii, M. Murray, D. Noonan, S. Sanders, J.S. Wood, V. Zhukova

Kansas State University, Manhattan, USA

A.f. Barfuss, T. Bolton, I. Chakaberia, A. Ivanov, S. Khalil, M. Makouski, Y. Maravin, S. Shrestha, I. Svintradze, Z. Wan

Lawrence Livermore National Laboratory, Livermore, USA

J. Gronberg, D. Lange, D. Wright

University of Maryland, College Park, USA

A. Baden, M. Boutemour, S.C. Eno, D. Ferencek, J.A. Gomez, N.J. Hadley, R.G. Kellogg, M. Kirn, Y. Lu, A.C. Mignerey, K. Rossato, P. Rumerio, F. Santanastasio, A. Skuja, J. Temple, M.B. Tonjes, S.C. Tonwar, E. Twedt

Massachusetts Institute of Technology, Cambridge, USA

B. Alver, G. Bauer, J. Bendavid, W. Busza, E. Butz, I.A. Cali, M. Chan, V. Dutta, P. Everaerts, G. Gomez Ceballos, M. Goncharov, K.A. Hahn, P. Harris, Y. Kim, M. Klute, Y.-J. Lee, W. Li, C. Loizides, P.D. Luckey, T. Ma, S. Nahn, C. Paus, D. Ralph, C. Roland, G. Roland, M. Rudolph, G.S.F. Stephans, F. Stöckli, K. Sumorok, K. Sung, E.A. Wenger, S. Xie, M. Yang, Y. Yilmaz, A.S. Yoon, M. Zanetti

University of Minnesota, Minneapolis, USA

S.I. Cooper, P. Cushman, B. Dahmes, A. De Benedetti, P.R. Duderø, G. Franzoni, J. Haupt, K. Klapoetke, Y. Kubota, J. Mans, V. Rekovic, R. Rusack, M. Sasseville, A. Singovsky

University of Mississippi, University, USA

L.M. Cremaldi, R. Godang, R. Kroeger, L. Perera, R. Rahmat, D.A. Sanders, D. Summers

University of Nebraska-Lincoln, Lincoln, USA

K. Bloom, S. Bose, J. Butt, D.R. Claes, A. Dominguez, M. Eads, J. Keller, T. Kelly, I. Kravchenko, J. Lazo-Flores, H. Malbouisson, S. Malik, G.R. Snow

State University of New York at Buffalo, Buffalo, USA

U. Baur, A. Godshalk, I. Iashvili, S. Jain, A. Kharchilava, A. Kumar, S.P. Shipkowski, K. Smith

Northeastern University, Boston, USA

G. Alverson, E. Barberis, D. Baumgartel, O. Boeriu, M. Chasco, S. Reucroft, J. Swain, D. Trocino, D. Wood, J. Zhang

Northwestern University, Evanston, USA

A. Anastassov, A. Kubik, N. Odell, R.A. Ofierzynski, B. Pollack, A. Pozdnyakov, M. Schmitt, S. Stoynev, M. Velasco, S. Won

University of Notre Dame, Notre Dame, USA

L. Antonelli, D. Berry, M. Hildreth, C. Jessop, D.J. Karmgard, J. Kolb, T. Kolberg, K. Lannon, W. Luo, S. Lynch, N. Marinelli, D.M. Morse, T. Pearson, R. Ruchti, J. Slaunwhite, N. Valls, M. Wayne, J. Ziegler

The Ohio State University, Columbus, USA

B. Bylsma, L.S. Durkin, J. Gu, C. Hill, P. Killewald, K. Kotov, T.Y. Ling, M. Rodenburg, G. Williams

Princeton University, Princeton, USA

N. Adam, E. Berry, P. Elmer, D. Gerbaudo, V. Halyo, P. Hebda, A. Hunt, J. Jones, E. Laird, D. Lopes Pegna, D. Marlow, T. Medvedeva, M. Mooney, J. Olsen, P. Piroué, X. Quan, H. Saka, D. Stickland, C. Tully, J.S. Werner, A. Zuranski

University of Puerto Rico, Mayaguez, USA

J.G. Acosta, X.T. Huang, A. Lopez, H. Mendez, S. Oliveros, J.E. Ramirez Vargas, A. Zatserklyaniy

Purdue University, West Lafayette, USA

E. Alagoz, V.E. Barnes, G. Bolla, L. Borrello, D. Bortoletto, A. Everett, A.F. Garfinkel, L. Gutay, Z. Hu, M. Jones, O. Koybasi, M. Kress, A.T. Laasanen, N. Leonardo, C. Liu, V. Maroussov, P. Merkel, D.H. Miller, N. Neumeister, I. Shipsey, D. Silvers, A. Svyatkovskiy, H.D. Yoo, J. Zablocki, Y. Zheng

Purdue University Calumet, Hammond, USA

P. Jindal, N. Parashar

Rice University, Houston, USA

C. Boulahouache, V. Cuplov, K.M. Ecklund, F.J.M. Geurts, B.P. Padley, R. Redjimi, J. Roberts, J. Zabel

University of Rochester, Rochester, USA

B. Betchart, A. Bodek, Y.S. Chung, R. Covarelli, P. de Barbaro, R. Demina, Y. Eshaq, H. Flacher, A. Garcia-Bellido, P. Goldenzweig, Y. Gotra, J. Han, A. Harel, D.C. Miner, D. Orbaker, G. Petrillo, D. Vishnevskiy, M. Zielinski

The Rockefeller University, New York, USA

A. Bhatti, R. Ciesielski, L. Demortier, K. Goulios, G. Lungu, S. Malik, C. Mesropian, M. Yan

Rutgers, the State University of New Jersey, Piscataway, USA

O. Atramentov, A. Barker, D. Duggan, Y. Gershtein, R. Gray, E. Halkiadakis, D. Hidas, D. Hits, A. Lath, S. Panwalkar, R. Patel, A. Richards, K. Rose, S. Schnetzer, S. Somalwar, R. Stone, S. Thomas

University of Tennessee, Knoxville, USA

G. Cerizza, M. Hollingsworth, S. Spanier, Z.C. Yang, A. York

Texas A&M University, College Station, USA

R. Eusebi, J. Gilmore, A. Gurrola, T. Kamon, V. Khotilovich, R. Montalvo, I. Osipenkov, Y. Pakhotin, J. Pivarski, A. Safonov, S. Sengupta, A. Tatarinov, D. Toback, M. Weinberger

Texas Tech University, Lubbock, USA

N. Akchurin, C. Bardak, J. Damgov, C. Jeong, K. Kovitangoon, S.W. Lee, P. Mane, Y. Roh, A. Sill, I. Volobouev, R. Wigmans, E. Yazgan

Vanderbilt University, Nashville, USA

E. Appelt, E. Brownson, D. Engh, C. Florez, W. Gabella, M. Issah, W. Johns, P. Kurt, C. Maguire, A. Melo, P. Sheldon, B. Snook, S. Tuo, J. Velkovska

University of Virginia, Charlottesville, USA

M.W. Arenton, M. Balazs, S. Boutle, B. Cox, B. Francis, R. Hirosky, A. Ledovskoy, C. Lin, C. Neu, R. Yohay

Wayne State University, Detroit, USA

S. Gollapinni, R. Harr, P.E. Karchin, P. Lamichhane, M. Mattson, C. Milstène, A. Sakharov

University of Wisconsin, Madison, USA

M. Anderson, M. Bachtis, J.N. Bellinger, D. Carlsmith, S. Dasu, J. Efron, K. Flood, L. Gray, K.S. Grogg, M. Grothe, R. Hall-Wilton, M. Herndon, A. Hervé, P. Klabbers, J. Klukas, A. Lanaro, C. Lazaridis, J. Leonard, R. Loveless, A. Mohapatra, F. Palmonari, D. Reeder, I. Ross, A. Savin, W.H. Smith, J. Swanson, M. Weinberg

- †: Deceased
- 1: Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
 - 2: Also at Universidade Federal do ABC, Santo Andre, Brazil
 - 3: Also at Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France
 - 4: Also at Suez Canal University, Suez, Egypt
 - 5: Also at British University, Cairo, Egypt
 - 6: Also at Fayoum University, El-Fayoum, Egypt
 - 7: Also at Soltan Institute for Nuclear Studies, Warsaw, Poland
 - 8: Also at Massachusetts Institute of Technology, Cambridge, USA
 - 9: Also at Université de Haute-Alsace, Mulhouse, France
 - 10: Also at Brandenburg University of Technology, Cottbus, Germany
 - 11: Also at Moscow State University, Moscow, Russia
 - 12: Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
 - 13: Also at Eötvös Loránd University, Budapest, Hungary
 - 14: Also at Tata Institute of Fundamental Research - HECR, Mumbai, India
 - 15: Also at University of Visva-Bharati, Santiniketan, India
 - 16: Also at Sharif University of Technology, Tehran, Iran
 - 17: Also at Shiraz University, Shiraz, Iran
 - 18: Also at Isfahan University of Technology, Isfahan, Iran
 - 19: Also at Facoltà Ingegneria Università di Roma "La Sapienza", Roma, Italy
 - 20: Also at Università della Basilicata, Potenza, Italy
 - 21: Also at Laboratori Nazionali di Legnaro dell' INFN, Legnaro, Italy
 - 22: Also at Università degli studi di Siena, Siena, Italy
 - 23: Also at Faculty of Physics of University of Belgrade, Belgrade, Serbia
 - 24: Also at University of California, Los Angeles, Los Angeles, USA
 - 25: Also at University of Florida, Gainesville, USA
 - 26: Also at Université de Genève, Geneva, Switzerland
 - 27: Also at Scuola Normale e Sezione dell' INFN, Pisa, Italy
 - 28: Also at University of Athens, Athens, Greece
 - 29: Also at California Institute of Technology, Pasadena, USA
 - 30: Also at The University of Kansas, Lawrence, USA
 - 31: Also at Institute for Theoretical and Experimental Physics, Moscow, Russia
 - 32: Also at Paul Scherrer Institut, Villigen, Switzerland
 - 33: Also at University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia
 - 34: Also at Gaziosmanpasa University, Tokat, Turkey
 - 35: Also at Adiyaman University, Adiyaman, Turkey
 - 36: Also at Mersin University, Mersin, Turkey
 - 37: Also at Izmir Institute of Technology, Izmir, Turkey
 - 38: Also at Kafkas University, Kars, Turkey
 - 39: Also at Suleyman Demirel University, Isparta, Turkey
 - 40: Also at Ege University, Izmir, Turkey
 - 41: Also at Rutherford Appleton Laboratory, Didcot, United Kingdom
 - 42: Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
 - 43: Also at INFN Sezione di Perugia; Università di Perugia, Perugia, Italy
 - 44: Also at Utah Valley University, Orem, USA
 - 45: Also at Institute for Nuclear Research, Moscow, Russia
 - 46: Also at Los Alamos National Laboratory, Los Alamos, USA
 - 47: Also at Erzincan University, Erzincan, Turkey